

Acta Universitatis Sapientiae

Mathematica

Volume 12, Number 2, 2020

Sapientia Hungarian University of Transylvania
Scientia Publishing House

Acta Universitatis Sapientiae, Mathematica is covered by the following databases:

Baidu Scholar
CNKI Scholar (China National Knowledge Infrastructure)
CNPIEC – cnpLINKer
Dimensions
DOAJ (Directory of Open Access Journals)
EBSCO (relevant databases)
EBSCO Discovery Service
Genamics JournalSeek
Google Scholar
Japan Science and Technology Agency (JST)
J-Gate
JournalGuide
JournalTOCs
KESLI-NDSL (Korean National Discovery for Science Leaders)
Mathematical Reviews (MathSciNet)
Microsoft Academic
MyScienceWork
Naviga (Softweco)
Primo Central (ExLibris)
ProQuest (relevant databases)
Publons
QOAM (Quality Open Access Market)
ReadCube
SCImago (SJR)
SCOPUS
Semantic Scholar
Summon (ProQuest)
TDNet
Ulrich's Periodicals Directory/ulrichsweb
WanFang Data
Web of Science – Emerging Sources Citation Index
WorldCat (OCLC)
Zentralblatt Math (zbMATH)

Contents

<i>A. H. Ansari, M. S. Khan, V. Rakočević</i> Maia type fixed point results via C-class function	227
<i>P. Baliarsingh, L. Nayak, S. Samantaray</i> On the convergence difference sequences and the related operator norms	245
<i>D. Bród</i> On some properties of split Horadam quaternions	260
<i>J. Bukor, J. T. Tóth</i> On topological properties of the set of maldistributed sequences	272
<i>C. M. da Fonseca</i> On the connection between tridiagonal matrices, Chebyshev polynomials, and Fibonacci numbers	280
<i>P. V. Danchev</i> A note on nil-clean rings	287
<i>H. Ö. Güney, S. Owa</i> Generalized operator for Alexander integral operator	294
<i>S. F. Namiq, E. Rosas</i> On $\lambda^D - R_0$ and $\lambda^D - R_1$ spaces	307

<i>Ş. Özlü, A. Sezgin</i> Soft covered ideals in semigroups	317
<i>M. Shahraki, S. Sedghi, S. M. A. Aleomraninejad, Z. D. Mitrović</i> Some fixed point results on S-metric spaces	347
<i>V. Soleymanivarniab, R. Nikandish, A. Tehranian</i> On the metric dimension of strongly annihilating-ideal graphs of commutative rings	358
<i>V. Agnes S. J. Lavanya, M. P. Jeyaraman, H. Aaisha Farzana</i> On application of differential subordination for Carathéodory functions	370
<i>A. Rawshdeh, H. H. Al-Jarrah, K. Y. Al- Zoubi, W. A. Shatanawi</i> On γ -countably paracompact sets	383
<i>Cs. Szántó, I. Szöllősi</i> On some Hall polynomials over a quiver of type \tilde{D}_4	395
Contents of volume 12, 2020	405



Maia type fixed point results via C-class function

Arslan Hojat Ansari
Department of Mathematics,
Karaj Branch, Islamic Azad University,
Karaj, Iran
email: mathanalsisamir4@gmail.com

Mohammad Saeed Khan
Department of Mathematics &
Statistics, Sultan Qaboos University,
P. O. Box 36, Al-Khoud 123, Muscat,
Sultanate of Oman
email: mohammad@squ.edu.om

Vladimir Rakočević
Faculty of Sciences and Mathematics,
Višegradska 33, 18000 Niš, Serbia
email: vrakoc@sbb.rs (corresponding author)

Abstract. In 1968, M. G. Maia [16] generalized Banach's fixed point theorem for a set X endowed with two metrics. In 2014, Ansari [2] introduced the concept of C-class functions and generalized many fixed point theorems in the literature. In this paper, we prove some Maia's type fixed point results via C-class function in the setting of two metrics space endowed with a binary relation. Our results, generalized and extended many existing fixed point theorems, for generalized contractive and quasi-contractive mappings, in a metric space endowed with binary relation.

1 Introduction and preliminaries

The classical Banach contraction mapping is one of the most useful in metric fixed point theory. It is very popular tool for solving existence and uniqueness

2010 Mathematics Subject Classification: 46N40, 47H10, 54H25, 46T99

Key words and phrases: fixed point, generalized contraction mappings, ordered metric spaces, C-class function

problems in many different fields of mathematics. Due to its importance and applications potential, the Banach Contraction Principle has been investigated heavily by many authors. Consequently, a number of generalizations of this celebrated principle have appeared in the literature. For some recent significant book from fixed point theory, we refer to ([1, 7, 11, 14, 17, 24, 26]).

We first recall Maia's fixed point theorem:

Theorem 1 [16] *Let (X, d, δ) be a bimetric space and $T : X \rightarrow X$. Assume that the following conditions are satisfied:*

- (i) $d(x, y) \leq \delta(x, y)$ for all $x, y \in X$,
- (ii) X is complete with respect to d ,
- (iii) T is continuous with respect to d ,
- (iv) there exists a constant $\alpha \in [0, 1)$ such that

$$\delta(Tx, Ty) \leq \alpha \delta(x, y), \quad \text{for all } x, y \in X.$$

Then T has a unique fixed point in X .

Singh [28] proved that the above theorem is true under much less restricted condition, that is we do not need the continuity of T with respect to d on X , but only the continuity at a point. Many papers deal with fixed point theorems of Maia type and with applications (see eg., [6, 5, 19, 23, 22, 25, 20, 30]) and references therein). In these direction, in 2019. Petrusel and Rus [20] consider the following: Let X be nonempty set endowed with a metric d , an order relation \preceq and an operator $f : X \rightarrow X$, which satisfies two main assumptions:

- (1) f is generalized monotone with respect to \preceq ;
- (2) f is a (generalized) contraction with respect to d on a certain subset Y of $X \times X$.

Then, they apply these results to study some problems related to integral and differential equations, and several open questions are discussed. We point out that Turinici [30] have showed that the Ran-Reurings [21] fixed point theorem is but a particular case of Maia's.

In 2012. Samet and Turinici [27] introduced the notion of contractive mapping in a metric space endowed with amorphous binary relation. They showed a theorem subsumes many known results in the literature. For further study about contractive mappings in a metric space endowed with binary relation, we refer the reader to [8] [27] and [31].

In the sequel let (X, d, δ) be a bimetric space and $T : X \rightarrow X$ be a mapping. Denote by

$$\text{Fix}(T) = \{x^* \in X : x^* = Tx^*\},$$

the set of all fixed points of T in X .

Let \mathcal{R} be a binary relation on X and let \mathcal{S} be the *symmetric binary relation* defined by

$$x, y \in X, \quad x\mathcal{S}y \quad \Longleftrightarrow \quad x\mathcal{R}y \quad \text{or} \quad y\mathcal{R}x.$$

For $x_0 \in X$ we define the sequence $\{x_n\}$ by

$$x_n = Tx_{n-1} \quad \text{for all } n \in \mathbb{N}.$$

Definition 1 Let (X, δ) be metric space and $n \in \mathbb{N} \cup \{0\}$. For $A \subset X$ we denote by $\text{diam}(A) := \sup\{\delta(a, b) : a, b \in A\}$ the diameter of A . For each $x_0 \in X$ the orbit sets of T at x_0 are defined as following

$$O_n(x_0) = \{x_0, x_1, \dots, x_n\} \quad \text{and} \quad O_\infty(x_0) = \{x_0, x_1, x_2, \dots\}.$$

We say that (X, δ) is T -orbitally complete iff every δ -Cauchy sequence from $O_\infty(x)$ for some $x \in X$ converges in X .

Definition 2 [27] A subset D of X is called \mathcal{R} -directed if for every $x, y \in D$, there exists $z \in X$ such that $z\mathcal{R}x$ and $z\mathcal{R}y$.

Definition 3 A mapping $T : X \rightarrow X$ is called \mathcal{R} -preserving mapping if

$$x, y \in X, \quad x\mathcal{R}y \quad \Longrightarrow \quad Tx\mathcal{R}Ty.$$

Next, we define the set Φ of functions $\varphi : [0, +\infty) \rightarrow [0, +\infty)$ satisfying:

(I) φ is nondecreasing,

(II) $\sum_{n=1}^{\infty} \varphi^n(t) < \infty$ for each $t > 0$, where φ^n is the n -th iterate of φ .

Remark 1 Let $\varphi \in \Phi$. We have $\varphi(t) < t$ for all $t > 0$.

Remark 2 Let $\varphi \in \Phi$. We have $\lim_{n \rightarrow \infty} \varphi^n(t) = 0$ for all $t > 0$.

Definition 4 [15] Assume that for $T : X \rightarrow X$ there exists $\varphi \in \Phi$ such that

$$\delta(Tx, Ty) \leq \varphi(M_\delta(x, y)) \quad \text{for all } x, y \in X \quad \text{with } xSy.$$

A mapping T is called a generalized contractive with respect to δ if

$$M_\delta(x, y) = \max \left\{ \delta(x, y), \frac{\delta(x, Tx) + \delta(y, Ty)}{2}, \frac{\delta(x, Ty) + \delta(Tx, y)}{2} \right\}.$$

A mapping T is called a generalized quasi-contractive (see [10, 11, 17]) with respect to δ if

$$M_\delta(x, y) = \max\{\delta(x, y), \delta(x, Tx), \delta(y, Ty), \delta(x, Ty), \delta(Tx, y)\}.$$

Lemma 1 (Lemma 1 of [15]) Let (X, δ) be a metric space, and \mathcal{R} a transitive binary relation over X . Assume that for $T : X \rightarrow X$, the following conditions are satisfied:

- (b1) there exists $x_0 \in X$ such that $x_0 \mathcal{R} Tx_0$,
- (b2) T is \mathcal{R} -preserving mapping,
- (b3) T is generalized quasi-contractive with respect to δ .

Then,

$$\delta(x_i, x_j) \leq \varphi(\text{diam}(O_n(x_0))),$$

for all $i, j \in \{1, \dots, n\}$.

In 2014 the concept of C-class functions were introduced by A.H.Ansari [2]. By using this concept we can generalize many fixed point theorems in the literature. C-class functions have been studied by many authors and some fixed point results with applications (see eg., [3, 12, 18, 4, 13]).

Definition 5 [2] A mapping $F : [0, \infty)^2 \rightarrow \mathbb{R}$ is called C-class function if it is continuous and satisfies following axioms:

- (1) $F(s, t) \leq s$,
- (2) $F(s, t) = s$ implies that either $s = 0$ or $t = 0$; for all $s, t \in [0, \infty)$.

Note for some F we have that $F(0, 0) = 0$.

We denote C-class functions as \mathcal{C} .

Example 1 [2] The following functions $F : [0, \infty)^2 \rightarrow \mathbb{R}$ are elements of \mathcal{C} , for all $s, t \in [0, \infty)$:

$$(1) F(s, t) = s - t, F(s, t) = s \Rightarrow t = 0;$$

$$(2) F(s, t) = ms, 0 < m < 1, F(s, t) = s \Rightarrow s = 0;$$

$$(3) F(s, t) = \frac{s}{(1+t)^r}, r \in (0, \infty), F(s, t) = s \Rightarrow s = 0 \text{ or } t = 0;$$

$$(4) F(s, t) = \log(t + a^s)/(1 + t), a > 1, F(s, t) = s \Rightarrow s = 0 \text{ or } t = 0;$$

$$(5) F(s, t) = \ln(1 + a^s)/2, a > e, F(s, 1) = s \Rightarrow s = 0;$$

$$(6) F(s, t) = (s + l)^{(1/(1+t)^r)} - l, l > 1, r \in (0, \infty), F(s, t) = s \Rightarrow t = 0;$$

$$(7) F(s, t) = s \log_{t+a} a, a > 1, F(s, t) = s \Rightarrow s = 0 \text{ or } t = 0;$$

$$(8) F(s, t) = s - \left(\frac{1+s}{2+s}\right)\left(\frac{t}{1+t}\right), F(s, t) = s \Rightarrow t = 0;$$

$$(9) F(s, t) = s\beta(s), \beta : [0, \infty) \rightarrow [0, 1], \text{ and is continuous, } F(s, t) = s \Rightarrow s = 0;$$

$$(10) F(s, t) = s - \frac{t}{k+t}, F(s, t) = s \Rightarrow t = 0;$$

$$(11) F(s, t) = s - \varphi(s), F(s, t) = s \Rightarrow s = 0, \text{ here } \varphi : [0, \infty) \rightarrow [0, \infty) \text{ is a continuous function such that } \varphi(t) = 0 \Leftrightarrow t = 0;$$

$$(12) F(s, t) = sh(s, t), F(s, t) = s \Rightarrow s = 0, \text{ here } h : [0, \infty) \times [0, \infty) \rightarrow [0, \infty) \text{ is a continuous function such that } h(t, s) < 1 \text{ for all } t, s > 0;$$

$$(13) F(s, t) = s - \left(\frac{2+t}{1+t}\right)t, F(s, t) = s \Rightarrow t = 0;$$

$$(14) F(s, t) = \sqrt[n]{\ln(1 + s^n)}, F(s, t) = s \Rightarrow s = 0;$$

$$(15) F(s, t) = \phi(s), F(s, t) = s \Rightarrow s = 0, \text{ here } \phi : [0, \infty) \rightarrow [0, \infty) \text{ is a upper semicontinuous function such that } \phi(0) = 0, \text{ and } \phi(t) < t \text{ for } t > 0;$$

$$(16) F(s, t) = \frac{s}{(1+s)^r}, r \in (0, \infty), F(s, t) = s \Rightarrow s = 0 ;$$

$$(17) F(s, t) = \vartheta(s), \vartheta : \mathbb{R}^+ \times \mathbb{R}^+ \rightarrow \mathbb{R} \text{ is a generalized Mizoguchi-Takahashi type function, } F(s, t) = s \Rightarrow s = 0;$$

$$(18) F(s, t) = \frac{s}{\Gamma(1/2)} \int_0^\infty \frac{e^{-x}}{\sqrt{x+t}} dx, \text{ where } \Gamma \text{ is the Euler Gamma function.}$$

Denote by Ψ the family of continuous and monotone nondecreasing functions $\psi : [0, \infty) \rightarrow [0, \infty)$ such that $\psi(t) = 0$ if and only if $t = 0$ and by Φ_u the family of continuous functions $\varphi : [0, \infty) \rightarrow [0, \infty)$ such that $\varphi(t) > 0$ for all $t > 0$.

Definition 6 Assume that for $T : X \rightarrow X$ there exists $\varphi \in \Phi_u, \psi \in \Psi, F \in \mathcal{C}$ such that

$$\psi(\delta(Tx, Ty)) \leq F(\psi(M_\delta(x, y)), \varphi(M_\delta(x, y))), \quad \text{for all } x, y \in X \text{ with } xSy. \quad (1)$$

A mapping T is called a generalized $F\psi\varphi$ -contractive with respect to δ , if

$$M_\delta(x, y) = \max \left\{ \delta(x, y), \frac{\delta(x, Tx) + \delta(y, Ty)}{2}, \frac{\delta(x, Ty) + \delta(Tx, y)}{2} \right\}.$$

A mapping T is called a generalized quasi- $F\psi\varphi$ -contractive with respect to δ , if

$$M_\delta(x, y) = \max\{\delta(x, y), \delta(x, Tx), \delta(y, Ty), \delta(x, Ty), \delta(Tx, y)\}.$$

Lemma 2 [9] Suppose (X, δ) be a metric space. Let $\{x_n\}$ be a sequence in X such that $\delta(x_n, x_{n+1}) \rightarrow 0$ as $n \rightarrow \infty$. If $\{x_n\}$ is not a Cauchy sequence then there exist an $\varepsilon > 0$ and sequences of positive integers $\{m(k)\}$ and $\{n(k)\}$ with $m(k) > n(k) > k$ such that $\delta(x_{m(k)}, x_{n(k)}) \geq \varepsilon$, $\delta(x_{m(k)-1}, x_{n(k)}) < \varepsilon$ and

$$(i) \lim_{k \rightarrow \infty} \delta(x_{m(k)-1}, x_{n(k)+1}) = \varepsilon,$$

$$(ii) \lim_{k \rightarrow \infty} \delta(x_{m(k)}, x_{n(k)}) = \varepsilon,$$

$$(iii) \lim_{k \rightarrow \infty} \delta(x_{m(k)-1}, x_{n(k)}) = \varepsilon,$$

$$(iv) \lim_{k \rightarrow \infty} \delta(x_{m(k)+1}, x_{n(k)+1}) = \varepsilon,$$

$$(v) \lim_{k \rightarrow \infty} \delta(x_{m(k)}, x_{n(k)-1}) = \varepsilon.$$

Definition 7 We say (ψ, ϕ, F) is monotone if $x \leq y \implies F(\psi(x), \phi(x)) \leq F(\psi(y), \phi(y))$.

Example 2 Let $F(s, t) = s - t$, $\phi(x) = \sqrt{x}$

$$\psi(x) = \begin{cases} \sqrt{x} & \text{if } 0 \leq x \leq 1, \\ x^2, & \text{if } x > 1. \end{cases}$$

Then (ψ, ϕ, F) is monotone.

Example 3 Let $F(s, t) = s - t$, $\phi(x) = x^2$

$$\psi(x) = \begin{cases} \sqrt{x} & \text{if } 0 \leq x \leq 1, \\ x^2, & \text{if } x > 1. \end{cases}$$

Then (ψ, ϕ, F) is not monotone.

In this paper we prove some Maia type fixed point results via C-class function in the setting of two metrics space endowed with a binary relation. Our results generalized and extended many existing fixed point theorems for generalized contractive and quasi-contractive mappings in a metric space endowed with binary relation.

2 Main results

Our first main result is the following theorem.

Theorem 2 *Let (X, d, δ) be a bimetric space and $T : X \rightarrow X$. Assume that the following conditions are satisfied:*

- (A1) $d(x, y) \leq \delta(x, y)$ for all $x, y \in X$,
- (A2) (X, d) is T -orbitally complete,
- (A3) T is continuous with respect to d ,
- (A4) T is \mathcal{S} -preserving,
- (A5) there exists $x_0 \in X$ with x_0STx_0 ,
- (A6) T is a generalized $F\psi\varphi$ -contractive with respect to δ .

Then T has a fixed point x^ in X . Moreover, if in addition $\text{Fix}(T)$ is \mathcal{S} -directed then x^* is the unique fixed point of T in X .*

Proof. From (A5), there exists $x_0 \in X$ with x_0STx_0 and from (A4) T is \mathcal{S} -preserving, we get

$$x_nSTx_n \quad \text{for all } n \in \mathbb{N}. \quad (2)$$

If $x_n = Tx_n$ then x_n is a fixed point of T . Suppose that $x_n \neq Tx_n$ for all n . Since (2) is satisfied for all $n \geq 1$, by applying the contraction condition (A6), and note that ψ is nondecreasing, we have

$$\begin{aligned} \psi(\delta(x_n, x_{n+1})) &= \psi(\delta(x_n, Tx_n)) \leq F(\psi(M_\delta(x_{n-1}, x_n)), \varphi(M_\delta(x_{n-1}, x_n))) \\ &< \psi(M_\delta(x_{n-1}, x_n)) \\ &\leq \psi(\max\{\delta(x_{n-1}, x_n), \delta(x_n, x_{n+1})\}). \end{aligned}$$

Now, we will show that $\{x_n\}$ is a Cauchy sequence in (X, δ) . If for some $n \geq 1$ we have $\delta(x_{n-1}, x_n) \leq \delta(x_n, x_{n+1})$, then we get

$$\psi(\delta(x_n, x_{n+1})) = F(\psi(\delta(x_n, x_{n+1})), \varphi(\delta(x_n, x_{n+1}))).$$

Thus, $\psi(\delta(x_n, x_{n+1})) = 0$ or $\varphi(\delta(x_n, x_{n+1})) = 0$, and therefore $\delta(x_n, x_{n+1}) = 0$ which is contradiction. We get $\delta(x_{n-1}, x_n) > \delta(x_n, x_{n+1})$ and

$$\psi(\delta(x_n, x_{n+1})) = F(\psi(\delta(x_{n-1}, x_n)), \varphi(\delta(x_{n-1}, x_n))). \quad (3)$$

Hence $\{\delta(x_n, x_{n+1})\}$ is a non-increasing sequence of positive real numbers. Thus there exist $L \geq 0$ such that

$$\lim_{n \rightarrow \infty} \delta(x_n, x_{n+1}) = L. \quad (4)$$

Taking the limit in equation (3) as $n \rightarrow \infty$ and using (4) and the properties of F and φ , we have

$$\psi(L) \leq F(\psi(L), \varphi(L)).$$

Thus $\psi(L) = 0$ or $\varphi(L) = 0$, and so $L = 0$. Therefore

$$\lim_{n \rightarrow \infty} \delta(x_n, x_{n+1}) = 0. \quad (5)$$

Let us show that $\{x_n\}$ is a Cauchy sequence. Suppose to the contrary that $\{x_n\}$ is not a Cauchy sequence.

By Lemma 2 there exists $\varepsilon > 0$ for which we can find subsequences $\{x_{n(k)}\}$ and $\{x_{m(k)}\}$ of $\{x_n\}$ with $n(k) > m(k) > k$ such that

$$\delta(x_{m(k)}, x_{n(k)+1}), \delta(x_{m(k)}, x_{n(k)}), \delta(x_{m(k)-1}, x_{n(k)+1}), \delta(x_{m(k)-1}, x_{n(k)}) \rightarrow \varepsilon.$$

Now from (1) we have

$$\begin{aligned} \psi(\delta(x_{m(k)}, x_{n(k)})) &= \psi(\delta(Tx_{m(k)-1}, Tx_{n(k)-1})) \\ &\leq F(\psi(M_\delta(x_{m(k)-1}, x_{n(k)-1})), \varphi(M_\delta(x_{m(k)-1}, x_{n(k)-1}))) \end{aligned} \quad (6)$$

where

$$\begin{aligned} M_\delta(x_{m(k)-1}, x_{n(k)-1}) &= \max \left\{ \delta(x_{m(k)-1}, x_{n(k)-1}), \right. \\ &\quad \frac{\delta(x_{m(k)-1}, Tx_{m(k)-1}) + \delta(x_{n(k)-1}, Tx_{n(k)-1})}{2}, \\ &\quad \left. \frac{\delta(x_{m(k)-1}, Tx_{n(k)-1}) + \delta(Tx_{m(k)-1}, x_{n(k)-1})}{2} \right\}. \end{aligned}$$

From above and (6), as $k \rightarrow \infty$ we have

$$\psi(\varepsilon) \leq F(\psi(\varepsilon), \varphi(\varepsilon)).$$

Thus $\psi(\varepsilon) = 0$ or $\varphi(\varepsilon) = 0$ and therefore $\varepsilon = 0$ which is contradiction. Consequently the sequence $\{x_n\}$ is δ -Cauchy, so by (A1), $\{x_n\}$ is d -Cauchy too. Since from (A2), we have that the metric space (X, d) is T -orbitally complete, then there exists $x^* \in X$ such that

$$\lim_{n \rightarrow \infty} \delta(x_n, x^*) = 0. \quad (7)$$

From (A3), we have that T is continuous with respect to d , and, so it follows that $x^* = \lim_{n \rightarrow \infty} Tx_n = T(\lim_{n \rightarrow \infty} x_n) = Tx^*$, that is, x^* is a fixed point of T .

Next suppose that $\text{Fix}(T)$ is \mathcal{S} -directed, and we will show that x^* is the unique fixed point of T in X . Suppose that $y^* \in \text{Fix}(T)$ is another fixed point of T . Then, there exists $z_0 \in X$ such that $z_0 \mathcal{S} x^*$ and $z_0 \mathcal{S} y^*$. Define the sequence $\{z_n\}$ in X by $z_{n+1} = Tz_n$ for all $n \geq 0$. Since T is \mathcal{S} -preserving, for all $n \geq 0$ we have $z_n \mathcal{S} x^*$ and $z_n \mathcal{S} y^*$. Applying (A6), for all $n \geq 0$ and note $\delta(z_n, z_{n+1}) \leq \delta(z_n, x^*) + \delta(z_{n+1}, x^*)$ we get

$$\begin{aligned} \psi(\delta(z_{n+1}, x^*)) &= \psi(\delta(Tz_n, Tx^*)) = F(\psi(M_\delta(z_n, x^*)), \varphi(M_\delta(z_n, x^*))) \\ &\leq F\left(\psi\left(\max\left\{\delta(z_n, x^*), \frac{\delta(z_n, Tz_n) + \delta(x^*, Tx^*)}{2}, \frac{\delta(z_n, Tx^*) + \delta(x^*, Tz_n)}{2}\right\}\right), \right. \\ &\quad \left. \varphi\left(\max\left\{\delta(z_n, x^*), \frac{\delta(z_n, Tz_n) + \delta(x^*, Tx^*)}{2}, \frac{\delta(z_n, Tx^*) + \delta(x^*, Tz_n)}{2}\right\}\right)\right) \\ &= F\left(\psi\left(\max\left\{\delta(z_n, x^*), \frac{\delta(z_n, z_{n+1}) + \delta(x^*, x^*)}{2}, \frac{\delta(z_n, x^*) + \delta(x^*, z_{n+1})}{2}\right\}\right), \right. \\ &\quad \left. \varphi\left(\max\left\{\delta(z_n, x^*), \frac{\delta(z_n, z_{n+1}) + \delta(x^*, x^*)}{2}, \frac{\delta(z_n, x^*) + \delta(x^*, z_{n+1})}{2}\right\}\right)\right) \\ &\leq F(\psi(\max\{\delta(z_n, x^*), \delta(z_{n+1}, x^*)\}), \varphi(\max\{\delta(z_n, x^*), \delta(z_{n+1}, x^*)\})). \end{aligned}$$

Now we will show that $\lim_{n \rightarrow \infty} \delta(z_n, x^*) = 0$. Without the loss of generality suppose that $\delta(z_n, x^*) > 0$ for all n . Assume that for some n we have $\delta(z_n, x^*) \leq \delta(x^*, z_{n+1})$. Hence we get

$$\psi(\delta(z_{n+1}, x^*)) = F(\psi(\delta(z_{n+1}, x^*)), \varphi(\delta(z_{n+1}, x^*))).$$

Hence $\psi(\delta(z_{n+1}, x^*)) = 0$ or $\varphi(\delta(z_{n+1}, x^*)) = 0$ and therefore $\delta(z_{n+1}, x^*) = 0$, which is a contradiction. Then, for all $n \geq 0$ we have $\delta(z_n, x^*) > \delta(x^*, z_{n+1})$. Consequently, for all n we obtain

$$\psi(\delta(z_{n+1}, x^*)) = F(\psi(\delta(z_n, x^*)), \varphi(\delta(z_n, x^*))) \leq \psi(\delta(z_n, x^*)) \quad (8)$$

that is, $\{\delta(z_n, x^*)\}$ is a non-increasing sequence of positive real numbers. Thus there exist $L \geq 0$ such that

$$\lim_{n \rightarrow \infty} \delta(z_n, x^*) = L. \quad (9)$$

Taking the limit in equation (8) as $n \rightarrow \infty$ and using (9) and the properties of F and φ we have

$$\psi(L) \leq F(\psi(L), \varphi(L)).$$

Thus $\psi(L) = 0$ or $\varphi(L) = 0$ and therefore $L = 0$. Thus

$$\lim_{n \rightarrow \infty} \delta(z_n, x^*) = 0.$$

Similarly we can prove that $\lim_{n \rightarrow \infty} \delta(z_n, y^*) = 0$. Hence $x^* = y^*$. \square

To prove our next main result we need the following lemmas which will be used in the sequel.

Lemma 3 *Let $n \in \mathbb{N}$, (X, δ) be a metric space, and \mathcal{R} a transitive binary relation over X . Assume that for $T : X \rightarrow X$ the following conditions are satisfied:*

- (a1) *there exists $x_0 \in X$ such that $x_0 \mathcal{R} Tx_0$,*
- (a2) *T is \mathcal{R} -preserving mapping,*
- (a3) *T is generalized quasi- $F\varphi$ -contractive with respect to δ and (ψ, φ, F) is monotone.*

Then

$$\psi(\delta(x_i, x_j)) \leq F(\psi(\text{diam}(O_n(x_0))), \varphi(\text{diam}(O_n(x_0)))), \quad (10)$$

for all $i, j \in \{1, \dots, n\}$.

Proof. From (a1) there exists $x_0 \in X$ such that $x_0 \mathcal{R} Tx_0$. Hence by (a2) we get $x_k \mathcal{R} x_{k+1}$ for all k . Since \mathcal{R} is transitive, then

$$x_{i-1} \mathcal{R} x_{j-1} \quad \text{for all } 1 \leq i < j \leq n. \quad (11)$$

We note that $x_{i-1}, x_i, x_{j-1}, x_j \in O_n(x_0)$. Now using (a3) and (11) we get

$$\begin{aligned} \psi(\delta(Tx_{i-1}, Tx_{j-1})) &\leq F(\psi(M_\delta(x_{i-1}, x_{j-1})), \varphi(M_\delta(x_{i-1}, x_{j-1}))) \\ &= F(\psi(\max\{\delta(x_{i-1}, x_{j-1}), \delta(x_{i-1}, Tx_{i-1}), \delta(x_{j-1}, Tx_{j-1})\}), \varphi(\max\{\delta(x_{i-1}, x_{j-1}), \delta(x_{i-1}, Tx_{i-1}), \delta(x_{j-1}, Tx_{j-1})\})). \end{aligned}$$

$$\begin{aligned} & \delta(x_{j-1}, Tx_{j-1}), \delta(x_{i-1}, Tx_{j-1}), \delta(Tx_{i-1}, x_{j-1})), \\ & \varphi(\max\{\delta(x_{i-1}, x_{j-1}), \delta(x_{i-1}, Tx_{i-1}), \\ & \delta(x_{j-1}, Tx_{j-1}), \delta(x_{i-1}, Tx_{j-1}), \delta(Tx_{i-1}, x_{j-1})\}), \end{aligned}$$

which implies (10). \square

Now we are ready to state our second main result.

Theorem 3 *Let (X, d, δ) be a bimetric space, \mathcal{R} a transitive binary relation over X and $T : X \rightarrow X$. Assume that the following conditions are satisfied:*

- (B1) $d(x, y) \leq \delta(x, y)$ for all $x, y \in X$,
- (B2) (X, d) is T -orbitally complete,
- (B3) T is continuous with respect to d ,
- (B4) T is \mathcal{R} -preserving,
- (B5) there exists $x_0 \in X$ with $x_0 \mathcal{R} Tx_0$,
- (B6) T is a generalized quasi- $F\psi\varphi$ -contractive with respect to δ and (ψ, φ, F) is monotone.

Then T has a fixed point x^ in X . Moreover if in addition \mathcal{R} is symmetric and $\text{Fix}(T)$ is \mathcal{R} -directed then x^* is the unique fixed point of T in X .*

Proof. Let $x_0 \in X$ and $x_0 \mathcal{R} Tx_0$. Define a sequence $\{x_n\}$ in X by $x_{n+1} = Tx_n$, for all $n \geq 0$. Since T is an \mathcal{R} -preserving, then $x_n \mathcal{R} x_{n+1}$ for all n . Let n and m , $n < m$ be any positive integers. From (B6) and Lemma 3 it follows

$$\begin{aligned} \psi(\delta(T^n x_0, T^m x_0)) &= \psi(\delta(TT^{n-1} x_0, T^{m-n+1} T^{n-1} x_0)) \\ F(\psi(\text{diam}(O_{m-n+1}(T^{n-1} x_0))), \varphi(\text{diam}(O_{m-n+1}(T^{n-1} x_0)))) & \end{aligned}$$

From Remark 1 there exists an integer k_1 , $1 \leq k_1 \leq m - n + 1$ such that

$$\text{diam}(O_{m-n+1}(T^{n-1} x_0)) = \delta(T^{n-1} x_0, T^{k_1} T^{n-1} x_0).$$

Using Lemma 3 again we get combining the above inequalities

$$\begin{aligned} \psi(\delta(T^n x_0, T^m x_0)) &= \psi(\delta(TT^{n-1} x_0, T^{m-n+1} T^{n-1} x_0)) \\ F(\psi(\text{diam}(O_{m-n+1}(T^{n-1} x_0))), \varphi(\text{diam}(O_{m-n+1}(T^{n-1} x_0)))) & \\ \psi(\text{diam}(O_{m-n+1}(T^{n-1} x_0))) &= \psi(\delta(T^{n-1} x_0, T^{k_1} T^{n-1} x_0)) \end{aligned}$$

$$\begin{aligned} &\leq F(\psi(\text{diam}(O_{k_1+1}(T^{n-2}x_0))), \varphi(\text{diam}(O_{k_1+1}(T^{n-2}x_0)))) \\ &\leq F(\psi(\text{diam}(O_{m-n+2}(T^{n-2}x_0))), \varphi(\text{diam}(O_{m-n+2}(T^{n-2}x_0)))). \end{aligned}$$

Continue this process we obtain

$$\begin{aligned} \psi(\delta(T^n x_0, T^m x_0)) &= \psi(\delta(TT^{n-1}x_0, T^{m-n+1}T^{n-1}x_0)) \\ &\quad F(\psi(\text{diam}(O_{m-n+1}(T^{n-1}x_0))), \varphi(\text{diam}(O_{m-n+1}(T^{n-1}x_0)))) \\ &\quad \psi(\text{diam}(O_{m-n+1}(T^{n-1}x_0))) = \psi(\delta(T^{n-1}x_0, T^k T^{n-1}x_0)) \\ &\leq F(\psi(\text{diam}(O_{k_1+1}(T^{n-2}x_0))), \varphi(\text{diam}(O_{k_1+1}(T^{n-2}x_0)))) \\ &\leq F(\psi(\text{diam}(O_{m-n+2}(T^{n-2}x_0))), \varphi(\text{diam}(O_{m-n+2}(T^{n-2}x_0)))) \\ &\quad \vdots \\ &\leq F(\psi(\text{diam}(O_m(x_0))), \varphi(\text{diam}(O_m(x_0)))) \\ &\leq F(\psi(\delta(T^{n-1}x_0, T^{m-1}x_0)), \varphi(\delta(T^{n-1}x_0, T^{m-1}x_0))). \end{aligned}$$

Hence

$$\psi(\varepsilon) \leq F(\psi(\varepsilon), \varphi(\varepsilon)).$$

Thus $\psi(\varepsilon) = 0$ or $\varphi(\varepsilon) = 0$, that is $\varepsilon = 0$. It follows that the sequence $\{T^n x_0\}$ is a δ -Cauchy sequence. Therefore by (B1) the sequence $\{T^n x_0\}$ is a d -Cauchy sequence too. Since the metric space (X, d) is T -orbitally complete we deduce that the sequence $\{T^n x_0\}$ converges to some x^* in X . From (B3) T is continuous with respect to d , so $x^* = \lim_{n \rightarrow \infty} T x_n = T(\lim_{n \rightarrow \infty} x_n) = T x^*$ and x^* is a fixed point of T .

Now suppose that $\text{Fix}(T)$ is \mathcal{R} -directed. We claim that the fixed point is unique. Let x^* and y^* be two fixed points of T . Suppose that $x^* \neq y^*$. Since $\text{Fix}(T)$ is \mathcal{R} -directed, then there exists $z \in X$ such that $z\mathcal{R}x^*$ and $z\mathcal{R}y^*$. By the transitivity of \mathcal{R} we have $x^*\mathcal{R}y^*$. Then we apply the contraction condition (B6) and get

$$\begin{aligned} \psi(\delta(x^*, y^*)) &= \psi(\delta(Tx^*, Ty^*)) \\ &\leq F(\psi(\max\{\delta(x^*, y^*), \delta(x^*, Tx^*), \delta(y^*, Ty^*), \delta(x^*, Ty^*), \delta(Tx^*, y^*)\}), \\ &\quad \varphi(\max\{\delta(x^*, y^*), \delta(x^*, Tx^*), \delta(y^*, Ty^*), \delta(x^*, Ty^*), \delta(Tx^*, y^*)\})). \end{aligned}$$

Hence

$$\psi(\delta(x^*, y^*)) \leq F(\psi(\delta(x^*, y^*)), \varphi(\delta(x^*, y^*)))$$

and $\psi(\delta(x^*, y^*)) = 0$ or $\varphi(\delta(x^*, y^*)) = 0$. Therefore $\delta(x^*, y^*) = 0$ and $x^* = y^*$.

□

The following results are an immediate consequences of Theorems 2 and 3.

Corollary 1 *Theorem 1 is a particular case of Theorem 2.*

Corollary 2 *Let (X, \preceq) be a partially ordered set and (X, d, δ) be a bimetric space and $T : X \rightarrow X$. Assume that the following conditions are satisfied:*

(C1) $d(x, y) \leq \delta(x, y)$, for all $x, y \in X$,

(C2) X is complete with respect to d ,

(C3) T is continuous with respect to d ,

(C4) T is monotone nondecreasing mapping,

(C5) there exists $x_0 \in X$ with $x_0 \preceq Tx_0$,

(C6) there exists $\varphi \in \Phi$ such that

$$\psi(\delta(Tx, Ty)) \leq \psi(M_\delta(x, y)) - \varphi(M_\delta(x, y)), \quad \text{for all } x, y \text{ in } X.$$

Then T has a unique fixed point in X .

Corollary 3 *Let (X, \preceq) be a partially ordered set and (X, d) be a complete metric space. Assume that for $T : X \rightarrow X$, the following conditions are satisfied:*

(D1) T is continuous;

(D2) T is monotone nondecreasing mapping;

(D3) there exists $x_0 \in X$ with $x_0 \preceq Tx_0$;

(D4) there exists a constant $\alpha \in [0, 1)$ such that

$$\psi(\delta(Tx, Ty)) \leq \alpha \psi(M_\delta(x, y)), \quad \text{for all } x, y \text{ in } X.$$

Then T has a unique fixed point in X .

Corollary 4 *Let (X, d) be a complete metric space and $T : X \rightarrow X$ be continuous mapping. Suppose there exists $\varphi \in \Phi$ such that*

$$\psi(\delta(Tx, Ty)) \leq \frac{\psi(M_\delta(x, y))}{1 + \varphi(M_\delta(x, y))}, \quad \text{for all } x, y \text{ in } X.$$

Then T has a unique fixed point in X .

Remark 3

(1) If in Theorem 2 we put $F(s, t) = mt$, $0 \leq m < 1$, $\psi(t) = \phi(t) = t$, then we obtain Maia's Theorem 1.

(2) If we use the same notations as in (1), and if we define relation \mathcal{S} by $x\mathcal{S}y$ if and only if $\alpha d(xTx) \leq d(x, y)$ implies $d(Tx, Ty) \leq \beta d(x, y)$, where $\alpha \in (0, 1/2)$, $\beta \in (0, 1)$, then when T is continuous Theorem 2 implies Theorem 2.2 in [29].

(3) Our results, when we put $F(s, t) = mt$, $0 \leq m < 1$, imply results from [15].

(4) Using Theorem of Singh [28] we note that our results are true under much less restricted condition, that is we do not need the continuity of T with respect to d on X , but only the continuity at a point.

3 Application to Cauchy problem

In this section, we study the Cauchy problem for a class of nonlinear differential equations, using the results obtained in the previous section. We just state the application part and we point out that the proof is on the lines of M.S. Khan at all [15]. So we omit it.

Example 4 Consider the nonlinear differential equation

$$\psi(x) = \begin{cases} x'(t) = f(t, x(t)) & t \in [a, b], \\ x(t_0) = x_0 \end{cases} \quad (12)$$

where $a, b, t_0 \in \mathbb{R}$ and $f : [a, b] \times \mathbb{R} \rightarrow \mathbb{R}$. Let $X = C([a, b], \mathbb{R})$ denotes the space of all continuous \mathbb{R} -valued functions on $[a, b]$ with the metric d given by

$$d(u, v) = \sup_{t \in [a, b]} |f(u(t), v(t))|, \text{ for all } u, v \in X$$

It is well known that (X, d) is a complete metric space. We define an order relation \preceq on X by

$$u \preceq v \Leftrightarrow u(t) \leq v(t), \text{ for all } t \in [a, b]$$

Consider the mapping $T : C([a, b], \mathbb{R}) \rightarrow C([a, b], \mathbb{R})$ defined by

$$Tx(t) = x_0 + \int_{t_0}^t f(s, x(s)) ds ; \quad t \in [a, b]$$

for all $x \in C([a, b], \mathbb{R})$. Clearly, $x^* \in C([a, b], \mathbb{R})$ is a solution of (12) if and only if x^* is a fixed point of T .

Furthermore, we consider the following assumptions:

(H1) $f : [a, b] \times \mathbb{R} \rightarrow \mathbb{R}$ is continuous;

(H2) $f : [a, b] \times \mathbb{R} \rightarrow \mathbb{R}$ is nondecreasing with respect to the second variable;

(H3) $|f(t, x(t)) - f(t, y(t))| \leq L|x(t) - y(t)|$ for all $x(t) \leq y(t)$ and $t \in [a, b]$.

It is worth noting that condition (H3) is weaker compared to those used by Maia for studying Cauchy problem in [16], that is, f is L -Lipschitzien function on the whole space.

By the proof of Theorem 6 in [15] we have

$$\delta(x, y) \leq \exp(L(b - a))d(x, y)$$

and for $\lambda = \sqrt{1 - \exp(L(a - b))}$, then we have

$$d(T^n x, T^n y) \leq \exp(L(b - a))d(x, y) \quad \text{for all } x \preceq y. \quad (13)$$

We deduce by using Corollary 3 (see also [15]) that T has a unique fixed point $x^* \in C([a, b], \mathbb{R})$.

Acknowledgements

The authors are grateful to the anonymous reviewers for their careful reviews and very useful comments.

References

- [1] R. P. Agarwal, E. Karapinar, D. O'Regan, Antonio Francisco Roldan-Lopez-de-Hierro, Fixed Point Theory in Metric Type Spaces, Springer International Publishing, Switzerland 2015.
- [2] A. H. Ansari, Note on $\varphi - \psi$ - contractive type mappings and related fixed point, The 2nd Regional Conference on Mathematics and Applications, Payame Noor University, 2014, pages 377–380.

- [3] A. H. Ansari, M. Demma, L. Guran and J. R. Lee, Fixed point results for C-class functions in modular metric spaces, *J. Fixed Point Theory Appl.*, (2018) 20: 103.
- [4] A. H. Ansari, V. Gupta, N. Mani, C-class functions on Some Coupled Fixed Point Theorems in partially ordered S-metric spaces, Communications Faculty of Sciences University of Ankara Series A1: Mathematics and Statistics, **68** (2) (2019), 1694–1708,.
- [5] M. -E. Balazs, Maia type fixed point theorems for Ćirić–Prešić operators, *Acta Univ. Sapientiae, Mathematica*, **10**, 1 (2018), 18–31.
- [6] V. Berinde, *A fixed point theorem of Maia type in K-metric spaces*, Seminar on Fixed Point Theory (Cluj-Napoca), **3** (1991), 7–14.
- [7] V. Berinde, *Iterative Approximation of Fixed Points*, Springer, Berlin, 2007.
- [8] M. Berzig, Coincidence and common fixed point results on metric spaces endowed with an arbitrary binary relation and applications, *J. Fixed Point Theory Appl.*, **12**.1-2, (2012), 221–238.
- [9] P. Chuadchawna, A. Kaewcharoen, S. Plubtieng, Fixed point theorems for generalized $\alpha - \eta - \psi$ -Geraghty contraction type mappings in $\alpha - \eta$ complete metric spaces, *J. Nonlinear Sci. Appl.*, **9** (2016), 471–485.
- [10] Lj. B. Ćirić, A generalization of Banach's contraction principle, *Proc. Amer. Math. Soc.*, **45**.2 (1974), 267–273.
- [11] Lj. B. Ćirić, *Some recent results in metrical fixed point theory*, University of Belgrade, Beograd 2003.
- [12] D. Dhamodharan, Yumnam Rohen and Arslan Hojat Ansari, Fixed point theorems of C-class functions in S_b -metric space, *Res. Fixed Point Theory Appl.* Volume 2018, Article ID 2018018, 20 pages.
- [13] V. Gupta, N. Mani, N. Sharma, Fixed point theorems for weak $(,)$ - mappings satisfying generalized C-condition and its application to boundary value problem, *Computational and Mathematical Methods*, **1** (4), e1041, 1–12, 2019.
- [14] W. Kirk, N. Shahzad, *Fixed Point Theory in Distance Spaces*, Springer International Publishing, Switzerland 2014.

- [15] M.S. Khan, M. Berzig and S. Chandok, Fixed point theorems in bimetric space endowed with binary relation and applications, *Miskolc Mathematical Notes*, Vol. **16** (2) (2015), 939–951.
- [16] M. G. Maia, *Un'osservazione sulle contrazioni metriche*, Rend. Semin. Mat. Univ. Padova, **40** (1968), 139–143.
- [17] E. Malkowsky and V. Rakočević, *Advanced Functional Analysis*, CRC Press, Taylor & Francis Group, Boca Raton, FL, 2019.
- [18] B. Moeini, *C-class functions and some fixed point results with applications*, LAMBERT Academic Publishing, 2018.
- [19] A. S. Muresan, From Maia fixed point theorem to the fixed point theory in a set with two metrics, *Carpatian J. Math.*, **23** (2007), 133–140.
- [20] A. Petrusel and I. A. Rus, Fixed point theory in terms of a metric and of an order relation, *Fixed Point Theory*, **20** (2) (2019), 601–622.
- [21] A. C. M. Ran and M. C. Reurings, A fixed point theorem in partially ordered sets and some applications to matrix equations, *Proc. Amer. Math. Soc.*, **132** (2004), 1435–1443.
- [22] D. O'Regan and A. Petrusel, Fixed point theorems for generalized contractions in ordered metric spaces, *J. Math. Anal. Appl.*, **341**(2008), 1241–1252.
- [23] I. A. Rus, *Basic problem for Maia's theorem*, Sem. on Fixed Point Theory, Preprint, **3** (1981), Babes-Bolyai Univ., Cluj-Napoca, 112–115.
- [24] I. A. Rus, *Generalized Contractions and Applications*, Cluj University Press, Cluj-Napoca, 2001.
- [25] I. A. Rus, Data dependence of the fixed points in a set with two metrics, *Fixed Point Theory*, Volume **8** No. 1 (2007), 115–123.
- [26] I. A. Rus, A. Petrusel, G. Petrusel, *Fixed Point Theory*, Cluj University Press Cluj-Napoca, 2008.
- [27] B. Samet and M. Turinici, Fixed point theorems on a metric space endowed with an arbitrary binary relation and applications, *Commun. Math. Anal.*, **13.2** (2012), 82–97.

- [28] S. P. Singh, On a fixed point theorem in metric space, *Rend. Semin. Mat. Univ. Padova*, **43** (1970), 229–232.
- [29] T. Suzuki, A new type of fixed point theorem in metric spaces, *Nonlinear Analysis*, **71** (2009), 5313–5317.
- [30] M. Turinici, *Ran-Reurings theorems in ordered Metric Spaces*, preprint, arxiv:1103.5207(2011).
- [31] M. Turinici, Contractive maps in locally transitive relational metric spaces, *The Scientific World Journal*, vol. 2014, article ID 169358.

Received: February 2, 2020



On the convergence difference sequences and the related operator norms

P. Baliarsingh

Department of Mathematics, Gangadhar
Meher University, Sambalpur-768004,
Odisha, India
email: pb.math10@gmail.com

L. Nayak*

Department of Mathematics, School
of Applied Sciences, Kalinga
Institute of Industrial Technology,
Bhubaneswar-751024, India
email: laxmipriyamath@gmail.com

S. Samantaray

Department of Mathematics, School of Applied
Sciences, Kalinga Institute of Industrial Technology,
Bhubaneswar-751024, India
Kalinga Institute of Industrial Technology,
Bhubaneswar-751024, India
email: samantaraysnigdha1995@gmail.com

Abstract. In this note, we discuss the definitions of the difference sequences defined earlier by Kızmaz (1981), Et and Çolak (1995), Malkowsky et al. (2007), Başar(2012), Baliarsingh (2013, 2015) and many others. Several authors have defined the difference sequence spaces and studied their various properties. It is quite natural to analyze the convergence of the corresponding sequences. As a part of this work, a convergence analysis of difference sequence of fractional order defined earlier is presented. It is demonstrated that the convergence of the fractional difference sequence is dynamic in nature and some of the results involved are also inconsistent. We provide certain stronger conditions on the primary sequence and the results due to earlier authors are substantially modified. Some illustrative examples are provided for each point of the modifications. Results on certain operator norms related to the difference operator of fractional order are also determined.

2010 Mathematics Subject Classification: 40A07; 40C09; 46B08

Key words and phrases: sequence spaces, difference sequence spaces, difference operators Δ^α and Δ^m , operator norms

*corresponding author

1 Introduction

Recently, one of the most interesting areas of research in Mathematics is the study of difference operators and related sequence spaces which has been attracted in different areas of Mathematical sciences especially in applied and computational fields involving calculus, matrix and approximation theory. The idea of difference sequence spaces plays a key role in most of the scientific problems involving the spectral properties of bounded linear operators (see [2, 7, 11, 15, 16, 28, 29, 30]), related topological structures (see [3, 4, 19, 20, 22, 26, 27]), matrix transformations (see [5, 12, 18, 19, 21, 23]), compact operators (see [1, 14, 24, 25]), fractional calculus [8, 9, 10], etc.

In fact, the study of all the ideas discussed earlier is only feasible and even possible if the related sequences are convergent.

Let $x = (x_k)$ be any sequence in w , the family of all real valued sequences. Let \mathbb{N} be the set of all positive integers and $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$. A sequence $x = (x_k)$ is said to be of order k^α , i.e., $x_k = \mathcal{O}(k^\alpha)$ if for a positive constant \mathcal{C} , we can write

$$|x_k| \leq \mathcal{C}k^\alpha, \quad k = 0, 1, 2, 3, \dots$$

By $\ell_{\infty, c}$ and c_0 , we denote the spaces of all bounded, convergent and null sequences, respectively, normed by

$$\|x\|_{\infty} = \sup_k |x_k|.$$

We use the notation ℓ_p , ($1 \leq p < \infty$) for the space of all p -summable sequence with the norm

$$\|x\|_p = \left(\sum_{k=0}^{\infty} |x_k|^p \right)^{1/p}.$$

The 1st order difference sequence space $X(\Delta)$ for $X \in \{\ell_{\infty, c}, c_0\}$ was introduced by Kizmaz [20] using forward difference operator Δ , where

$$\Delta x_k = x_k - x_{k+1}, \quad (k \in \mathbb{N}_0). \quad (1)$$

Later on, this idea has been generalized to the case of difference sequence spaces of integer order m by Et and Çolak [17] using the operator Δ^m and

$$\Delta^m x_k = \sum_{i=0}^m (-1)^i \binom{m}{i} x_{k+i}, \quad (k \in \mathbb{N}_0). \quad (2)$$

Using Euler gamma function for a proper fraction α , the fractional difference sequence $\Delta^\alpha x$ of order α was defined by Baliarsingh [4] (see also [5, 6]) as

$$\Delta^\alpha x_k = \sum_{i=0}^{\infty} (-1)^i \frac{\Gamma(\alpha+1)}{i! \Gamma(\alpha-i+1)} x_{k+i}, \quad (k \in \mathbb{N}_0). \quad (3)$$

By taking inverse transform $\Delta^{-\alpha}$ on the sequence $x = (x_k)$, we write the Eqn. (3) as

$$\Delta^{-\alpha} x_k = x_k + \alpha x_{k+1} + \frac{\alpha(\alpha+1)}{2!} x_{k+2} + \frac{\alpha(\alpha+1)(\alpha+1)}{3!} x_{k+3} + \dots \quad (4)$$

An infinite series has no meaning unless it converges. It is important to mention that in the previous papers, the convergence of the fractional difference sequence defined by (3) and (4) have been presumed without taking any further investigations. Now, in particular, we illustrate the following examples regarding the convergence of these series:

Example 1 Let α be a proper fraction and $x = (x_k)$ be the convergent sequence defined by $x_k = \frac{1}{3^k}$ for all $k \in \mathbb{N}_0$. Then, we can easily calculate

$$\Delta^\alpha x_k = \sum_{i=0}^{\infty} (-1)^i \frac{\Gamma(\alpha+1)}{i! \Gamma(\alpha-i+1)} \frac{1}{3^{k+i}} = \frac{1}{3^k} \left(\frac{2}{3} \right)^\alpha = \frac{2^\alpha}{3^{k+\alpha}} \rightarrow 0 \text{ as } k \rightarrow \infty,$$

and

$$\Delta^{-\alpha} x_k = \sum_{i=0}^{\infty} \frac{\alpha(\alpha+1) \dots (\alpha+i-1)}{\Gamma(i+1)} \frac{1}{3^{k+i}} = \frac{1}{3^k} \left(\frac{2}{3} \right)^{-\alpha} = \frac{3^{\alpha-k}}{2^\alpha} \rightarrow 0 \text{ as } k \rightarrow \infty.$$

Example 2 Let $x = (x_k)$ be the constant sequence with $x_k = 1$ for all $k \in \mathbb{N}_0$. Although the sequence $x = (x_k)$ is convergent, but for a proper fraction α , $\Delta^\alpha x_k \rightarrow 0$ as $k \rightarrow \infty$ whereas, $\Delta^{-\alpha} x_k \rightarrow \infty$ as $k \rightarrow \infty$.

Example 3 Let $x = (x_k)$ be the oscillating sequence, defined by $x_k = (-1)^k$ for all $k \in \mathbb{N}_0$. Clearly, the sequence $x = (x_k)$ is divergent and for a proper fraction α , we have

$$\Delta^\alpha x_k = \begin{cases} 2^\alpha, & (k \text{ is even}) \\ -2^\alpha, & (k \text{ is odd}), \end{cases}$$

and

$$\Delta^{-\alpha} x_k = \begin{cases} 2^{-\alpha}, & (k \text{ is even}) \\ -2^{-\alpha}, & (k \text{ is odd}) \end{cases}$$

are also divergent.

Example 4 Let $x = (x_k)$ be the divergent sequence defined by $x_k = k$ for all $k \in \mathbb{N}_0$. Although the sequence $x = (x_k)$ is divergent, but for an integer $\alpha > 1$, $\Delta^\alpha x_k \rightarrow 0$ as $k \rightarrow \infty$ whereas, $\Delta^{-\alpha} x_k \rightarrow \infty$ as $k \rightarrow \infty$. For a proper fraction α , both of the difference sequences go to ∞ as $k \rightarrow \infty$.

It is remarked that the infinite series defined in (3) and (4) need not be convergent for any arbitrary sequence $x = (x_k)$ and any proper fraction α . Therefore, it is quite difficult to study and analyze the behaviors of the related sequence spaces for fractional cases. As the convergence of the difference sequence $\Delta^\alpha x$ depends on the nature and behavior of the sequence x and the value α , it has been observed that the properties such as linearity and exponent rules of the difference operator Δ^α are violating in certain particular cases. As a consequence of these violations, it is concluded that Theorems 1, 2 and 3 due to [4, 5] are not stable and need certain additional conditions in order to provide their substantial modifications.

The primary objective of this note is to study the convergence of the fractional difference sequences, the dynamic nature of the fractional difference operator Δ^α in detail and apply the same to modify Theorems 1, 2 and 3 of [4, 5]. Now, we analyze the convergence of the difference sequence $\Delta^\alpha x$ for different choice of α in detail, (i.e., $\alpha > 0, \alpha < 0$ and $\alpha \in \mathbb{N}$) by using the following theorems.

2 Main results

Theorem 1 The series defined in (3) is convergent for any $\alpha = n \in \mathbb{N}$ if the sequence $x = (x_k)$ is convergent. The converse of the statement may not hold in general.

Proof. Let $x = (x_k)$ be a convergent sequence. Then for given $\varepsilon > 0$, there exists a natural number N and real or complex number l such that, for every $k \geq N$, we have $|x_k - l| < \varepsilon$. Now, we have

$$\begin{aligned} |\Delta^n x_k| &= \left| \sum_{i=0}^n (-1)^i \binom{n}{i} x_{k+i} \right| \\ &= \left| \sum_{i=0}^n (-1)^i \binom{n}{i} x_{k+i} - \sum_{i=0}^n (-1)^i \binom{n}{i} l + \sum_{i=0}^n (-1)^i \binom{n}{i} l \right| \\ &= \left| \sum_{i=0}^n (-1)^i \binom{n}{i} (x_{k+i} - l) + l \sum_{i=0}^n (-1)^i \binom{n}{i} \right| \end{aligned}$$

$$\begin{aligned} &\leq \sum_{i=0}^n (-1)^i \binom{n}{i} |(x_{k+i} - l)| + |l| \left| \sum_{i=0}^n (-1)^i \binom{n}{i} \right| \\ &\leq \varepsilon \sum_{i=0}^n (-1)^i \binom{n}{i} = 0, \text{ for every } k \geq N. \end{aligned}$$

Therefore, $|\Delta^n x_k| \rightarrow 0$ as $k \rightarrow \infty$. For the converse part we take the following counter example:

For a natural number m , consider the sequence $x = (x_k)$, defined by $x_k = k^m$ for all $k \in \mathbb{N}_0$. Clearly, $x = (x_k)$ is divergent, but its associated difference sequence is

$$\begin{aligned} \Delta^n x_k &= \sum_{i=0}^n (-1)^i \binom{n}{i} (k+i)^m \\ &= k^m - \binom{n}{1} \left[k^m + \binom{m}{1} k^{m-1} + \binom{m}{2} + \cdots + \binom{m}{m} \right] \\ &\quad + \binom{n}{2} \left[k^m + 2 \binom{m}{1} k^{m-1} + 2^2 \binom{m}{2} + \cdots + \binom{m}{m} 2^m \right] + \cdots \\ &\quad + (-1)^n \binom{n}{n} \left[k^m + n \binom{m}{1} k^{m-1} + n^2 \binom{m}{2} + \cdots + \binom{m}{m} n^m \right] \\ &= k^m \left[1 - \binom{n}{1} + \binom{n}{2} - \binom{n}{3} + \cdots + (-1)^n \right] \\ &\quad + k^{m-1} \binom{m}{1} \left[-\binom{n}{1} + 2 \binom{n}{2} - 3 \binom{n}{3} + \cdots + n(-1)^n \right] \\ &\quad + k^{m-2} \binom{m}{2} \left[-\binom{n}{1} + 2^2 \binom{n}{2} - 3^2 \binom{n}{3} + \cdots + n^2(-1)^n \right] + \cdots \\ &\quad + k^{m-m} \binom{m}{m} \left[-\binom{n}{1} + 2^m \binom{n}{2} - 3^m \binom{n}{3} + \cdots + n^m(-1)^n \right] \\ &= \begin{cases} 0, & (n > m) \\ n!, & (n = m) \\ \infty, & (n < m) \end{cases}. \end{aligned}$$

Therefore, we conclude that for $n \geq m$ the difference sequence $(\Delta^n(k^m))_k$ is convergent while the primary sequence $x = (k^m)$ is divergent. \square

Theorem 2 *The series defined in (3) is convergent for any proper fraction $\alpha > 0$ if the sequence $x = (x_k)$ is convergent. The converse of the statement is true if the sequence involving infinite series*

$$\sum_{i=k}^{\infty} \binom{i-k+\alpha-1}{i-k} \Delta^{\alpha}(x_i) \text{ converges.} \quad (5)$$

Proof. The proof of the sufficient part is similar to that of Theorem (1), hence omitted.

For the necessary part we assume that the difference sequence $\Delta^{\alpha}x_k$ and the infinite series $\sum_{i=k}^{\infty} \binom{i-k+\alpha-1}{i-k} \Delta^{\alpha}(x_i)$ converge for all $k \in \mathbb{N}_0$. Let α be a proper fraction, i.e., $0 < \alpha < 1$. On simplifying (5), we obtain that

$$\begin{aligned} & \sum_{i=k}^{\infty} \binom{i-k+\alpha-1}{i-k} \Delta^{\alpha}(x_i) \\ &= \binom{\alpha-1}{0} \Delta^{\alpha}(x_k) + \binom{\alpha}{1} \Delta^{\alpha}(x_{k+1}) + \binom{\alpha+1}{2} \Delta^{\alpha}(x_{k+2}) + \dots \\ &= x_k - \binom{\alpha}{1} x_{k+1} + \binom{\alpha}{2} x_{k+2} - \binom{\alpha}{3} x_{k+3} + \dots \\ & \quad + \binom{\alpha}{1} \left[x_{k+1} - \binom{\alpha}{1} x_{k+2} + \binom{\alpha}{2} x_{k+3} - \binom{\alpha}{3} x_{k+4} + \dots \right] \\ & \quad + \binom{\alpha+1}{2} \left[x_{k+2} - \binom{\alpha}{1} x_{k+3} + \binom{\alpha}{2} x_{k+4} - \binom{\alpha}{3} x_{k+5} + \dots \right] + \dots \\ &= x_k. \end{aligned}$$

Thus, from the hypothesis, the sequence (x_k) is convergent. However, from Example 5, it is noticed that for a unbounded sequence $x = (x_k)$ with $x_k = k$ for all $k \in \mathbb{N}_0$, for a proper fraction α , corresponding difference sequence $\Delta^{\alpha}x_k \rightarrow \infty$, as $k \rightarrow \infty$. This completes the proof. \square

Theorem 3 *The series defined in (4) is convergent for any proper $\alpha > 0$ or $\alpha = n \in \mathbb{N}_0$ if the sequence $x = (x_k)$ is convergent with $x_k = \mathcal{O}(k^{-\alpha-1})$. The converse of the statement is true if the sequence involving infinite series*

$$\sum_{i=k}^{\infty} (-1)^{i-k} \binom{\alpha}{i-k} \Delta^{-\alpha}(x_i) \text{ converges.} \quad (6)$$

Proof. We know that the infinite series in (4) represents the inverse fractional difference sequence of the sequence (x_k) , thus it always suggests the idea analog

to integration or summation. Since the equation is a sum of infinite terms with all positive coefficients of x_k , most of the cases it gives ∞ even if the primary sequence is convergent. As a result, we need to consider strictly the order of the convergence of the primary sequence (x_k) in such a way that the final sum of the series (4) will be dominated.

Let us consider the convergent sequence $x = (x_k)$ with $x_k = \mathcal{O}(k^{-\alpha-1})$ and $\alpha > 0$. Then, there exists a constant M such that

$$\sup_k |x_k| \leq \frac{M}{k^{\alpha+1}}.$$

In fact, the above sequence is a null sequence and the corresponding inverse difference sequence is given below:

$$\begin{aligned} \Delta^{-\alpha} x_k &= \sum_{i=0}^{\infty} \frac{\alpha(\alpha+1)\dots(\alpha+i-1)}{\Gamma(i+1)} x_{k+i} \\ &= x_k + \alpha x_{k+1} + \frac{\alpha(\alpha+1)}{2!} x_{k+2} + \frac{\alpha(\alpha+1)(\alpha+1)}{3!} x_{k+3} + \dots \\ &\leq \frac{M}{k^{\alpha+1}} \left(1 + \alpha + \frac{\alpha(\alpha+1)}{2!} + \frac{\alpha(\alpha+1)(\alpha+1)}{3!} + \dots \right). \end{aligned}$$

The right hand side of the above equation is tending to 0 as $k \rightarrow \infty$. The equation contains two terms out of which the term $\frac{M}{k^{\alpha+1}}$ is dominating since it contains $(\alpha+1)$ as power of $1/k$ whereas other term contains α , only, which is a constant. It is rapidly tending to 0 as comparison to the rate at which the other term goes to ∞ . The converse part of this theorem is similar to that of Theorem 5. \square

Theorem 4 *Let $\alpha > 0$ be either a fraction or a natural number and $\Delta^\alpha : \mathcal{W} \rightarrow \mathcal{W}$ is a linear operator provided the series in (3) is convergent.*

Theorems (1), (2) and (3) can be verified in the light of the above theorem, it can be shown that most of the results are not satisfied in general.

Theorem 5 *For any proper fractions α, α_1 and α_2 , in general we have*

- (i) $\Delta^{\alpha_1}(\Delta^{\alpha_2} x_k) \neq \Delta^{\alpha_1+\alpha_2}(x_k)$ and $\Delta^{\alpha_2}(\Delta^{\alpha_1} x_k) \neq \Delta^{\alpha_1+\alpha_2}(x_k)$,
- (ii) $\Delta^\alpha(\Delta^{-\alpha} x_k) \neq x_k$ and $\Delta^{-\alpha}(\Delta^\alpha x_k) \neq x_k$,

Proof. We prove theorem by using suitable counter examples.

Example 5 Consider the sequence $x = (x_k)$, defined by $x_k = k$ for all $k \in \mathbb{N}_0$. Clearly it is a divergent sequence. Let us take $\alpha_1 = 1/2 = \alpha_2$ and therefore, $\alpha_1 + \alpha_2 = 1$. Then, we can calculate

$$\begin{aligned}\Delta^{\alpha_2} x_k &= (\Delta^{1/2} k)_k = k - \binom{1/2}{1}(k+1) + \binom{1/2}{2}(k+2) - \binom{1/2}{3}(k+3) + \dots \\ &= k \left[1 - \binom{1/2}{1} + \binom{1/2}{2} - \binom{1/2}{3} + \dots \right] \\ &\quad - \frac{1}{2} \left[1 - \binom{-1/2}{1} + \binom{-1/2}{2} - \binom{-1/2}{3} + \dots \right] \\ &= \infty.\end{aligned}$$

Now, $\Delta^{\alpha_1}(\Delta^{\alpha_2}(x_k)) = \Delta^{1/2}(\Delta^{1/2}(k)) = \Delta^{1/2}(\infty) = \infty$, but $\Delta^{\alpha_1+\alpha_2}(x_k) = \Delta^{1/2+1/2}(k) = \Delta(k) = k - (k+1) = -1$. Interchanging α_1 and α_2 in above expression we can prove the second condition. This completes the proof of Part (i) of Theorem 5.

Example 6 Let us consider the sequence $x = (x_k)$, defined by $x_k = r$ for all $k \in \mathbb{N}_0$ and $r \in \mathbb{R}$, the set of all real numbers. Clearly, $x = (x_k)$ is a convergent sequence. Taking $\alpha = 1/2$, we have

$$\begin{aligned}\Delta^{-\alpha} x_k &= (\Delta^{-1/2} r)_k = r \left[1 - \binom{-1/2}{1} + \binom{-1/2}{2} - \binom{-1/2}{3} + \dots \right] \\ &= \infty.\end{aligned}$$

Thus, the left hand side of 1st equation of Part (ii) is $\Delta^{\alpha}(\Delta^{-\alpha}(x_k)) = \Delta^{1/2}(\Delta^{-1/2}(r)) = \Delta^{1/2}(\infty) = \infty$, whereas the right hand side is $x_k = r$. Again by interchanging the positions of α and $-\alpha$, it is also noticed that

$$\begin{aligned}\Delta^{\alpha} x_k &= (\Delta^{1/2} r)_k = r \left[1 - \binom{1/2}{1} + \binom{1/2}{2} - \binom{1/2}{3} + \dots \right] \\ &= 0.\end{aligned}$$

Now, the left hand side of the second equation of Part (ii) can be found as $\Delta^{-\alpha}(\Delta^{\alpha}(x_k)) = \Delta^{-1/2}(\Delta^{1/2}(r)) = \Delta^{-1/2}(0) = 0$ which is not equal to the right hand side i.e., $x_k = r$. This completes the proof of Part (ii) of Theorem 5.

□

Above examples conclude that linearity and exponent rules involving the fractional difference operator Δ^{α} for any sequence in w are not uniformly

posed. Eventually, these rules are deviating due to lack of convergence of related infinite series. In fact, the convergence of the related infinite series is completely depending on the nature of the primary sequence (x_k) and the choice of the values of α . It is understood that if the primary sequence (x_k) and the value α are suitably chosen then obviously, this deviation can be restricted to a given domain. This idea suggests that Theorems 1, 2 and 3 of [4] need relevant modifications and the modified results are as follows.

Theorem 6 *For any positive proper fractions α, α_1 and α_2 , we have*

(i) *Let the sequence $x = (x_k)$ be convergent, then*

$$\Delta^{\alpha_1}(\Delta^{\alpha_2}(x_k)) = \Delta^{\alpha_1+\alpha_2}(x_k) = \Delta^{\alpha_2}(\Delta^{\alpha_1}(x_k)),$$

(ii) *Let the sequence $(\Delta^{-\alpha}x_k)$ be convergent, then*

$$\Delta^{\alpha}(\Delta^{-\alpha}x_k) = x_k,$$

(iii) *Let the sequence $(\Delta^{\alpha}x_k)$ be of $\mathcal{O}(k^{-\alpha-1})$, then*

$$\Delta^{-\alpha}(\Delta^{\alpha}x_k) = x_k.$$

Combining all points, Theorem 6 can be restated as follows:

Remark 1 *Let $\alpha > 0$ and β be a real such that $\alpha + \beta > 0$ and the sequence (x_k) be of $\mathcal{O}(k^{-m-1})$, where $m = \min(|\alpha|, |\beta|)$, then*

$$\Delta^{\alpha}(\Delta^{\beta}(x_k)) = \Delta^{\alpha+\beta}(x_k) = \Delta^{\beta}(\Delta^{\alpha}(x_k)).$$

Corollary 1 *For any $n \in \mathbb{N}$, let Δ^{-n} be the negative integral difference operator, then*

(i) $\Delta^{-1}(x_k) = \sum_{i=1}^{\infty} x_{k+i}$, if the sequence (x_k) is convergent with $x_k = \mathcal{O}(k^{-2})$,

(ii) $\Delta^{-2}(x_k) = \sum_{i=1}^{\infty} (i+1)x_{k+i}$, if the sequence (x_k) is convergent with $x_k = \mathcal{O}(k^{-3})$,

(iii) $\Delta^{-3}(x_k) = \sum_{i=1}^{\infty} (s_i)x_{k+i}$, where $s_i = \sum_{j=1}^i j$, if the sequence (x_k) is convergent with $x_k = \mathcal{O}(k^{-4})$.

To next, we discuss some operator norms involving the difference operator of fractional order.

Let $A = (a_{nk})$ be an infinite matrix with $a_{nk} \geq 0$ for all $n, k \in \mathbb{N}_0$. Then we have the following theorems on operator norms via the infinite matrix A :

Theorem 7 *Let $X \in \{c_0, c, \ell_\infty\}$. Then the infinite matrix A is a bounded operator from X to $X(\Delta^\alpha)$ if*

$$\mathcal{M} = \sup_n \left\{ \sum_{k=0}^{\infty} \left| \sum_{i=0}^{\infty} \frac{(-\alpha)(-\alpha+1)\dots(-\alpha+i-1)}{\Gamma(i+1)} a_{n+i,k} \right| \right\} < \infty$$

and

$$\|A\|_{(\infty, \Delta^\alpha)} = \mathcal{M}.$$

Proof. Suppose $X = \ell_\infty$ and $x \in X$. Then, we have

$$\begin{aligned} \|Ax\|_{(\infty, \Delta^\alpha)} &= \sup_n \left| \sum_{k=0}^{\infty} \sum_{i=0}^{\infty} \frac{(-\alpha)(-\alpha+1)\dots(-\alpha+i-1)}{\Gamma(i+1)} a_{n+i,k} x_k \right| \\ &\leq \sup_n \left\{ \sum_{k=0}^{\infty} \left| \sum_{i=0}^{\infty} \frac{(-\alpha)(-\alpha+1)\dots(-\alpha+i-1)}{\Gamma(i+1)} a_{n+i,k} x_k \right| \right\} \\ &\leq \mathcal{M} \|x\|_\infty. \end{aligned}$$

Also, for $x = e = (1, 1, 1, \dots)$, we have

$$\begin{aligned} \|Ae\|_{(\infty, \Delta^\alpha)} &= \sup_n \left| \sum_{k=0}^{\infty} \sum_{i=0}^{\infty} \frac{(-\alpha)(-\alpha+1)\dots(-\alpha+i-1)}{\Gamma(i+1)} a_{n+i,k} \right| \\ &= \sup_n \left\{ \sum_{k=0}^{\infty} \left| \sum_{i=0}^{\infty} \frac{(-\alpha)(-\alpha+1)\dots(-\alpha+i-1)}{\Gamma(i+1)} a_{n+i,k} \right| \right\} \\ &= \mathcal{M}. \end{aligned}$$

This proves the result. □

Theorem 8 *The infinite matrix A is a bounded operator from ℓ_1 to $\ell_1(\Delta^\alpha)$ if*

$$\overline{\mathcal{M}} = \sup_k \left\{ \sum_{n=0}^{\infty} \left| \sum_{i=0}^{\infty} \frac{(-\alpha)(-\alpha+1)\dots(-\alpha+i-1)}{\Gamma(i+1)} a_{n+i,k} \right| \right\} < \infty,$$

and

$$\|A\|_{(1, \Delta^\alpha)} = \overline{\mathcal{M}}.$$

Proof. Suppose that $x \in \ell_1$ and A be an infinite matrix, then

$$\begin{aligned} \|Ax\|_{(1,\Delta^\alpha)} &= \sum_{n=0}^{\infty} \left| \sum_{k=0}^{\infty} \sum_{i=0}^{\infty} \frac{(-\alpha)(-\alpha+1)\dots(-\alpha+i-1)}{\Gamma(i+1)} a_{n+i,k} x_k \right| \\ &\leq \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \left| \sum_{i=0}^{\infty} \frac{(-\alpha)(-\alpha+1)\dots(-\alpha+i-1)}{\Gamma(i+1)} a_{n+i,k} x_k \right| \\ &\leq \sum_{k=0}^{\infty} \sum_{n=0}^{\infty} \left| \sum_{i=0}^{\infty} \frac{(-\alpha)(-\alpha+1)\dots(-\alpha+i-1)}{\Gamma(i+1)} a_{n+i,k} \right| |x_k| \\ &\leq \overline{\mathcal{M}} \|x\|_1. \end{aligned}$$

Now, for the sequence $x = e^{(m)}$ (having 1 at m -th place and 0 otherwise), one can get

$$\begin{aligned} \|Ae^{(m)}\|_{(1,\Delta^\alpha)} &= \sum_{n=0}^{\infty} \left| \sum_{k=0}^{\infty} \sum_{i=0}^{\infty} \frac{(-\alpha)(-\alpha+1)\dots(-\alpha+i-1)}{\Gamma(i+1)} a_{n+i,k} x_k \right| \\ &= \sum_{n=0}^{\infty} \left| \sum_{i=0}^{\infty} \frac{(-\alpha)(-\alpha+1)\dots(-\alpha+i-1)}{\Gamma(i+1)} a_{n+i,m} \right| \\ &= \overline{\mathcal{M}}. \end{aligned}$$

This concludes the proof. \square

Theorem 9 *The infinite matrix A is a bounded operator from ℓ_p , ($1 \leq p < \infty$) to $\ell_p(\Delta^\alpha)$ if*

$$\overline{\mathcal{M}}_p = \sup_k \left\{ \sum_{n=0}^{\infty} \left| \sum_{i=0}^{\infty} \frac{(-\alpha)(-\alpha+1)\dots(-\alpha+i-1)}{\Gamma(i+1)} a_{n+i,k} \right|^p \right\} < \infty,$$

and

$$\|A\|_{(p,\Delta^\alpha)}^p = \overline{\mathcal{M}}_p.$$

Proof. This follows from the proof of Theorem 8. \square

Theorem 10 *The identity matrix I is a bounded operator from X to $X(\Delta^\alpha)$ for $X \in \{c, c_0, \ell_\infty, \ell_1\}$ and*

$$\|I\|_{(\infty,\Delta^\alpha)} = \|I\|_{(1,\Delta^\alpha)} = 2^\alpha.$$

Proof. Suppose the infinite matrix $A = I$, then from Theorem 7, we can write

$$\begin{aligned}\mathcal{M}_n &= \sum_{k=0}^{\infty} \left| \sum_{i=0}^{\infty} \frac{(-\alpha)(-\alpha+1)\dots(-\alpha+i-1)}{\Gamma(i+1)} a_{n+i,k} \right| \\ &= \sum_{k=n}^{\infty} \left| \frac{(-\alpha)(-\alpha+1)\dots(-\alpha+k-n-1)}{\Gamma(k-n+1)} \right|.\end{aligned}$$

Therefore, we have

$$\|I\|_{(\infty, \Delta^\alpha)} = \sup_n \mathcal{M}_n = 2^\alpha.$$

Similarly, using Theorem 8, one can prove $\|I\|_{(1, \Delta^\alpha)} = 2^\alpha$. \square

Conclusion

We have investigated some idea on the convergence of difference sequence for fractional-order which may be very similar to that of integer orders but most of the cases they are nonuniform and dynamic in nature. As an application of this idea, some existing results in the literature have been modified. Certain operator norms involving the difference operator of fractional order is determined.

In the next study, we will extend this idea to the case of the statistical convergence of difference sequence and study the variations in the cases of integer and fractional orders.

References

- [1] A. Alotaibi, M. Mursaleen, B. A. S. Alamri, S. A. Mohiuddine, Compact operators on some Fibonacci difference sequence spaces, *J. Inequal. Appl.*, (2015), 2015–203.
- [2] B. Altay, F. Başar, The fine spectrum and the matrix domain the difference operator Δ on the sequence space ℓ_p , ($0 < p < 1$), *Commun. Math. Anal.*, **2** (2) (2007), 1–11.
- [3] C. Aydın, F. Başar, Some new difference sequence spaces, *Appl. Math. Comput.*, **157** (3) (2004), 677–693.
- [4] P. Baliarsingh, Some new difference sequence spaces of fractional order and their dual spaces, *Appl. Math. Comput.*, **219** (18) (2013), 9737–9742.

- [5] P. Baliarsingh, S. Dutta, On the classes of fractional order difference sequence spaces and their matrix transformations, *Appl. Math. Comput.*, **250** (2015), 665–674.
- [6] P. Baliarsingh, S. Dutta, A unifying approach to the difference operators and their applications, *Bol. Soc. Parana. Mat.*, **33** (1) (2015), 49–57.
- [7] P. Baliarsingh, S. Dutta, On a spectral classification of the operator Δ_v^r over the Sequence Space c_0 , *Proc. Natl. Acad. Sci., India, Sect. A Phys. Sci.*, **84** (4) (2014) 555–561.
- [8] P. Baliarsingh, On a fractional difference operator, *Alexandria Eng. J.*, **55** (2) (2016), 1811–1816.
- [9] P. Baliarsingh, L. Nayak, A note fractional difference operators, *Alexandria Eng. J.*, **57** (2) (2018), 1051–1054.
- [10] P. Baliarsingh, On certain dynamic properties of difference sequences and the fractional derivatives, *Math. Metho. Appl. Sci.*, (2020) doi:10.1002/mma.6417.
- [11] F. Başar, *Summability theory and its applications*, Bentham Science Publishers, e-books, Monographs, İstanbul, 2012.
- [12] F. Başar, M. Kirişçi, Almost convergence and generalized difference matrix, *Comput. Math. Appl.*, **61** (3) (2011), 602–611.
- [13] F. Başar, B. Altay, On the space of sequences of p -bounded variation and related matrix mappings, (English, Ukrainian summary) *Ukrain. Mat. Zh.*, **55** (1) (2003), 108–118; reprinted in *Ukrainian Math. J.*, **55** (1) (2003), 136–147.
- [14] M. Basarır, E. E. Kara, On some difference sequence spaces of weighted mean and compact operators, *Ann. Funct. Anal.*, **2** (2) (2011), 114–129.
- [15] S. Dutta, P. Baliarsingh, On the fine spectra of the generalized r th difference operator Δ_v^r on the sequence space ℓ_1 , *Appl. Math. Comput.*, **219** (2012), 1776–1784.
- [16] S. Dutta, P. Baliarsingh, On the spectrum of 2-nd order generalized difference operator Δ^2 over the sequence space c_0 , *Bol. Soc. Paran. Mat.* **31** (2) (2013), 235–244.

- [17] M. Et, R. Çolak, On some generalized difference sequence spaces, *Soochow J. Math.*, **21** (4) (1995), 377–386.
- [18] U. Kadak, P. Baliarsingh, On certain Euler difference sequence spaces of fractional order and related dual properties, *J. Nonlinear Sci. Appl.*, **8** (2015), 997–1004.
- [19] M. Kirişçi, F. Başar, Some new sequence spaces derived by the domain of generalized difference matrix, *Comput. Math. Appl.*, **60** (5) (2010), 1299–1309.
- [20] H. Kızmaz, On Certain Sequence spaces, *Canad. Math. Bull.*, **24** (2) (1981) 169–176.
- [21] E. Malkowsky, M. Mursaleen, S. Suantai, The dual spaces of sets of difference sequences of order m and matrix transformations, *Acta Math. Sin. (Engl. Ser.)*, **23** (3) (2007), 521–532.
- [22] S. A. Mohiuddine, B. Hazarika, Some classes of ideal convergent sequences and generalized difference matrix operator, *Filomat*, **31** (6) (2017), 1827–1834.
- [23] M. Mursaleen, A. K. Noman, On some new difference sequence spaces of non-absolute type, *Math. Comput. Modelling*, **52** (2010), 603–617.
- [24] M. Mursaleen, A. K. Noman, Compactness of matrix operators on some new difference sequence spaces, *Linear Algebra Appl.*, **436** (1) (2012), 41–52.
- [25] M. Mursaleen, V. Karakaya, H. Polat, N. Simsek, Measure of noncompactness of matrix operators on some difference sequence spaces of weighted means, *Comput. Math. Appl.*, **62** (2011), 814–820.
- [26] B. C. Tripathy, On a New class of sequences, *Demonstratio Math.*, **37** (2) (2004), 377– 381.
- [27] B. C. Tripathy, S. Mahanta, On a class of difference sequences related to the ℓ^p space defined by Orlicz functions, *Math. Slovaca*, **57** (2) (2007), 171–178.
- [28] B. C. Tripathy, A. Paul, The spectrum of the operator $D(r, 0, s, 0, t)$ over the sequence spaces c_0 and c , *Journal Math.* , **2013**, (2013), Article ID 430965.

- [29] B. C. Tripathy, A. Paul, The spectrum of the operator $D(r, 0, 0, s)$ over the sequence space c_0 and c , *Kyungpook Math. Journal*, **53** (2) (2013), 247–256.
- [30] B. C. Tripathy, A. Paul, The spectrum of the operator $D(r, 0, 0, s)$ over the sequence spaces ℓ_p and bv_p , *Hacettepe J. Math. Stat.*, **43** (3) (2014), 425–434.

Received: March 5, 2020

On some properties of split Horadam quaternions

Dorota Bród

Rzeszow University of Technology,
Faculty of Mathematics and Applied Physics,
al. Powstańców Warszawy 12,
35-959 Rzeszów, Poland
email: dorotab@prz.edu.pl

Abstract. In this paper we introduce and study the split Horadam quaternions. We give some identities, among others Binet's formula, Catalan's, Cassini's and d'Ocagne's identities for these numbers.

1 Introduction

Let \mathbb{C} be the field of complex numbers. A quaternion x is a hyper-complex number represented by

$$\mathbb{H} = \{x = a_0 + a_1\mathbf{i} + a_2\mathbf{j} + a_3\mathbf{k} : a_s \in \mathbb{R}, s = 0, 1, 2, 3\},$$

where $\{1, \mathbf{i}, \mathbf{j}, \mathbf{k}\}$ is an orthonormal basis in \mathbb{R}^4 , which satisfies the quaternion multiplication rules:

$$\mathbf{i}^2 = \mathbf{j}^2 = \mathbf{k}^2 = \mathbf{ijk} = -1,$$

$$\mathbf{ij} = \mathbf{k} = -\mathbf{ji}, \mathbf{jk} = \mathbf{i} = -\mathbf{kj}, \mathbf{ki} = \mathbf{j} = -\mathbf{ik}.$$

The quaternions were introduced by W. R. Hamilton in 1843.

2010 Mathematics Subject Classification: 11B37, 11R52

Key words and phrases: Horadam numbers, quaternion, split quaternion, split Horadam quaternion

Another extension of the complex numbers is the algebra of split quaternions. The split quaternions were introduced by J. Cockle in 1849 [2]. The set of split (or coquaternions) can be represented as

$$\hat{\mathbb{H}} = \{y = b_0 + b_1i + b_2j + b_3k : b_s \in \mathbb{R}, s = 0, 1, 2, 3\},$$

where $\{1, i, j, k\}$ is the basis of $\hat{\mathbb{H}}$ satisfying the following equalities

$$i^2 = -j^2 = -k^2 = -1, \quad (1)$$

$$ij = k = -ji, \quad jk = -i = -kj, \quad ki = j = -ik. \quad (2)$$

The split quaternion can be rewritten as

$$y = (b_0 + b_1i) + (b_2 + b_3i)j = z_1 + z_2j, \quad z_1, z_2 \in \mathbb{C}.$$

The split quaternions contain nontrivial zero divisors, nilpotent elements and idempotents. The conjugate of a split quaternion $y = b_0 + b_1i + b_2j + b_3k$, denoted by \bar{y} , is given by $\bar{y} = b_0 - b_1i - b_2j - b_3k$. The norm of y is defined as

$$N(y) = y\bar{y} = b_0^2 + b_1^2 - b_2^2 - b_3^2. \quad (3)$$

Let $y_1, y_2 \in \hat{\mathbb{H}}$, $y_1 = a_1 + b_1i + c_1j + d_1k$, $y_2 = a_2 + b_2i + c_2j + d_2k$. Then addition and subtraction of the split quaternions is defined as follows

$$y_1 \pm y_2 = (a_1 \pm a_2) + (b_1 \pm b_2)i + (c_1 \pm c_2)j + (d_1 \pm d_2)k.$$

Multiplication of the split quaternions is defined by

$$\begin{aligned} y_1 \cdot y_2 = & a_1a_2 - b_1b_2 + c_1c_2 + d_1d_2 + (a_1b_2 + b_1a_2 - c_1d_2 + d_1c_2)i \\ & + (a_1c_2 + c_1a_2 - b_1d_2 + d_1b_2)j + (a_1d_2 + d_1a_2 + b_1c_2 - c_1b_2)k. \end{aligned} \quad (4)$$

For the basics on split quaternions theory, see [5].

2 The Horadam numbers

In [3] Horadam introduced a sequence $\{W_n\}$ defined by the following relation

$$W_0 = a, W_1 = b, W_n = pW_{n-1} + qW_{n-2} \text{ for } n \geq 2 \quad (5)$$

for arbitrary $a, b, p, q \in \mathbb{Z}$. This sequence is a certain generalization of famous sequences such as Fibonacci sequence $\{F_n\}$ ($a = 0, b = 1, p = q = 1$), Lucas

sequence $\{L_n\}$ ($a = 2, b = 1, p = q = 1$), Jacobsthal sequence $\{J_n\}$ ($a = 0, b = 1, p = 1, q = 2$), Pell sequence $\{P_n\}$ ($a = 0, b = 1, p = 2, q = 1$), Pell-Lucas sequence $\{PL_n\}$ ($a = b = 1, p = 2, q = 1$). The sequences defined by (5) are called sequences of the Fibonacci type.

The characteristic equation associated with the recurrence (5) is

$$r^2 - pr - q = 0.$$

Assuming that $p^2 + 4q > 0$, the equation has the following roots

$$r_1 = \frac{p + \sqrt{p^2 + 4q}}{2}, \quad r_2 = \frac{p - \sqrt{p^2 + 4q}}{2}. \quad (6)$$

Note that

$$r_1 + r_2 = p, \quad (7)$$

$$r_1 - r_2 = \sqrt{p^2 + 4q}, \quad (8)$$

$$r_1 r_2 = -q. \quad (9)$$

The Binet's formula for the sequence $\{W_n\}$ has the following form

$$W_n = \frac{(b - ar_2)r_1^n - (b - ar_1)r_2^n}{r_1 - r_2}.$$

Let

$$\alpha = \frac{b - ar_2}{r_1 - r_2}, \quad \beta = \frac{b - ar_1}{r_1 - r_2}. \quad (10)$$

Then

$$W_n = \alpha r_1^n - \beta r_2^n. \quad (11)$$

In the next section we will use the following result.

Theorem 1 *Let n, p, q be integers such that $n \geq 0, p^2 + 4q > 0$. Then*

$$\sum_{l=0}^{n-1} W_l = \frac{W_n + qW_{n-1} + a(p-1) - b}{p + q - 1}. \quad (12)$$

Proof. Using formula (11), (7) and (9), we get

$$\sum_{l=0}^{n-1} W_l = \sum_{l=0}^{n-1} (\alpha r_1^l - \beta r_2^l) = \alpha \frac{1 - r_1^n}{1 - r_1} - \beta \frac{1 - r_2^n}{1 - r_2}$$

$$\begin{aligned}
&= \frac{\alpha - \beta - (\alpha r_2 - \beta r_1) - (\alpha r_1^n - \beta r_2^n) + r_1 r_2 (\alpha r_1^{n-1} - \beta r_2^{n-1})}{1 - (r_1 + r_2) + r_1 r_2} \\
&= \frac{\alpha - \beta - (\alpha r_2 - \beta r_1) - W_n - q W_{n-1}}{1 - p - q}.
\end{aligned}$$

By simple calculations we have $\alpha - \beta = a$, $\alpha r_2 - \beta r_1 = ap - b$. Hence

$$\sum_{l=0}^{n-1} W_l = \frac{W_n + q W_{n-1} + a(p-1) - b}{p + q - 1}.$$

□

Numbers of the Fibonacci type appear in many subjects of mathematics. In [4] Horadam defined the Fibonacci and Lucas quaternions. In [1] the split Fibonacci quaternions Q_n and split Lucas quaternions T_n were introduced by the following relations

$$\begin{aligned}
Q_n &= F_n + iF_{n+1} + jF_{n+2} + kF_{n+3}, \\
T_n &= L_n + iL_{n+1} + jL_{n+2} + kL_{n+3},
\end{aligned}$$

where F_n , L_n is n th Fibonacci and Lucas number, resp. and $\{i, j, k\}$ is the standard basis of split quaternions. In the literature there are many generalizations of the Fibonacci and Lucas sequences, among others k -Fibonacci sequence $\{F_{k,n}\}$, k -Lucas sequence $\{L_{k,n}\}$, defined for $k \in \mathbb{N}$ in the following way

$$\begin{aligned}
F_{k,0} &= 0, F_{k,1} = 1, F_{k,n} = kF_{k,n-1} + F_{k,n-2} \text{ for } n \geq 2, \\
L_{k,0} &= 2, L_{k,1} = k, L_{k,n} = kL_{k,n-1} + L_{k,n-2} \text{ for } n \geq 2.
\end{aligned}$$

Some interesting results for the split k -Fibonacci and split k -Lucas quaternions can be found in [6]. In [7] the authors studied split Pell quaternions SP_n and split Pell-Lucas quaternions SPL_n defined by

$$\begin{aligned}
SP_n &= P_n + iP_{n+1} + jP_{n+2} + kP_{n+3}, \\
SPL_n &= PL_n + iPL_{n+1} + jPL_{n+2} + kPL_{n+3},
\end{aligned}$$

where P_n and PL_n is n th Pell and Pell-Lucas number, resp.

We will focus on split Horadam quaternions. We will present some identities for the split Horadam quaternions, which generalize the results for the split Fibonacci quaternions, the split Lucas quaternions, the split Pell quaternions and the split Pell-Lucas quaternions.

3 The split Horadam quaternions

For $n \geq 0$ define the split Horadam quaternion H_n by

$$H_n = W_n + iW_{n+1} + jW_{n+2} + kW_{n+3}, \quad (13)$$

where W_n is the n th Horadam number and i, j, k are split quaternionic units which satisfy the multiplication rules given by (1) and (2).

By (5) and (13) we obtain

$$\begin{aligned} H_0 &= a + bi + j(pb + qa) + k(p^2b + pqa + qb) \\ H_1 &= b + i(pb + qa) + j(p^2b + pqa + qb) + k(p^3b + p^2qa + 2pqb + q^2a) \\ H_2 &= pb + qa + i(p^2b + pqa + qb) + j(p^3b + p^2qa + 2pqb + q^2a) \\ &\quad + k(p^4b + p^3qa + 2pq(pb + qa) + p^2qb + q^2b). \end{aligned} \quad (14)$$

For any $n \geq 0$ we obtain the norm of H_n .

Proposition 1 *Let n, p, q be integers such that $n \geq 0$, $p^2 + 4q > 0$. Then*

$$\begin{aligned} N(H_n) &= (1 - q^2 - p^2q^2)W_n^2 + (1 - p^2 - (p^2 + q^2)^2)W_{n+1}^2 \\ &\quad - 2pq(1 + p^2 + q)W_nW_{n+1}. \end{aligned}$$

Proof. Using formula (3) and (13), we get

$$\begin{aligned} N(H_n) &= W_n^2 + W_{n+1}^2 - W_{n+2}^2 - W_{n+3}^2 \\ &= W_n^2 + W_{n+1}^2 - (pW_{n+1} + qW_n)^2 - \left((p^2 + q)W_{n+1} + pqW_n\right)^2 \\ &= W_n^2 + W_{n+1}^2 - (p^2W_{n+1}^2 + 2pqW_nW_{n+1} + q^2W_n^2) \\ &\quad - ((p^2 + q)^2W_{n+1}^2 + 2pq(p^2 + q)W_nW_{n+1} + p^2q^2W_n^2). \end{aligned}$$

By simple calculations we get the result. □

By (13) we get a recurrence relation for the split Horadam quaternions.

Proposition 2 *Let n, p, q be integers such that $n \geq 2$, $p^2 + 4q > 0$. Then*

$$H_n = pH_{n-1} + qH_{n-2},$$

where H_0, H_1 are given by (14).

Proof. By formula (13) and (5) we get

$$\begin{aligned}
 pH_{n-1} + qH_{n-2} &= p(W_{n-1} + iW_n + jW_{n+1} + kW_{n+2}) \\
 &\quad + q(W_{n-2} + iW_{n-1} + jW_n + kW_{n+1}) \\
 &= pW_{n-1} + qW_{n-2} + i(pW_n + qW_{n-1}) \\
 &\quad + j(pW_{n+1} + qW_n) + k(pW_{n+2} + qW_{n+1}) \\
 &= W_n + iW_{n+1} + jW_{n+2} + kW_{n+3} = H_n,
 \end{aligned}$$

which ends the proof. \square

Theorem 2 Let n, p, q be integers such that $n \geq 0, p^2 + 4q > 0$. Then

$$(i) \ H_n + \overline{H_n} = 2W_n,$$

$$(ii) \ N(H_n) = 2W_n H_n - H_n^2.$$

Proof. (i) Using the definition of the conjugate of a split quaternion we obtain the result.

(ii) By formula (13) we have

$$\begin{aligned}
 H_n^2 &= W_n^2 - W_{n+1}^2 + W_{n+2}^2 + W_{n+3}^2 \\
 &\quad + 2iW_n W_{n+1} + 2jW_n W_{n+2} + 2kW_n W_{n+3} \\
 &= -W_n^2 - W_{n+1}^2 + W_{n+2}^2 + W_{n+3}^2 \\
 &\quad + 2(W_n^2 + iW_n W_{n+1} + jW_n W_{n+2} + kW_n W_{n+3}) \\
 &= 2W_n(W_n + iW_{n+1} + jW_{n+2} + kW_{n+3}) \\
 &\quad - W_n^2 - W_{n+1}^2 + W_{n+2}^2 + W_{n+3}^2 \\
 &= 2W_n H_n - N(H_n).
 \end{aligned}$$

Hence we get the result. \square

The next theorem presents the Binet's formula for the split Horadam quaternions.

Theorem 3 (Binet's formula) Let n, p, q be integers such that $n \geq 0, p^2 + 4q > 0$. Then

$$H_n = \alpha \hat{r}_1 r_1^n - \beta \hat{r}_2 r_2^n, \quad (15)$$

where r_1, r_2, α, β are given by (6), (10), resp. and $\hat{r}_1 = 1 + ir_1 + jr_1^2 + kr_1^3$, $\hat{r}_2 = 1 + ir_2 + jr_2^2 + kr_2^3$.

Proof. By (11) we have

$$\begin{aligned}
 H_n &= W_n + iW_{n+1} + jW_{n+2} + kW_{n+3} \\
 &= \alpha r_1^n - \beta r_2^n + i(\alpha r_1^{n+1} - \beta r_2^{n+1}) + j(\alpha r_1^{n+2} - \beta r_2^{n+2}) \\
 &\quad + k(\alpha r_1^{n+3} - \beta r_2^{n+3}) \\
 &= \alpha r_1^n (1 + ir_1 + jr_1^2 + kr_1^3) - \beta r_2^n (1 + ir_2 + jr_2^2 + kr_2^3) \\
 &= \alpha \hat{r}_1 r_1^n - \beta \hat{r}_2 r_2^n.
 \end{aligned}$$

□

Using the Binet's formula (15), we can obtain some new identities for the split Horadam quaternions. We will use the following lemma.

Lemma 1 *Let $\hat{r}_1 = 1 + ir_1 + jr_1^2 + kr_1^3$, $\hat{r}_2 = 1 + ir_2 + jr_2^2 + kr_2^3$, where r_1, r_2 are given by (6). Then*

$$\begin{aligned}
 \hat{r}_1 \hat{r}_2 &= 1 + q + q^2 - q^3 + i(p + q^2 \sqrt{p^2 + 4q}) \\
 &\quad + j(p^2 + 2q - pq \sqrt{p^2 + 4q}) + k(p^3 + 3pq + q \sqrt{p^2 + 4q}), \quad (16)
 \end{aligned}$$

$$\begin{aligned}
 \hat{r}_2 \hat{r}_1 &= 1 + q + q^2 - q^3 + i(p - q^2 \sqrt{p^2 + 4q}) \\
 &\quad + j(p^2 + 2q + pq \sqrt{p^2 + 4q}) + k(p^3 + 3pq - q \sqrt{p^2 + 4q}). \quad (17)
 \end{aligned}$$

Proof. Using formula (4), we have

$$\begin{aligned}
 \hat{r}_1 \hat{r}_2 &= 1 - r_1 r_2 + (r_1 r_2)^2 + (r_1 r_2)^3 + i(r_1 + r_2 + (r_1 r_2)^2(r_1 - r_2)) \\
 &\quad + j(r_1^2 + r_2^2 + r_1 r_2(r_1^2 - r_2^2)) + k(r_1^3 + r_2^3 - r_1 r_2(r_1 - r_2)), \\
 \hat{r}_2 \hat{r}_1 &= 1 - r_1 r_2 + (r_1 r_2)^2 + (r_1 r_2)^3 + i(r_1 + r_2 - (r_1 r_2)^2(r_1 - r_2)) \\
 &\quad + j(r_1^2 + r_2^2 - r_1 r_2(r_1^2 - r_2^2)) + k(r_1^3 + r_2^3 + r_1 r_2(r_1 - r_2)).
 \end{aligned}$$

By (7) and (9) we get

$$\begin{aligned}
 r_1^2 + r_2^2 &= (r_1 + r_2)^2 - 2r_1 r_2 = p^2 + 2q, \\
 r_1^3 + r_2^3 &= (r_1 + r_2)^3 - 3r_1 r_2(r_1 + r_2) = p^3 + 3pq.
 \end{aligned}$$

Hence

$$\begin{aligned}
 \hat{r}_1 \hat{r}_2 &= 1 + q + q^2 - q^3 + i(p + q^2 \sqrt{p^2 + 4q}) \\
 &\quad + j(p^2 + 2q - pq \sqrt{p^2 + 4q}) + k(p^3 + 3pq + q \sqrt{p^2 + 4q}),
 \end{aligned}$$

$$\begin{aligned}\hat{r}_1\hat{r}_1 &= 1 + q + q^2 - q^3 + i(p - q^2\sqrt{p^2 + 4q}) \\ &\quad + j(p^2 + 2q + pq\sqrt{p^2 + 4q}) + k(p^3 + 3pq - q\sqrt{p^2 + 4q}).\end{aligned}$$

□

Corollary 1

$$\hat{r}_1\hat{r}_2 + \hat{r}_2\hat{r}_1 = 2(1 + q + q^2 - q^3 + pi + j(p^2 + 2q) + k(p^3 + 3pq)). \quad (18)$$

Theorem 4 (*Catalan's identity*) Let n, m, p, q be integers such that $n \geq m$, $p^2 + 4q > 0$. Then

$$H_{n-m}H_{n+m} - H_n^2 = \alpha\beta(-q)^{n-m}[(-q)^m(\hat{r}_1\hat{r}_2 + \hat{r}_2\hat{r}_1) - r_2^{2m}\hat{r}_1\hat{r}_2 - r_1^{2m}\hat{r}_2\hat{r}_1],$$

where $\alpha, \beta, \hat{r}_1\hat{r}_2 + \hat{r}_2\hat{r}_1, \hat{r}_1\hat{r}_2, \hat{r}_2\hat{r}_1$ are given by (10), (18), (16), (17), resp.

Proof. By (15) we get

$$\begin{aligned}H_{n-m}H_{n+m} - H_n^2 &= (\alpha\hat{r}_1r_1^{n-m} - \beta\hat{r}_2r_2^{n-m})(\alpha\hat{r}_1r_1^{n+m} - \beta\hat{r}_2r_2^{n+m}) \\ &\quad - (\alpha\hat{r}_1r_1^n - \beta\hat{r}_2r_2^n)(\alpha\hat{r}_1r_1^n - \beta\hat{r}_2r_2^n) \\ &= \alpha\beta(r_1r_2)^{n-m}[(r_1r_2)^m(\hat{r}_1\hat{r}_2 + \hat{r}_2\hat{r}_1) \\ &\quad - r_2^{2m}\hat{r}_1\hat{r}_2 - r_1^{2m}\hat{r}_2\hat{r}_1].\end{aligned}$$

Using formula (9), we obtain

$$H_{n-m}H_{n+m} - H_n^2 = \alpha\beta(-q)^{n-m} \left((-q)^m(\hat{r}_1\hat{r}_2 + \hat{r}_2\hat{r}_1) - r_2^{2m}\hat{r}_1\hat{r}_2 - r_1^{2m}\hat{r}_2\hat{r}_1 \right).$$

□

Corollary 2 (*Cassini's identity*) Let n, p, q be integers such that $n \geq 0$, $p^2 + 4q > 0$. Then

$$H_{n-1}H_{n+1} - H_n^2 = -\alpha\beta(-q)^{n-1} \left(q(\hat{r}_1\hat{r}_2 + \hat{r}_2\hat{r}_1) + r_2^2\hat{r}_1\hat{r}_2 + r_1^2\hat{r}_2\hat{r}_1 \right).$$

Note that for $p = q = 1$ we get the Cassini's identity for the split Fibonacci quaternions Q_n and the split Lucas quaternions T_n ([1]).

Corollary 3 Let $n \geq 1$ be an integer. Then

$$(i) \quad Q_{n-1}Q_{n+1} - Q_n^2 = (-1)^n(2Q_1 - 2i - 3k),$$

$$(ii) \quad T_{n-1}T_{n+1} - T_n^2 = 5(-1)^{n+1}(2Q_1 - 2i - 3k).$$

Proof. (i) Using Lemma 1, for $p = q = 1$ we get

$$\begin{aligned}\hat{r}_1\hat{r}_2 &= 2 + (1 + \sqrt{5})i + (3 - \sqrt{5})j + (4 + \sqrt{5})k, \\ \hat{r}_2\hat{r}_1 &= 2 + (1 - \sqrt{5})i + (3 + \sqrt{5})j + (4 - \sqrt{5})k, \\ \hat{r}_1\hat{r}_2 + \hat{r}_2\hat{r}_1 &= 4 + 2i + 6j + 8k.\end{aligned}$$

Hence and by Corollary 2 we have

$$\begin{aligned}Q_{n-1}Q_{n+1} - Q_n^2 &= -\frac{1}{5}(-1)^{n-1}[4 + 2i + 6j + 8k \\ &\quad + \frac{3 - \sqrt{5}}{2}(2 + (1 + \sqrt{5})i + (3 - \sqrt{5})j + (4 + \sqrt{5})k) \\ &\quad + \frac{3 + \sqrt{5}}{2}(2 + (1 - \sqrt{5})i + (3 + \sqrt{5})j + (4 - \sqrt{5})k)] \\ &= (-1)^n(2 + 4j + 3k) = (-1)^n(2Q_1 - 2i - 3k).\end{aligned}$$

We omit the proof of (ii). □

Proposition 3 *Let n, p, q be integers such that $n \geq 0$, $p^2 + 4q > 0$. Then*

$$H_{n+1}H_{n-1} - H_n^2 = -\alpha\beta(-q)^{n-1} \left(q(\hat{r}_1\hat{r}_2 + \hat{r}_2\hat{r}_1) + r_1^2\hat{r}_1\hat{r}_2 + r_2^2\hat{r}_2\hat{r}_1 \right).$$

For $p = 2$ and $q = 1$ we get the Cassini's identity for the split Pell quaternions SP_n and the split Pell-Lucas quaternions SPL_n ([7]).

Corollary 4 *Let $n \geq 1$ be an integer. Then*

$$\begin{aligned}SP_{n+1}SP_{n-1} - SP_n^2 &= (-1)^n(2 + 4i + 2j + 16k), \\ SPL_{n+1}SPL_{n-1} - SPL_n^2 &= (-1)^{n-1}(4 + 8i + 4j + 32k).\end{aligned}$$

Theorem 5 (*d'Ocagne's identity*) *Let m, n, p, q be integers such that $n \geq 0$, $p^2 + 4q > 0$. Then*

$$H_nH_{m+1} - H_{n+1}H_m = \frac{(-q)^m(b - ar_2)(b - ar_1)}{r_1 - r_2} (r_1^{n-m}\hat{r}_1\hat{r}_2 - r_2^{n-m}\hat{r}_2\hat{r}_1),$$

where $\hat{r}_1\hat{r}_2, \hat{r}_2\hat{r}_1$ are given by (16), (17), resp.

Proof. By (15) we get

$$\begin{aligned}
 H_n H_{m+1} - H_{n+1} H_m &= (\alpha \hat{r}_1 r_1^n - \beta \hat{r}_2 r_2^n)(\alpha \hat{r}_1 r_1^{m+1} - \beta \hat{r}_2 r_2^{m+1}) \\
 &\quad - (\alpha \hat{r}_1 r_1^{n+1} - \beta \hat{r}_2 r_2^{n+1})(\alpha \hat{r}_1 r_1^m - \beta \hat{r}_2 r_2^m) \\
 &= \alpha \beta (r_1 - r_2) (r_1^n r_2^m \hat{r}_1 \hat{r}_2 - r_1^m r_2^n \hat{r}_2 \hat{r}_1) \\
 &= \alpha \beta (r_1 - r_2) (r_1 r_2)^m (r_1^{n-m} \hat{r}_1 \hat{r}_2 - r_2^{n-m} \hat{r}_2 \hat{r}_1) \\
 &= \frac{(b - ar_2)(b - ar_1)(-q)^m}{r_1 - r_2} (r_1^{n-m} \hat{r}_1 \hat{r}_2 - r_2^{n-m} \hat{r}_2 \hat{r}_1).
 \end{aligned}$$

□

In the next theorem we give a summation formula for the split Horadam quaternions.

Theorem 6 Let n, p, q be integers such that $n \geq 0$, $p^2 + 4q > 0$. Then

$$\sum_{l=0}^n H_l = \frac{H_{n+1} + qH_n + (ap - a - b)(1 + i + j + k)}{p + q - 1} - ia - j(a + b) - k(a + b + pb + qa).$$

Proof. By formula (12) we get

$$\begin{aligned}
 \sum_{l=0}^n H_l &= \sum_{l=0}^n H_l + i \sum_{l=0}^n H_{l+1} + j \sum_{l=0}^n H_{l+2} + k \sum_{l=0}^n H_{l+3} \\
 &= \frac{1}{p + q - 1} [W_{n+1} + qW_n + a(p - 1) - b + i(W_{n+2} + qW_{n+1} + a(p - 1) - b) \\
 &\quad + j(W_{n+3} + qW_{n+2} + a(p - 1) - b) + k(W_{n+4} + qW_{n+3} + a(p - 1) - b)] \\
 &\quad - iW_0 - j(W_0 + W_1) - k(W_0 + W_1 + W_2).
 \end{aligned}$$

Hence we obtain

$$\begin{aligned}
 \sum_{l=0}^n H_l &= \frac{1}{p + q - 1} [W_{n+1} + iW_{n+2} + jW_{n+3} + kW_{n+4} \\
 &\quad + q(W_n + iW_{n+1} + jW_{n+2} + kW_{n+3} + (ap - a - b)(1 + i + j + k)) \\
 &\quad - ia - j(a + b) - k(a + b + pb + qa)] \\
 &= \frac{H_{n+1} + qH_n + (ap - a - b)(1 + i + j + k)}{p + q - 1} \\
 &\quad - ia - j(a + b) - k(a + b + pb + qa).
 \end{aligned}$$

□

For $p = q = 1$ and $a = 0, b = 1$ we get the result for the split Fibonacci quaternions Q_n ([1]).

Corollary 5 $\sum_{l=1}^n Q_l = Q_{n+2} - Q_2$.

Now we will give the generating function of the split Horadam quaternions.

Theorem 7 *The generating function of the split Horadam quaternions is*

$$f(x) = \frac{H_0 + (H_1 - pH_0)x}{1 - px - qx^2}.$$

Proof. Let $f(x) = H_0 + H_1x + H_2x^2 + \dots + H_nx^n + \dots$. Then

$$\begin{aligned} pxf(x) &= pH_0x + pH_1x^2 + pH_2x^3 + \dots + pH_{n-1}x^n + \dots \\ qx^2f(x) &= qH_0x^2 + qH_1x^3 + qH_2x^4 + \dots + qH_{n-2}x^n + \dots \end{aligned}$$

Hence, by Proposition 2, we get

$$\begin{aligned} f(x) - pxf(x) - qx^2f(x) &= H_0 + (H_1 - pH_0)x + (H_2 - pH_1 - qH_0)x^2 + \dots \\ &= H_0 + (H_1 - pH_0)x. \end{aligned}$$

Thus

$$f(x) = \frac{H_0 + (H_1 - pH_0)x}{1 - px - qx^2}.$$

Moreover, by (14) we obtain

$$\begin{aligned} H_0 &= a + ib + j(pb + qa) + k(p^2b + pqa + qb), \\ H_1 - pH_0 &= b - pa + iqa + jqb + k(pqb + q^2a). \end{aligned}$$

□

References

- [1] M. Akyiğit, H. H. Kösal, M. Tosun, Split Fibonacci Quaternions, *Adv. Appl. Clifford Algebr.*, **23** (2013), 535–545.
- [2] J. Cockle, On systems of algebra involving more than one imaginary, *Philosophical Magazine III*, **35(238)** (1849), 434–435.
- [3] A. F. Horadam, Basic properties of a certain generalized sequence of numbers, *Fibonacci Quart.*, **3.3** (1965), 161–176.

- [4] A. F. Horadam, Complex Fibonacci Numbers and Fibonacci Quaternions, *Amer. Math. Monthly*, **70** (1963), 289–291.
- [5] A. A. Pogoruy, R. M. R. Rodrigues-Dagnino, Some algebraic and Analytical Properties of Coquaternion Algebra, *Adv. Appl. Clifford Algebr.*, **20** (2010), 79–84.
- [6] E. Polatli, C. Kizilates, S. Kesim, On Split k -Fibonacci and k -Lucas Quaternions, *Adv. Appl. Clifford Algebr.*, **26** (2016), 353–362.
- [7] Ü. Tokeşer, Z. Ünal, G. Bilgici, Split Pell and Pell-Lucas Quaternions, *Adv. Appl. Clifford Algebr.*, **27** (2017), 1881–1893.

Received: August 21, 2019

On topological properties of the set of maldistributed sequences

József Bukor

Department of Informatics,
J. Selye University,
Komárno, Slovakia
email: bukorj@ujvs.sk

János T. Tóth

Department of Mathematics,
J. Selye University,
Komárno, Slovakia
email: tothj@ujvs.sk

Abstract. The real sequence (x_n) is maldistributed if for any non-empty interval I , the set $\{n \in \mathbb{N} : x_n \in I\}$ has upper asymptotic density 1. The main result of this note is that the set of all maldistributed real sequences is a residual set in the set of all real sequences (i.e., the maldistribution is a typical property in the sense of Baire categories). We also generalize this result.

1 Introduction

Following the concept of statistical convergence for real sequences, J. A. Fridy [2] introduced the concept of statistical cluster points of a sequence (x_n) . A number α is called a statistical cluster point of the sequence (x_n) provided that for every $\varepsilon > 0$ the set $\{n \in \mathbb{N} : |x_n - \alpha| < \varepsilon\}$ has a positive upper asymptotic density.

G. Myerson [7] calls a sequence (x_n) maldistributed if for any non-empty interval I the set $\{n \in \mathbb{N} : x_n \in I\}$ has upper asymptotic density 1. In [12] the maldistribution property is characterized by one-jump distribution functions. Examples of maldistributed sequences are given in [12] and [3]. Using the idea from [4] (Example VII) for the generalization of the concept of statistical

2010 Mathematics Subject Classification: 54E52

Key words and phrases: maldistributed sequence, weighted density, Baire category

convergence, we can extend the maldistribution property of sequences with the help of weighted densities.

The concept of weighted density as a generalization of asymptotic density was introduced in [1] and [10]. Let $f : \mathbb{N} \rightarrow (0, \infty)$ be a weight function with the properties

$$\sum_{n=1}^{\infty} f(n) = \infty, \quad \lim_{n \rightarrow \infty} \frac{f(n)}{\sum_{a \leq n} f(a)} = 0. \quad (1)$$

For $A \subset \mathbb{N}$ define by

$$\underline{d}_f(A) = \liminf_{n \rightarrow \infty} \frac{\sum_{a \leq n, a \in A} f(a)}{\sum_{a \leq n} f(a)} \quad \text{and} \quad \bar{d}_f(A) = \limsup_{n \rightarrow \infty} \frac{\sum_{a \leq n, a \in A} f(a)}{\sum_{a \leq n} f(a)}$$

the lower and upper f -densities of A , respectively. Note that the asymptotic densities correspond to $f(n) = 1$ and the logarithmic densities to $f(n) = \frac{1}{n}$. It is well-known that each set which has asymptotic density also has the logarithmic one but a set may have a logarithmic density without having an asymptotic one.

The main tool to compare weighted densities is the classical result of C. T. Rajagopal (cf. [9], Theorem 3) which, in terms of weighted densities, says the following.

Let $f, g : \mathbb{N} \rightarrow (0, \infty)$ be weight functions with properties (1). If $\frac{f(n)}{g(n)}$ is decreasing, then for any $A \subset \mathbb{N}$ we have

$$\underline{d}_g(A) \leq \underline{d}_f(A) \leq \bar{d}_f(A) \leq \bar{d}_g(A). \quad (2)$$

Now we give a generalization of maldistributed sequences.

Definition 1 Let $f : \mathbb{N} \rightarrow (0, \infty)$ be a weight function with properties (1). The sequence (x_n) is said to be f -maldistributed, if for any non-empty interval I the set $\{n \in \mathbb{N} : x_n \in I\}$ has upper f -density 1.

Comparing to asymptotic density, logarithmic density is less sensitive to certain perturbations. For example, if a sequence is maldistributed, then it is not necessary f -maldistributed for $f(n) = \frac{1}{n}$ (which defines the logarithmic density).

Let us denote by \mathcal{M}_f the set of all f -maldistributed sequences. The purpose of this note is to show that for any weight function f satisfying (1) the set \mathcal{M}_f is residual in the Fréchet metric space of all real sequences.

Let \mathbf{s} be the Fréchet metric space of all sequences of real numbers with the metric

$$\rho(\mathbf{x}, \mathbf{y}) = \sum_{k=1}^{\infty} \frac{1}{2^k} \frac{|x_k - y_k|}{1 + |x_k - y_k|},$$

where $\mathbf{x} = (x_k)$, $\mathbf{y} = (y_k)$. It is known that (\mathbf{s}, ρ) is a complete metric space.

In [5] it was proved that the set of all uniformly distributed sequences is a dense subset of the first Baire category in \mathbf{s} . The same is true for the set of all statistically convergent sequences of real numbers (cf. [11]).

2 Main results

The main result of this paper is as follows.

Theorem 1 *Let $f : \mathbb{N} \rightarrow (0, \infty)$ be a weight function with properties (1). Then the set of all f -maldistributed sequences \mathcal{M}_f is residual in the the Fréchet metric space of all sequences of real numbers \mathbf{s} .*

For the proof of the theorem we shall use the following lemma.

Lemma 1 *For the interval $I = [a, b]$ denote by $\mathcal{A}(I, \alpha)$ the set of all $\mathbf{x} = (x_k) \in \mathbf{s}$ for which*

$$\overline{d}_f(\{n \in \mathbb{N} : x_n \in I\}) \leq \alpha,$$

where $\alpha \in (0, 1)$. Then $\mathcal{A}(I, \alpha)$ is a set of the first Baire category in \mathbf{s} .

Proof of Lemma 1. We define a continuous function $h : \mathbb{R} \rightarrow [0, 1]$ by

$$h(x) = \begin{cases} \frac{2x-2a}{b-a} & \text{for } x \in [a, \frac{a+b}{2}] \\ \frac{2b-2x}{b-a} & \text{for } x \in [\frac{a+b}{2}, b] \\ 0 & \text{for } x \in \mathbb{R} \setminus [a, b] \end{cases}$$

We choose an arbitrary real number $\beta \in (\alpha, 1)$. Using the function h we define for $\mathbf{x} = (x_k) \in \mathbf{s}$ and fixed n the function $g_n : \mathbf{s} \rightarrow [0, 1]$ in the following way:

$$g_n(\mathbf{x}) = \max \left\{ \beta, \frac{\sum_{k=1}^n h(x_k) \cdot f(k)}{\sum_{k=1}^n f(k)} \right\}.$$

Denote $\mathcal{A}^*(I, \alpha)$ the set of all $\mathbf{x} = (x_k) \in \mathbf{s}$ for which there exists the limit $\lim_{n \rightarrow \infty} g_n(\mathbf{x})$.

One can easily check that for each $\mathbf{x} = (x_k) \in \mathbf{s}$ and natural number n we have

$$\frac{\sum_{k=1}^n h(x_k).f(k)}{\sum_{k=1}^n f(k)} \leq \frac{\sum_{k \leq n, x_k \in I} f(k)}{\sum_{k \leq n} f(k)}. \quad (3)$$

For any $\mathbf{x} \in \mathcal{A}(I, \alpha)$, the right hand side of (3) does not exceed α if n is large enough. Therefore $\lim_{n \rightarrow \infty} g_n(\mathbf{x}) = \beta$, and then $\mathcal{A}(I, \alpha) \subset \mathcal{A}^*(I, \alpha)$.

Put $g(\mathbf{x}) = \lim_{n \rightarrow \infty} g_n(\mathbf{x})$ for $\mathbf{x} \in \mathcal{A}^*(I, \alpha)$. We shall prove that

- (a) the function g_n ($n = 1, 2, \dots$) is a continuous function on \mathbf{s} ,
- (b) g is discontinuous at each point of $\mathcal{A}^*(I, \alpha)$.

(a) Let $\mathbf{x}^0 = (x_k^0)_{k=1}^\infty$, $\mathbf{x}^{(j)} = (x_k^{(j)})_{k=1}^\infty \in \mathbf{s}$ ($j = 1, 2, \dots$) and $\mathbf{x}^{(j)} \rightarrow \mathbf{x}^0$ (for $j \rightarrow \infty$).

Then from the convergence in the space \mathbf{s} for each fixed k we have $\lim_{j \rightarrow \infty} x_k^{(j)} = x_k^0$. The continuity of function h implies $\lim_{j \rightarrow \infty} g_n(\mathbf{x}^{(j)}) = g_n(\mathbf{x}^0)$. Thus g_n ($n = 1, 2, \dots$) is continuous on \mathbf{s} .

(b) Let $\mathbf{y} = (y_k) \in \mathcal{A}^*(I, \alpha)$. We have the following two possibilities.

- (1) $g(\mathbf{y}) < 1$,
- (2) $g(\mathbf{y}) = 1$.

In case (1) we choose a positive ε such that $\varepsilon < 1 - g(\mathbf{y})$. It is suffice to prove that in each ball $K(\mathbf{y}, \delta) = \{\mathbf{x} \in \mathcal{A}^*(I, \alpha), \rho(\mathbf{x}, \mathbf{y}) < \delta\}$ ($\delta > 0$) of the subspace $\mathcal{A}^*(I, \alpha)$ of \mathbf{s} there exists an element $\mathbf{x} = (x_k) \in \mathbf{s}$ with $g(\mathbf{x}) - g(\mathbf{y}) > \varepsilon$.

Let $\delta > 0$. Choose a positive integer m such that $\sum_{k=m+1}^\infty 2^{-k} < \delta$, and define the sequence $\mathbf{x} = (x_k)$ in the following way:

$$x_k = \begin{cases} y_k, & \text{if } k \leq m, \\ \frac{a+b}{2}, & \text{if } k > m. \end{cases}$$

Hence $\rho(\mathbf{x}, \mathbf{y}) < \delta$, further $h(x_k) = 1$ for $k > m$. Then

$$\frac{\sum_{k=1}^n h(x_k).f(k)}{\sum_{k=1}^n f(k)} \geq \frac{\sum_{k=m+1}^n f(k)}{\sum_{k=1}^n f(k)} = 1 - \frac{\sum_{k=1}^m f(k)}{\sum_{k=1}^n f(k)} \rightarrow 1 \text{ for } n \rightarrow \infty,$$

and therefore $g(\mathbf{x}) = \lim_{n \rightarrow \infty} g_n(\mathbf{x}) = 1$. Then immediately follows

$$g(\mathbf{x}) - g(\mathbf{y}) = 1 - g(\mathbf{y}) > \varepsilon.$$

In case (2) we have $g(\mathbf{y}) = 1$. Let δ, m, \mathbf{x} have the previous meaning. Put

$$x_k = \begin{cases} y_k, & \text{if } k \leq m, \\ a, & \text{if } k > m. \end{cases}$$

Then, clearly $\rho(\mathbf{x}, \mathbf{y}) < \delta$, and $h(x_k) = 0$ for $k > m$. Then

$$\frac{\sum_{k=1}^n h(x_k) \cdot f(k)}{\sum_{k=1}^n f(k)} \leq \frac{\sum_{k=1}^m f(k)}{\sum_{k=1}^n f(k)} \rightarrow 0 \quad \text{for } n \rightarrow \infty.$$

So, we have $g(\mathbf{x}) = \lim_{n \rightarrow \infty} g_n(\mathbf{x}) = \beta$, and therefore $g(\mathbf{y}) - g(\mathbf{x}) = 1 - \beta > 0$.

Hence the discontinuity of g at $\mathbf{y} \in \mathcal{A}^*(I, \alpha)$ has been proved.

The function g is a limit function (on $\mathcal{A}^*(I, \alpha)$) of the sequence of continuous functions $(g_n)_{n=1}^\infty$ on $\mathcal{A}^*(I, \alpha)$. Then the function g is a function in the first Baire class on $\mathcal{A}^*(I, \alpha)$. According to the well-known fact that the set of discontinuity points of an arbitrary function of the first Baire class is a set of the first Baire category (cf. [8], p. 32), we see that the set $\mathcal{A}^*(I, \alpha)$ is of the first Baire category in $\mathcal{A}^*(I, \alpha)$. Thus $\mathcal{A}^*(I, \alpha)$ is in \mathbf{s} , too. Since $\mathcal{A}(I, \alpha) \subset \mathcal{A}^*(I, \alpha)$, the assertion follows. \square

Proof of Theorem 1. Denote by \mathbb{Q} the set of all rational numbers. Denote by \mathcal{H} the set of all $\mathbf{x} = (x_k) \in \mathbf{s}$ for which there exists an interval I with

$$\bar{d}_f(\{n \in \mathbb{N} : x_n \in I\}) \leq \alpha$$

for some $\alpha \in (0, 1)$. Combining Lemma 1 and the fact that for each interval I there exist rational numbers a, b such that $I \subset [a, b]$, we have

$$\mathcal{H} \subset \bigcup_{a, b \in \mathbb{Q}, a < b} \bigcup_{i \in \mathbb{N}, i \geq 2} A\left([a, b], 1 - \frac{1}{i}\right)$$

from which follows at once that \mathcal{H} is a meager set. But $\mathcal{M}_f = \mathbf{s} \setminus \mathcal{H}$ and therefore the assertion of theorem follows. Hence the property of f -maldistribution is a typical property of real sequences from the topological point of view. \square

We now introduce the concept of f -maldistributed integer sequences.

Definition 2 Let $f : \mathbb{N} \rightarrow (0, \infty)$ be a weight function with properties (1). The sequence (x_n) of positive integers is said to be f -maldistributed, if for any positive integers $m \geq 2$ and $j \in \{0, 1, \dots, m-1\}$ the set $\{n \in \mathbb{N} : x_n \equiv j \pmod{m}\}$ has upper f -density 1.

Let \mathbf{S} be the Baire's space of all sequences of positive integers with the metric ρ' defined in the following way.

Let $\mathbf{x} = (x_k) \in \mathbf{S}$, and $\mathbf{y} = (y_k) \in \mathbf{S}$. If $\mathbf{x} = \mathbf{y}$, then $\rho'(\mathbf{x}, \mathbf{y}) = 0$, otherwise

$$\rho'(\mathbf{x}, \mathbf{y}) = \frac{1}{\min\{n : x_n \neq y_n\}}.$$

The space (\mathbf{S}, ρ') is a complete metric space. In [6] the topological properties of the set of all uniformly distributed sequences of positive integers in Baire's space were investigated.

The following auxiliary result is similar to Lemma 1.

Lemma 2 For the positive integers $m \geq 2$ and $j \in \{0, 1, \dots, m-1\}$ denote by $\mathcal{A}(j, m, \alpha)$ the set of all $\mathbf{x} = (x_k) \in \mathbf{S}$ for which

$$\overline{d}_f(\{n \in \mathbb{N} : x_n \equiv j \pmod{m}\}) \leq \alpha,$$

where $\alpha \in (0, 1)$. Then $\mathcal{A}(j, m, \alpha)$ is a set of the first Baire category in \mathbf{S} .

The proof is analogous to the proof of Lemma 1. The crucial role is played by the function $g_n : \mathbf{S} \rightarrow [0, 1]$ given by

$$g_n(\mathbf{x}) = \max \left\{ \sqrt{\alpha}, \frac{\sum_{\substack{k \leq n \\ x_k \equiv j \pmod{m}}} f(k)}{\sum_{k=1}^n f(k)} \right\}.$$

The following theorem says that the set of all f -maldistributed integer sequences form a residual set in Baire's space.

Theorem 2 Let $f : \mathbb{N} \rightarrow (0, \infty)$ be a weight function with properties (1). Denote by \mathcal{G} the set of all $\mathbf{x} = (x_k) \in \mathbf{S}$ for which there exist $m \geq 2$ and $j \in \{0, 1, \dots, m-1\}$ such that

$$\overline{d}_f(\{n \in \mathbb{N} : x_n \equiv j \pmod{m}\}) \leq \alpha$$

for some $\alpha \in (0, 1)$. Then \mathcal{G} is a set of the first Baire category in \mathbf{S} .

Proof. Combining Lemma 2 with the fact that

$$\mathcal{G} \subset \bigcup_{m=2}^{+\infty} \bigcup_{j=0}^{m-1} \bigcup_{i=2}^{+\infty} A\left(j, m, 1 - \frac{1}{i}\right)$$

it immediately follows that \mathcal{G} is a meager set in \mathbf{S} . □

References

- [1] R. Alexander, Density and multiplicative structure of sets of integers, *Acta Arith.*, **12** (1976), 321–332.
- [2] J. A. Fridy, Statistical limit points, *Proc. Amer. Math. Soc.*, **118** (1993), 1187–1192.
- [3] P. Kostyrko, M. Mačaj, T. Šalát, O. Strauch, On statistical limit points, *Proc. Amer. Math. Soc.*, **129** (2000), 2647–2654.
- [4] P. Kostyrko, M. Mačaj, T. Šalát, O. Strauch, I-convergence and extremal I-limit points, *Math. Slovaca*, **55** (2005), 443–464.
- [5] V. László, T. Šalát, The structure of some sequence spaces, and uniform distribution (mod 1), *Periodica Math. Hung.*, **10** (1979), 89–98.
- [6] V. László, T. Šalát, Uniformly distributed sequences of positive integers in Baire’s space, *Math. Slovaca*, **41** (1991), 277–281.
- [7] G. Myerson, A sampler of recent developments in the distribution of sequences, *Lecture Notes in pure and applied Mathematics*, **147** (1993), 163–190.
- [8] J. C. Oxtoby. *Measure and Category*. Graduate texts in Mathematics. Springer, 1980.
- [9] C. T. Rajagopal, Some limit theorems, *Amer. J. Math.*, **70** (1948), 157–166.
- [10] H. Rohrbach, B. Volkmann, Verallgemeinerte asymptotische Dichten, *J. Reine Angew. Math.*, **194** (1955), 195–209.

- [11] T. Šalát, On statistically convergent sequences of real numbers, *Math. Slovaca*, **30** (1980), 139–150.
- [12] O. Strauch, Uniformly maldistributed sequences in a strict sense, *Monatsh. Math.*, **120** (1995), 153–164.

Received: February 20, 2019

On the connection between tridiagonal matrices, Chebyshev polynomials, and Fibonacci numbers

Carlos M. da Fonseca

Kuwait College of Science and Technology,
Doha District, Block 4, P.O. Box 27235,
Safat 13133, Kuwait
email: c.dafonseca@kcst.edu.kw

Abstract. In this note, we recall several connections between the determinant of some tridiagonal matrices and the orthogonal polynomials allowing the relation between Chebyshev polynomials of second kind and Fibonacci numbers. With basic transformations, we are able to recover some recent results on this matter, bringing them into one place.

1 Orthogonal polynomials and tridiagonal matrices

From the elementary orthogonal polynomials theory, we know that any monic sequence orthogonal polynomial sequence $\{P_n(x)\}$ is defined by the recurrence formula

$$P_n(x) = (x - c_n) P_{n-1}(x) - \lambda_n P_{n-2}(x), \quad \text{for } n = 1, 2, \dots \quad (1)$$

with initial conditions $P_{-1}(x) = 0$ and $P_0(x) = 1$, and for complex numbers c_n 's and λ_n 's, if and only if $\lambda_n \neq 0$ [6, Theorem 4.4]. Moreover, (1) is equivalent

2010 Mathematics Subject Classification: 11B37, 11B39, 15A15, 15B05, 15B36, 39A10, 42C05

Key words and phrases: tridiagonal matrices, Chebyshev polynomials of second kind, determinant, Fibonacci numbers

to

$$P_n(x) = \begin{vmatrix} x - c_1 & 1 & & & \\ \lambda_2 & x - c_2 & 1 & & \\ & \lambda_3 & \ddots & \ddots & \\ & & \ddots & \ddots & 1 \\ & & & \lambda_n & x - c_n \end{vmatrix} \quad (2)$$

(cf. [6, Exercise 4.12, p.26]).

Setting $\lambda_n = 1$ and $c_n = 0$, for any positive integer n , in (1) or (2), we obtain $U_n(x) = P_n(2x)$, the Chebyshev polynomial of second kind of degree n (cf. [1, (22.7.5)] or [6, Exercise 4.9, p.25]). The Chebyshev polynomials of second kind satisfy the three-term recurrence relations

$$U_n(x) = 2xU_{n-1}(x) - U_{n-2}(x), \quad \text{for all } n = 1, 2, \dots, \quad (3)$$

or, equivalently, we may state

$$U_n(x) = \begin{vmatrix} 2x & 1 & & & \\ 1 & \ddots & \ddots & & \\ & \ddots & \ddots & 1 & \\ & & 1 & 2x \end{vmatrix}_{n \times n}.$$

Among the most important explicit representations for the $U_n(x)$ we have

$$U_n(x) = \frac{\sin(n+1)\theta}{\sin \theta}, \quad \text{with } x = \cos \theta \quad (0 \leq \theta < \pi), \quad (4)$$

as we can see, for example, in [1, (22.3.16)] or [6, Exercise 1.2, p.5]. From de Moivre's formula we have

$$U_n(x) = \frac{(x + \sqrt{x^2 - 1})^{n+1} - (x - \sqrt{x^2 - 1})^{n+1}}{2\sqrt{x^2 - 1}}. \quad (5)$$

Another formula with some relevance is

$$U_n(x) = \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} (-1)^k \binom{n-k}{k} (2x)^{n-2k} \quad (6)$$

which can be found for example in [1, (22.3.7)]. There are many other representations and relations for $U_n(x)$. As stated in [8, p.187], many of them are

paraphrases of trigonometric identities, derivations from (4). For particular values of $U_n(x)$ the reader is referred to [1, (22.4.5)].

Regarding the generating function for $U_n(x)$, we have

$$\frac{1}{1 - 2zx + z^2} = \sum_{n=0}^{\infty} U_n(x) z^n \quad (7)$$

(cf. [1, (22.9.10)]).

From (3), we can easily deduce that

$$(\sqrt{ab})^n U_n \left(\frac{c}{2\sqrt{ab}} \right) = \begin{vmatrix} c & a & & & \\ b & \ddots & \ddots & & \\ & \ddots & \ddots & a & \\ & & b & c & \end{vmatrix}_{n \times n}. \quad (8)$$

provided $ab \neq 0$. Perhaps one of the most interesting applications of this identity is

$$F_n = (-i)^{n-1} U_{n-1} \left(\frac{i}{2} \right),$$

where F_n is the n th Fibonacci number (cf. for example [2, 5, 12]), setting $a = c = 1$ and $b = -1$.

Many of these results were recently recast. In this short note, we aim to bring them into attention and put into one place. At the same time, we provide shorter proofs to many of them.

2 Determinants of tridiagonal matrices

In [14], the authors guessed that the determinant $D_n(c)$ in (8), for $a = b = 1$, satisfies

$$D_n(c) = \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} (-1)^k \binom{n-k}{k} c^{n-2k}.$$

This is repeatedly proved in [13] and in [16], while indeed it is an immediate consequence of (6). For that propose the authors in [13] proved (7) for $D_n(c)$.

Next, it is proved in [13] that

$$D_n(c) = \frac{\alpha^{n+1} - \beta^{n+1}}{\alpha - \beta},$$

where $\alpha = \frac{1}{\beta} = \frac{c+\sqrt{c^2-4}}{2}$, and the trivial cases are ignored. In [16], three alternative proofs are provided to this fact. Nonetheless, this equality is immediate from (5), if one notices that $\beta = \frac{c-\sqrt{c^2-4}}{2}$. A particular case of (8) is also claimed to be proved by induction when $c = a + b$, using (5).

Another topic discussed in [13] is the inverse of

$$A = \begin{pmatrix} c & a & & & \\ b & \ddots & \ddots & & \\ & \ddots & \ddots & a & \\ & & b & c & \end{pmatrix}_{n \times n},$$

for $a = b = 1$. We remark that this matrix is occasionally called in [13] “diagonal matrix”, but this is obviously not appropriate. It is well-known that if A is nonsingular, i.e., $U_n(d) \neq 0$, with $d = c/(2\sqrt{ab})$, then its inverse is given by

$$(A^{-1})_{ij} = \begin{cases} (-1)^{i+j} \frac{a^{j-i}}{(\sqrt{ab})^{j-i+1}} \frac{U_{i-1}(d) U_{n-j}(d)}{U_n(d)} & \text{if } i \leq j \\ (-1)^{i+j} \frac{b^{i-j}}{(\sqrt{ab})^{i-j+1}} \frac{U_{j-1}(d) U_{n-i}(d)}{U_n(d)} & \text{if } i > j. \end{cases}$$

This can be found in [10, 19, 20, 21] or easily deduced from [11, p.28]. Yet, in [13] we can see this inverse for the particular case of $a = b = 1$. Moreover, in [16] the authors provide a detailed proof for it, while this particular case has been studied for the last 75 years (cf. [9, 17, 18]).

We also want to remark that the eigenpairs of A in [13] are wrong. Indeed, it is a standard result that, for example, the eigenvalues of A are

$$\lambda_k = c + 2\sqrt{ab} \cos\left(\frac{\ell\pi}{n+1}\right), \quad \text{for } \ell = 1, \dots, n.$$

On contrary to what is suggested in [13], the real entries of the tridiagonal matrix A are absolutely irrelevant for this formula.

Finally, [13] provides an intricate proof for the equality

$$\begin{vmatrix} -c & 1 & & & \\ 2 & -2c & 1 & & \\ & 6 & \ddots & \ddots & \\ & & \ddots & \ddots & 1 \\ & & & n(n-1) & -nc \end{vmatrix} = (-1)^n n! \begin{vmatrix} c & 1 & & & \\ 1 & c & 1 & & \\ & 1 & \ddots & \ddots & \\ & & \ddots & \ddots & 1 \\ & & & 1 & c \end{vmatrix}. \quad (9)$$

In [16], among others, the authors claim that (9) is “neither trivial nor obvious”. Next, using elementary matrix theory, we show that (9) is both trivial and obvious (see also [3]). In fact,

$$\begin{aligned}
 \begin{vmatrix} -c & 1 & & & \\ 2 & -2c & 1 & & \\ & 6 & \ddots & \ddots & \\ & & \ddots & \ddots & 1 \\ & & & n(n-1) & -nc \end{vmatrix} &= (-1)^n \begin{vmatrix} c & 1 & & & \\ 2 & 2c & 2 & & \\ & 3 & \ddots & \ddots & \\ & & \ddots & \ddots & n-1 \\ & & & n & nc \end{vmatrix} \\
 &= (-1)^n n! \begin{vmatrix} c & 1 & & & \\ 1 & c & 1 & & \\ & 1 & \ddots & \ddots & \\ & & \ddots & \ddots & 1 \\ & & & 1 & c \end{vmatrix}.
 \end{aligned}$$

We observe that the same identity can be found and proved using an involved approach in [15].

3 Central Delannoy numbers

Our final comment goes to the last section of the paper [13].

The Legendre polynomials $P_n(x)$ are a particular family of the ultraspherical polynomials [4, p.899]. Their generating function is

$$\frac{1}{\sqrt{1-2tx+t^2}}$$

and they can be defined, for example, by the contour integral

$$P_n(x) = \frac{1}{2\pi i} \oint \frac{1}{\sqrt{1-2xt+t^2}} \frac{1}{t^{n+1}} dt$$

where the contour encloses the origin and is traversed in a counterclockwise direction [4, p.416]. The central Delannoy numbers $D(n)$ are defined as ([7, p.81])

$$D(n) = P_n(3).$$

The authors recover a generalization for the central Delannoy numbers, $D_{a,b}(n)$, with generating function

$$\frac{1}{\sqrt{(x+a)(x+b)}} = \sum_{k=0}^{\infty} D_{a,b}(k) x^k.$$

However, next they claim that squaring both sides of the previous identity one gets

$$\frac{1}{(x+a)(x+b)} = \sum_{k=0}^{\infty} D_{a,b}(k) x^k,$$

which is obviously not true.

References

- [1] M. Abramowitz, I. A. Stegun, *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, ninth edition, Dover Publications, Inc., New York, 1970.
- [2] M. Anđelić, Z. Du, C. M. da Fonseca, E. Kılıç, A matrix approach to some second-order difference equations with sign-alternating coefficients, *J. Difference Equ. Appl.*, **26** (2020), 149–162.
- [3] M. Anđelić, C. M. da Fonseca, A determinantal formula for generalized Fibonacci numbers, *Matematiche (Catania)*, **74** (2019), 363–367.
- [4] G. B. Arfken, H. J. Weber, F. E. Harris, *Mathematical Methods for Physicists: A Comprehensive Guide*, Academic Press, Orlando, 1985.
- [5] R. G. Buschman, Fibonacci numbers, Chebyshev polynomials generalizations and difference equations, *Fibonacci Quart.* 1 (1963), 1–8, 19.
- [6] T. S. Chihara, *An Introduction to Orthogonal Polynomials*, Gordon and Breach, New York, 1978.
- [7] L. Comtet, *Advanced Combinatorics: The Art of Finite and Infinite Expansions*, Reidel, Dordrecht, Netherlands, 1974.
- [8] Erdélyi, et al., *Higher Transcendental Functions*, vol. 2, McGraw Hill, New York, 1953.
- [9] C. F. Fischer, R. A. Usmani, Properties of some tridiagonal matrices and their application to boundary value problems, *SIAM J. Numer. Anal.*, **6** (1969), 127–141.
- [10] C. M. da Fonseca, J. Petronilho, Explicit inverses of some tridiagonal matrices, *Linear Algebra Appl.*, **325** (2001), 7–21.

- [11] G. Heinig, K. Rost, *Algebraic Methods for Toeplitz-Like Matrices and Operators*, Birkhäuser OT 13, 1984.
- [12] A. F. Horadam, Basic properties of a certain generalized sequence of numbers, *Fibonacci Quart.*, **3** (1965), 161–176.
- [13] F. Qi, V. Čerňanová, Y. S. Semenov, Some tridiagonal determinants related to central Delannoy numbers, the Chebyshev polynomials, and the Fibonacci polynomials, *Politehn. Univ. Bucharest Sci. Bull. Ser. A Appl. Math. Phys.*, **81** (2019), 123–136.
- [14] F. Qi, V. Čerňanová, X.-T. Shi, B.-N. Guo, Some properties of central Delannoy numbers, *J. Comput. Appl. Math.*, **328** (2018), 101–115.
- [15] F. Qi, B.-N. Guo, Expressing the generalized Fibonacci polynomials in terms of a tridiagonal determinant, *Matematiche* (Catania), **72** (2017), 167–175.
- [16] F. Qi, A.-Q. Liu, Alternative proofs of some formulas for two tridiagonal determinants, *Acta Univ. Sapientiae, Mathematica*, **10** (2018), 287–297.
- [17] G. Meurant, A review on the inverse of symmetric tridiagonal and block tridiagonal matrices, *SIAM J. Matrix Anal. Appl.*, **13** (1992), 707–728.
- [18] D. Moskovitz, The numerical solution of Laplace’s and Poisson’s equations, *Quart. Appl. Math.*, **2** (1944), 14S–63.
- [19] P. Schlegel, *The explicit inverse of a tridiagonal matrix*, *Math. Comp.* **24** (1970), 665.
- [20] R. Usmani, Inversion of a tridiagonal Jabobi matrix, *Linear Algebra Appl.*, **212/213** (1994), 413–414.
- [21] T. Yamamoto, Y. Ikebe, Inversion of band matrices, *Linear Algebra Appl.*, **24** (1979), 105–111.

Received: November 20, 2019



A note on nil-clean rings

Peter V. Danchev

Institute of Mathematics and Informatics,
Bulgarian Academy of Sciences,
“Acad. G. Bonchev” str., bl. 8, 1113 Sofia, Bulgaria
email: danchev@math.bas.bg;
pvdanchev@yahoo.com

Abstract. We study a special kind of nil-clean rings, namely those nil-clean rings whose nilpotent elements are difference of two “left-right symmetric” idempotents, and prove that in some various cases they are strongly π -regular. We also show that all nil-clean rings having cyclic unit 2-groups are themselves strongly nil-clean of characteristic 2 (and thus they are again strongly π -regular).

1 Introduction and background

Everywhere in the text of the present paper, all our rings R are assumed to be associative, containing the identity element 1 , which in general differs from the zero element 0 of R . Our terminology and notations are mainly standard being in agreement with [9]. Exactly, $U(R)$ denotes the set of all units in R , $\text{Id}(R)$ the set of all idempotents in R , $\text{Nil}(R)$ the set of all nilpotents in R and $J(R)$ the Jacobson radical of R .

A ring R is called *von Neumann regular* or just *regular* for short if, for any element $r \in R$, there is an element $a \in R$ such that $r = rar$. In the case when $a = 1$, we have that $r = r^2$ and these rings are known to be *boolean*. Generalizing regularity, a ring R is called *π -regular* if, for each $r \in R$, there are $i \in \mathbb{N}$ and $b \in R$ both depending on r such that $r^i = r^i b r^i$. Likewise, a ring R is called *strongly π -regular* if, for every $r \in R$, there exist $j \in \mathbb{N}$ and

2010 Mathematics Subject Classification: 16U99; 16E50; 16W10; 13B99

Key words and phrases: nil-clean rings, nilpotents, idempotents, strongly π -regular rings

$c \in R$ both depending on r with $r^j = r^{j+1}c$. It is well known that strongly π -regularity implies π -regularity, while the converse is wrong as some critical examples show (see, e.g., [9]).

On the other hand, referring to [7] for a more account, we shall say that a ring is *nil-clean* provided each its element is a sum of a nilpotent and an idempotent. If these two elements commute, the nil-clean ring is said to be *strongly nil-clean*. While nil-clean rings are not completely characterized up to an isomorphism yet, this was successfully done in [4] by proving that a ring R is strongly nil-clean if, and only if, the quotient ring $R/J(R)$ is boolean and $J(R)$ is nil.

That is why, classifying the structure of some special types of nil-clean rings will be of some interest and importance. Our workable purpose here is to examine those nil-clean rings whose nilpotents are differences of two (special) idempotents. Specifically, we shall prove that in Theorem 1 presented below that every nil-clean ring having only nilpotents which are difference of two special (so-called “left-right symmetric”) idempotents is strongly π -regular. This contrasts an example due to Šter in [11] who constructed a nil-clean ring of unbounded index of nilpotence which is not strongly π -regular. Note that by an appeal to [6, Corollary 3.12] nil-clean rings of bounded index of nilpotence are always strongly π -regular. We also consider the challenging question of when a nil-clean ring with finite (in particular, cyclic) unit group is strongly nil-clean. It is necessarily such a group to be consisting only of elements of order being a power of 2, and the ring will be of characteristic 2 too.

2 Main results

We separate our chief results into two subsections as follows:

2.1 Nil-clean rings with nilpotents as a sum of two special idempotents

We start our assertions with the next one.

Proposition 1 *If R is a nil-clean ring such that each nilpotent is a difference of two commuting idempotents, then R is a boolean ring.*

Proof. We first claim that such a ring R is of characteristic 2. Indeed, as $2 \in \text{Nil}(R)$ (see, e.g., [7]), one writes that $2 = e - f$ for some $e, f \in \text{Id}(R)$. Hence, it easily follows that $ef = fe$ even not assuming this a priori and,

therefore, $2^3 = (e - f)^3 = e - f = 2$. This means that $6 = 0$, i.e., $2 = 0$ because $3 \in \mathcal{U}(R)$ and the claim is sustained.

Moreover, we assert that R has to be abelian, that is, all its idempotents are central. In fact, given an arbitrary $a \in R$ and an arbitrary $e \in \text{Id}(R)$, one sees that $ea(1 - e) \in \text{Nil}(R)$ and thus $ea(1 - e) = e_1 + e_2$ for some $e_1, e_2 \in \text{Id}(R)$ with $e_1 e_2 = e_2 e_1$. Squaring this, it follows at once that $0 = e_1 + e_2$ since $2 = 0$ which yields $ea = eae$. Similarly, one derives that $ae = eae$ by looking at the element $(1 - e)ae$, which allows us to conclude that $ae = ea$, as asserted.

We next arrive at the fact that R is semi-primitive, which is equivalent to $J(R) = \{0\}$. To verify this, given any element $z \in J(R)$, one may write that $z = e - f$ for some $e, f \in \text{Id}(R)$ with $ef = fe$ since $J(R)$ is nil (see, for instance, [7]). Now, taking into account that $2 = 0$, we find that $z^2 = z$ whence $z(z - 1) = 0$ ensuring that $z = 0$ because $z - 1 \in \mathcal{U}(R)$. Thus R is semi-primitive, as claimed.

Furthermore, we may apply either [4] or [7] to get the desired boolean property of R . \square

It was established in [8, Proposition 1] that any nilpotent matrix over a field is a difference of two idempotent matrices (for another approach see [10] as well). This major statement allows us to extract the following assertion, independently proved also in [10] and partially in [3].

Lemma 1 *In regular rings all nilpotent elements are difference of two idempotents.*

Proof. Consulting with the main result from [1] which shows that, in an arbitrary ring, a nilpotent with all powers regular can be thought of as locally just a nilpotent matrix in Jordan or Weyr form. With this at hand, the aforementioned matrix result in [8] gives the desired presentation. \square

Imitating [3], two idempotents e, f are called *left-right symmetric* if the two equalities $ef = e$ and $fe = f$ hold. It is evident that both e and f are somewhat “left-active” in the sense that they are “preserved on the left multiplication”.

So, we have accumulated all the information necessary to establish the following.

Theorem 1 *Every nil-clean ring in which all nilpotents are difference of two left-right symmetric idempotents are strongly π -regular.*

Proof. We foremost assert that for such a ring R it must be that $\text{char}(R) = 2$. To see that, as $2 \in \text{Nil}(R)$ holds in view of [7], one writes that $2 = e_1 - e_2$ for two $e_1, e_2 \in \text{Id}(R)$. This surely means that e_1 and e_2 do commute, so that

$2^3 = (e_1 - e_2)^3 = e_1 - e_2 = 2$ whence $6 = 0$. Consequently, $2 = 0$ because $3 \in \mathcal{U}(\mathbf{R})$, as asserted.

For such a ring \mathbf{R} , given an arbitrary $q \in \text{Nil}(\mathbf{R})$, we write that $q = e - f = e + f$ for some two $e, f \in \text{Id}(\mathbf{R})$ with $ef = e$ and $fe = f$. We, therefore, obtain by squaring that $q^2 = 2q = 0$. Thus \mathbf{R} is of bounded index of nilpotence and [6, Corollary 3.12] is a guarantor for the validity of our assertion that \mathbf{R} is strongly π -regular. \square

The given proof allows us to consider whether a more general situation in which we have slightly amended relationships between e and f , that are, $efe = e$ and $fef = f$. Certainly, $ef = e$ forces $efe = e$ as well as $fe = f$ forces $fef = f$. Furthermore, writing $q = e + f$ and squaring this, we deduce that $q^2 - q = ef + fe$. Again squaring the last equality, we derive that $q^4 + q^2 = (q^2 - q)^2 = efef + efef + fef + fefe = ef + e + f + fe = q^2$. Finally, $q^4 = 0$ and hence \mathbf{R} is with bounded index of nilpotence, too.

We can now mention some constructions of nil-clean rings having only nilpotent elements which are difference of two idempotents.

Remark 1 *By what we have just previously shown, a crucial example of such a sort of nil-clean rings is any nil-clean ring which is simultaneously regular – in fact, such is, for instance, the ring $\mathbb{M}_n(\mathbb{Z}_2)$ for all $n \geq 1$ by an appeal to [2] and to the well-known fact from [9] that it is a regular ring because so is \mathbb{Z}_2 . Indeed, this is not always possible as it was recently exhibited in [11] an ingenious example of a nil-clean ring of characteristic 2 which is not strongly π -regular as well as of a nil-clean ring of characteristic 4 which is not π -regular.*

An other interesting example of a nil-clean ring whose nilpotent elements are differences of two idempotents and which ring is not regular (due to the fact that it has a non-zero Jacobson radical) is the upper triangular matrix ring $\mathbb{T}_2(\mathbb{Z}_2)$, which fact we leave to the interested reader for a direct inspection. This ring is, however, strongly π -regular.

Moreover, the indecomposable nil-clean ring \mathbb{Z}_4 does not have the indicated above specific property of its nilpotents since $2 \neq 0$ in it.

We end our work in this subsection with the following challenging problem.

Problem 1 *Characterize nil-clean rings whose nilpotent elements are differences of two arbitrary idempotents.*

2.2 Nil-clean rings with cyclic unit group

In [5, p.81] it was asked of whether or not a clean ring with cyclic units is strongly clean. We shall resolve this question in the case of nil-clean rings (note that nil-clean rings are always clean and a *clean* ring is the one whose elements are sums of a unit and an idempotent; if these two elements commute, the clean ring is called *strongly clean*). It was established in [4, Corollary 4.10] that a nil-clean is strongly nil-clean if, and only if, its unit group is a 2-group.

We are now arriving at the following statement.

Theorem 2 *Suppose R is a nil-clean ring with cyclic $U(R)$. Then R is strongly nil-clean of characteristic 2 if, and only if, $U(R)$ is a 2-group.*

Proof. If we assume for a moment that $U(R) = \{1\}$, then $\text{Nil}(R) = \{0\}$ as $1 + \text{Nil}(R) \subseteq U(R)$, so that R must be boolean whence strongly nil-clean. So, we shall assume hereafter that $U(R) \neq \{1\}$.

Firstly, to prove the “right-to-left” implication, assume that $U(R)$ is a cyclic 2-group. Thus, as commented above, it follows immediately from [4, Corollary 4.10] that R is strongly nil-clean. What remain to show is that $2 = 0$ holds in R . Indeed, since $2 \in \text{Nil}(R)$, one observes that the infinite sequence $\{3, 5, 7, \dots, 2k-1, 2k+1, \dots\}$ will invert in R for any $k \in \mathbb{N}$. But as $U(R)$ is finite, there will exist a natural number k with $2k-1 = 2k+1$, so that $2 = 0$ is really fulfilled.

Secondly, the direct application of [4, Corollary 4.10] gives the “left-to-right” part, as desired. \square

We finish our work in this subsection with the following useful comments which shed some further light on the explored theme.

Remark 2 *For nil-clean rings with finite unit group the above theorem is not longer true: in fact, as an example we can consider the 2×2 matrix ring $M_2(\mathbb{Z}_2)$ which, in accordance with [2], is nil-clean but surely not strongly nil-clean (however, it is strongly π -regular being finite). This suggests to extract even the more general claim that nil-clean rings with finite unit group are strongly π -regular of characteristic 2. In fact, as unipotents (= the sum of 1 and a nilpotent) are always units, it readily follows that the set of nilpotents is also finite and so the ring is with bounded index of nilpotence. We, therefore, can apply [6, Corollary 3.12] to get the wanted claim. That $\text{char}(R) = 2$ follows now in the same manner as in the proof of Theorem 2.*

In closing, we pose a few intriguing problems of some interest and importance which immediately arise.

Problem 1. If R is a nil-clean ring with bounded $U(R)$, does it follow that R is (strongly) π -regular?

Problem 2. If R is a nil-clean ring of characteristic 2 and $U(R)$ is a p -group (or, respectively, a $2p$ -group) for some prime p , is it true that R is (strongly) π -regular?

For eventual counterexamples in case we have dropped some of the requirements, see Examples 3.1 and 3.2 from [11].

In regard to both sections explored above, one may state the following:

Problem 3. Is any nil-clean ring R such that its nilpotents are differences of two idempotents always π -regular? In particular, if $J(R) = 0$, is then R necessarily von Neumann regular.

In fact, each such nil-clean ring is of characteristic 2. If the above question holds in the affirmative, this will be in sharp contrast to the recent example by Šter from [11] showing that there is a nil-clean ring which is not π -regular.

Letting $Q\text{Nil}(R)$ be the set of all quasi-nilpotent elements of the ring R , we note that both inclusions $\text{Nil}(R) \subseteq Q\text{Nil}(R)$ and $J(R) \subseteq Q\text{Nil}(R)$ hold. We thereby come in mind to our next question as follows:

Problem 4. Examine those (nil-clean) rings for which the equality $U(R) = 1 + Q\text{Nil}(R)$ is true.

Notice that the condition $U(R) = 1 + \text{Nil}(R) + J(R)$ obviously implies the condition $U(R) = 1 + Q\text{Nil}(R)$, as in the latter situation we shall say that the ring R has *quasi-nilpotent units*.

Acknowledgement

The author is grateful to the referee for his/her careful reading of the manuscript and the competent comments and suggestions made as well as he is obliged to the chief-editor, Prof. Robert Szasz, for the professional editorial management of the current submission.

Funding: The article is partially supported by the Bulgarian National Science Fund under Grant KP-06 No 32/1 of December 07, 2019 and by project PД-08-136/04.02.2020 of the University of Shumen “Konstantin Preslavsky”.

References

- [1] K. I. Beidar, K. C. O'Meara and R. M. Raphael, On uniform diagonalisation of matrices over regular rings and one-accessible regular algebras, *Commun. Algebra*, **32** (2004), 3543–3562.
- [2] S. Breaz, G. Călugăreanu, P. Danchev and T. Micu, Nil-clean matrix rings, *Linear Algebra and Appl.*, (10) **439** (2013), 3115–3119.
- [3] P. V. Danchev, Weakly exchange rings whose units are sums of two idempotents, *Vestnik St. Petersburg Univ., Math., Mech. & Astronomy*, (2) **6** (64) (2019), 265–269.
- [4] P. V. Danchev and T. Y. Lam, Rings with unipotent units, *Publ. Math. Debrecen*, (3-4) **88** (2016), 449–466.
- [5] P. Danchev and J. Matczuk, \mathfrak{n} -Torsion clean rings, *Contemp. Math.*, **727** (2019), 71–82.
- [6] P. Danchev and J. Šter, Generalizing π -regular rings, *Taiwanese J. Math.*, (6) **19** (2015), 1577–1592.
- [7] A. J. Diesl, Nil clean rings, *J. Algebra*, **383** (2013), 197–211.
- [8] R. E. Hartwig and M. S. Putcha, When is a matrix a difference of two idempotents, *Linear and Multilinear Algebra*, (4) **26** (1990), 267–277.
- [9] T.-Y. Lam, *A First Course in Noncommutative Rings*, Second Edition, Graduate Texts in Math., Vol. 131, Springer-Verlag, Berlin-Heidelberg-New York, 2001.
- [10] K. C. O'Meara, *Nilpotents often the difference of two idempotents*, private correspondence on draft privately circulated on March 2018.
- [11] J. Šter, On expressing matrices over \mathbb{Z}_2 as the sum of an idempotent and a nilpotent, *Linear Algebra Appl.*, **544** (2018), 339–349.

Received: December 17, 2019

Generalized operator for Alexander integral operator

H. Özlem Güney

Dicle University, Faculty of Science,
Department of Mathematics,
21280 Diyarbakır-Turkey
email: ozlemg@dicle.edu.tr

Shigeyoshi Owa

Honorary Professor
“1 Decembrie 1918” University
Alba Iulia, Romania
email: shige21@ican.zaq.ne.jp

Abstract. Let T_n be the class of functions f which are defined by a power series

$$f(z) = z + a_{n+1}z^{n+1} + a_{n+2}z^{n+2} + \dots$$

for every z in the closed unit disc \overline{U} . With m different boundary points z_s , ($s = 1, 2, \dots, m$), we consider $\alpha_m \in e^{i\beta} A_{-j-\lambda} f(U)$, here $A_{-j-\lambda}$ is the generalized Alexander integral operator and U is the open unit disc. Applying $A_{-j-\lambda}$, a subclass $B_n(\alpha_m, \beta, \rho; j, \lambda)$ of T_n is defined with fractional integral for functions f . The object of present paper is to consider some interesting properties of f to be in $B_n(\alpha_m, \beta, \rho; j, \lambda)$.

1 Introduction

Let T_n be the class of functions

$$f(z) = z + \sum_{k=n+1}^{\infty} a_k z^k, \quad n \in \mathbb{N} = \{1, 2, 3, \dots\} \quad (1)$$

2010 Mathematics Subject Classification: 30C45, 30C50

Key words and phrases: analytic function, Alexander integral operator, fractional integral, gamma function, Miller and Mocanu lemma

that are analytic in the closed unit disc $\overline{\mathbb{U}} = \{z \in \mathbb{C} : |z| \leq 1\}$. For $f \in T_n$, J.W.Alexander [2] had defined the following the Alexander integral operator $A_{-1}f(z)$ given by

$$A_{-1}f(z) = \int_0^z \frac{f(t)}{t} dt = z + \sum_{k=n+1}^{\infty} \frac{a_k}{k} z^k. \quad (2)$$

The above the Alexander integral operator was applied for some subclasses of analytic functions in the open unit disc $\mathbb{U} = \{z \in \mathbb{C} : |z| < 1\}$ by M.Acu [1] and by K. Kugita et al. [4].

For the above the Alexander integral operator $A_{-1}f(z)$, we consider

$$A_{-j}f(z) = A_{-j+1}(A_{-1}f(z)) = z + \sum_{k=n+1}^{\infty} \frac{a_k}{k^j} z^k, \quad j \in \mathbb{N} \quad (3)$$

where $A_0f(z) = f(z)$.

From the various definitions of fractional calculus of $f \in T_n$ (that is, fractional integrals and fractional derivatives) given in the literature, we would like to recall here the following definitions for fractional calculus which were used by Owa [7] and Owa and Srivastava [8].

Definition 1 *The fractional integral of order λ for $f \in T_n$ is defined by*

$$D_z^{-\lambda}f(z) = \frac{1}{\Gamma(\lambda)} \int_0^z \frac{f(t)}{(z-t)^{1-\lambda}} dt, \quad (\lambda > 0) \quad (4)$$

where f is an analytic function in a simply-connected region of the z -plane containing the origin, and the multiplicity of $(z-t)^{\lambda-1}$ is removed by requiring $\log(z-t)$ to be real when $z-t > 0$ and Γ is the Gamma function.

With the above definition, we know that

$$D_z^{-\lambda}f(z) = \frac{1}{\Gamma(2+\lambda)} z^{1+\lambda} + \sum_{k=n+1}^{\infty} \frac{k!}{\Gamma(k+1+\lambda)} a_k z^{k+\lambda} \quad (5)$$

for $\lambda > 0$ and $f \in T_n$. Further applying the fractional integral for $f \in T_n$, we define a new operator $A_{-\lambda}f(z)$ given by

$$A_{-\lambda}f(z) = \frac{\Gamma\left(\frac{3+\lambda}{2}\right)}{\Gamma\left(\frac{3-\lambda}{2}\right)} z^{\frac{1-\lambda}{2}} D_z^{-\lambda} \left(z^{\frac{-1-\lambda}{2}} f(z) \right), \quad (6)$$

where $0 \leq \lambda \leq 1$. If $\lambda = 0$, then (6) becomes $A_0 f(z) = f(z)$ and if $\lambda = 1$, then (6) leads us that

$$A_{-1} f(z) = D_z^{-1} \left(\frac{f(z)}{z} \right) = \int_0^z \frac{f(t)}{t} dt. \quad (7)$$

With this integral operator, we know

$$A_{-j-\lambda} f(z) = A_{-j} (A_{-\lambda} f(z)) \quad (8)$$

where $j \in \mathbb{N}$ and $0 \leq \lambda \leq 1$. This operator $A_{-j-\lambda} f(z)$ is the generalization of the Alexander integral operator $A_{-1} f(z)$. Here, we note that

$$A_{-\lambda} f(z) = z + \sum_{k=n+1}^{\infty} \frac{\Gamma\left(\frac{3+\lambda}{2}\right) \Gamma\left(\frac{2k+1-\lambda}{2}\right)}{\Gamma\left(\frac{3-\lambda}{2}\right) \Gamma\left(\frac{2k+1+\lambda}{2}\right)} a_k z^k \quad (9)$$

and

$$A_{-j-\lambda} f(z) = z + \sum_{k=n+1}^{\infty} \frac{\Gamma\left(\frac{3+\lambda}{2}\right) \Gamma\left(\frac{2k+1-\lambda}{2}\right)}{\Gamma\left(\frac{3-\lambda}{2}\right) \Gamma\left(\frac{2k+1+\lambda}{2}\right)} k^j a_k z^k \quad (10)$$

where $j \in \mathbb{N}$ and $0 \leq \lambda \leq 1$. From the above, we know that

$$A_{-j-\lambda} f(z) = A_{-j} (A_{-\lambda} f(z)) = A_{-\lambda} (A_{-j} f(z)) \quad (11)$$

for $f \in T_n$. For m different boundary points $z_s (s = 1, 2, 3, \dots, m)$ with $|z_s| = 1$, we consider

$$\alpha_m = \frac{1}{m} \sum_{s=1}^m \frac{A_{-j-\lambda} f(z_s)}{z_s}, \quad (12)$$

where $\alpha_m \in e^{i\beta} A_{-j-\lambda} f(\mathbb{U})$, $\alpha_m \neq 1$ and $-\frac{\pi}{2} \leq \beta \leq \frac{\pi}{2}$. For such α_m , if $f \in T_n$ satisfies

$$\left| \frac{e^{i\beta} \frac{A_{-j-\lambda} f(z)}{z} - \alpha_m}{e^{i\beta} - \alpha_m} - 1 \right| < \rho, \quad z \in \mathbb{U} \quad (13)$$

for some real $\rho > 0$, we say that the function f belongs to the subclass $B_n(\alpha_m, \beta, \rho; j, \lambda)$ of T_n . With this definition for the class $B_n(\alpha_m, \beta, \rho; j, \lambda)$, we see that the condition (13) is equivalent to

$$\left| \frac{A_{-j-\lambda} f(z)}{z} - 1 \right| < \rho \left| e^{i\beta} - \alpha_m \right|, \quad z \in \mathbb{U}. \quad (14)$$

If we consider the function $f \in T_n$ given by

$$f(z) = z + \frac{\Gamma\left(\frac{3-\lambda}{2}\right) \Gamma\left(\frac{2n+3+\lambda}{2}\right)}{\Gamma\left(\frac{3+\lambda}{2}\right) \Gamma\left(\frac{2n+3-\lambda}{2}\right)} \rho (e^{i\beta} - \alpha_m) (n+1)^j z^{n+1} \quad (15)$$

then f satisfies

$$\left| \frac{A_{-j-\lambda} f(z)}{z} - 1 \right| = \rho \left| e^{i\beta} - \alpha_m \right| |z|^n < \rho \left| e^{i\beta} - \alpha_m \right|, \quad z \in \mathbb{U}. \quad (16)$$

Therefore, f given by (15) belongs to the class $B_n(\alpha_m, \beta, \rho; j, \lambda)$.

Discussing our problems for $f \in B_n(\alpha_m, \beta, \rho; j, \lambda)$, we have to introduce the following lemma due to S. S. Miller and P. T. Mocanu [5, 6] (also, due to I. S. Jack [3]).

Lemma 1 *Let the function w given by*

$$w(z) = a_n z^n + a_{n+1} z^{n+1} + a_{n+2} z^{n+2} + \dots, \quad (n \in \mathbb{N}) \quad (17)$$

be analytic in \mathbb{U} with $w(0) = 0$. If $|w(z)|$ attains its maximum value on the circle $|z| = r$ at a point z_0 , ($0 < |z_0| < 1$) then there exists a real number $k \geq n$ such that

$$\frac{z_0 w'(z_0)}{w(z_0)} = k \quad (18)$$

and

$$\operatorname{Re} \left(1 + \frac{z_0 w''(z_0)}{w'(z_0)} \right) \geq k. \quad (19)$$

2 Properties of functions concerning with the class $B_n(\alpha_m, \beta, \rho; j, \lambda)$

Our first property for $f \in T_n$ is as follows.

Theorem 1 *If $f \in T_n$ satisfies*

$$\left| \frac{A_{-j-\lambda+1} f(z)}{A_{-j-\lambda} f(z)} - 1 \right| < \frac{|e^{i\beta} - \alpha_m| n \rho}{1 + |e^{i\beta} - \alpha_m| \rho}, \quad z \in \mathbb{U} \quad (20)$$

for some α_m defined by (12) with $\alpha_m \neq 1$ such that $z_s \in \partial\mathbb{U}$ ($s = 1, 2, 3, \dots, m$), and for some real $\rho > 1$, then

$$\left| \frac{A_{-j-\lambda} f(z)}{z} - 1 \right| < \rho \left| e^{i\beta} - \alpha_m \right|, \quad z \in \mathbb{U} \quad (21)$$

that is, $f \in B_n(\alpha_m, \beta, \rho; j, \lambda)$.

Proof. We introduce the function w by

$$w(z) = \frac{e^{i\beta} \frac{A_{-j-\lambda}f(z)}{z} - \alpha_m}{e^{i\beta} - \alpha_m} - 1 = \frac{e^{i\beta}}{e^{i\beta} - \alpha_m} \left\{ \sum_{k=n+1}^{\infty} \frac{\Gamma\left(\frac{3+\lambda}{2}\right) \Gamma\left(\frac{2k+1-\lambda}{2}\right)}{\Gamma\left(\frac{3-\lambda}{2}\right) \Gamma\left(\frac{2k+1+\lambda}{2}\right) k^j} \alpha_k z^{k-1} \right\}. \quad (22)$$

Then, w is analytic in \mathbb{U} with $w(0) = 0$ and

$$\frac{A_{-j-\lambda}f(z)}{z} = 1 + (1 - e^{-i\beta} \alpha_m)w(z). \quad (23)$$

It follows from the above that

$$\frac{z(A_{-j-\lambda}f(z))'}{A_{-j-\lambda}f(z)} - 1 = \frac{(1 - e^{-i\beta} \alpha_m)zw'(z)}{1 + (1 - e^{-i\beta} \alpha_m)w(z)}. \quad (24)$$

Note that

$$z(A_{-j-\lambda}f(z))' = A_{-j-\lambda+1}f(z). \quad (25)$$

So, (24) is the same as

$$\frac{A_{-j-\lambda+1}f(z)}{A_{-j-\lambda}f(z)} - 1 = \frac{(1 - e^{-i\beta} \alpha_m)zw'(z)}{1 + (1 - e^{-i\beta} \alpha_m)w(z)}. \quad (26)$$

Thus, our condition (20) gives that

$$\left| \frac{A_{-j-\lambda+1}f(z)}{A_{-j-\lambda}f(z)} - 1 \right| = \left| \frac{(1 - e^{-i\beta} \alpha_m)zw'(z)}{1 + (1 - e^{-i\beta} \alpha_m)w(z)} \right| < \frac{|e^{i\beta} - \alpha_m|n\rho}{1 + |e^{i\beta} - \alpha_m|\rho}. \quad (27)$$

Now, we suppose that there exists a point z_0 , ($0 < |z_0| < 1$) such that

$$\max\{|w(z)|; |z| \leq |z_0|\} = |w(z_0)| = \rho > 1. \quad (28)$$

Then, we can write that $w(z_0) = \rho e^{i\theta}$, ($0 \leq \theta \leq 2\pi$) and $z_0 w'(z_0) = kw(z_0)$, ($k \geq n$) by Lemma 1. For such a point z_0 , ($0 < |z_0| < 1$) we see that

$$\begin{aligned} \left| \frac{A_{-j-\lambda+1}f(z_0)}{A_{-j-\lambda}f(z_0)} - 1 \right| &= \left| \frac{(1 - e^{-i\beta} \alpha_m)z_0 w'(z_0)}{1 + (1 - e^{-i\beta} \alpha_m)w(z_0)} \right| \\ &= \left| \frac{(1 - e^{-i\beta} \alpha_m)k\rho}{1 + (1 - e^{-i\beta} \alpha_m)\rho e^{i\theta}} \right| \\ &\geq \frac{|1 - e^{-i\beta} \alpha_m|n\rho}{1 + |1 - e^{-i\beta} \alpha_m|\rho} \\ &= \frac{|e^{i\beta} - \alpha_m|n\rho}{1 + |e^{i\beta} - \alpha_m|\rho}. \end{aligned} \quad (29)$$

Since (29) contradicts our condition (20), we know that there is no z_0 , ($0 < |z_0| < 1$) such that $|w(z_0)| = \rho > 1$. Therefore, using $|w(z)| < \rho$ for all $z \in \mathbb{U}$, we have that

$$|w(z)| = \left| \frac{e^{i\beta} \left(\frac{A_{-j-\lambda}f(z)}{z} - 1 \right)}{e^{i\beta} - \alpha_n} \right| < \rho, \quad z \in \mathbb{U}, \quad (30)$$

that is, that

$$\left| \frac{A_{-j-\lambda}f(z)}{z} - 1 \right| < \rho \left| e^{i\beta} - \alpha_n \right|, \quad z \in \mathbb{U}. \quad (31)$$

This completes the proof of the theorem. \square

Example 1 We consider the function $f \in T_n$ given by

$$f(z) = z + a_{n+1}z^{n+1}, \quad z \in \mathbb{U}. \quad (32)$$

Then, we see that

$$\begin{aligned} \left| \frac{A_{-j-\lambda+1}f(z)}{A_{-j-\lambda}f(z)} - 1 \right| &= \left| \frac{P(n, j, \lambda)na_{n+1}z^n}{1 + P(n, j, \lambda)a_{n+1}z^n} \right| \\ &< \frac{nP(n, j, \lambda)|a_{n+1}|}{1 - P(n, j, \lambda)|a_{n+1}|}, \quad z \in \mathbb{U}, \end{aligned} \quad (33)$$

where

$$0 < |a_{n+1}| < \frac{1 - P(n, j, \lambda)}{P(n, j, \lambda)} \quad (34)$$

and

$$P(n, j, \lambda) = \frac{\Gamma\left(\frac{3+\lambda}{2}\right) \Gamma\left(\frac{2n+3-\lambda}{2}\right)}{\Gamma\left(\frac{3-\lambda}{2}\right) \Gamma\left(\frac{2n+3+\lambda}{2}\right) (n+1)^j}. \quad (35)$$

Now, we consider five boundary points

$$z_1 = e^{-i \frac{\arg(a_{n+1})}{n}} \quad (36)$$

$$z_2 = e^{i \frac{\pi - 6\arg(a_{n+1})}{6n}} \quad (37)$$

$$z_3 = e^{i \frac{\pi - 4\arg(a_{n+1})}{4n}} \quad (38)$$

$$z_4 = e^{i \frac{\pi - 3\arg(a_{n+1})}{3n}} \quad (39)$$

and

$$z_5 = e^{i \frac{\pi - 2 \arg(a_{n+1})}{2n}}. \quad (40)$$

For such z_s ($s = 1, 2, 3, 4, 5$), we have that

$$\frac{A_{-j-\lambda}f(z_1)}{z_1} = 1 + P(n, j, \lambda)|a_{n+1}|, \quad (41)$$

$$\frac{A_{-j-\lambda}f(z_2)}{z_2} = 1 + P(n, j, \lambda)|a_{n+1}| \frac{\sqrt{3} + i}{2}, \quad (42)$$

$$\frac{A_{-j-\lambda}f(z_3)}{z_3} = 1 + P(n, j, \lambda)|a_{n+1}| \frac{\sqrt{2}(1 + i)}{2}, \quad (43)$$

$$\frac{A_{-j-\lambda}f(z_4)}{z_4} = 1 + P(n, j, \lambda)|a_{n+1}| \frac{1 + \sqrt{3}i}{2}, \quad (44)$$

and

$$\frac{A_{-j-\lambda}f(z_5)}{z_5} = 1 + P(n, j, \lambda)|a_{n+1}|i. \quad (45)$$

It follows from the above that

$$\alpha_5 = \frac{1}{5} \sum_{s=1}^5 \frac{A_{-j-\lambda}f(z_s)}{z_s} = 1 + \frac{(3 + \sqrt{2} + \sqrt{3})P(n, j, \lambda)|a_{n+1}|(1 + i)}{10} \quad (46)$$

and that

$$\left| 1 - e^{-i\beta} \alpha_5 \right| = \frac{\sqrt{2}(3 + \sqrt{2} + \sqrt{3})P(n, j, \lambda)|a_{n+1}|}{10} \quad (47)$$

with $\beta = 0$. For such α_5 and β , we consider $\rho > 1$ with

$$\frac{nP(n, j, \lambda)|a_{n+1}|}{1 - P(n, j, \lambda)|a_{n+1}|} \leq \frac{|e^{i\beta} - \alpha_5|n\rho}{1 + |e^{i\beta} - \alpha_5|\rho}. \quad (48)$$

This gives us that

$$\rho \geq \frac{10}{\sqrt{2}(3 + \sqrt{2} + \sqrt{3})(1 - (1 + |a_{n+1}|)P(n, j, \lambda))} > \frac{10}{\sqrt{2}(3 + \sqrt{2} + \sqrt{3})} > 1. \quad (49)$$

For such α_5 and $\rho > 1$, the function f satisfies

$$\left| \frac{A_{-j-\lambda}f(z)}{z} - 1 \right| < P(n, j, \lambda)|a_{n+1}| \leq \rho |e^{i\beta} - \alpha_5|, \quad z \in \mathbb{U}.$$

Next, we derive the following theorem.

Theorem 2 *If $f \in T_n$ satisfies*

$$\left| \left(\frac{A_{-j-\lambda+1}f(z)}{A_{-j-\lambda}f(z)} - 1 \right) \left(\frac{A_{-j-\lambda}f(z)}{z} - 1 \right) \right| < \frac{|e^{i\beta} - \alpha_m|^2 n \rho^2}{1 + |e^{i\beta} - \alpha_m| \rho}, \quad z \in \mathbb{U} \quad (50)$$

for some α_m defined by (12) with $\alpha_m \neq 1$ and for some real $\rho > 1$, then

$$\left| \frac{A_{-j-\lambda}f(z)}{z} - 1 \right| < \rho |e^{i\beta} - \alpha_m|, \quad z \in \mathbb{U} \quad (51)$$

that is, $f \in B_n(\alpha_m, \beta, \rho; j, \lambda)$.

Proof. Define the function w by (22). Applying (25), our condition (50) leads us that

$$\begin{aligned} \left| \left(\frac{A_{-j-\lambda+1}f(z)}{A_{-j-\lambda}f(z)} - 1 \right) \left(\frac{A_{-j-\lambda}f(z)}{z} - 1 \right) \right| &= \left| \frac{(1 - e^{-i\beta} \alpha_m)^2 z w(z) w'(z)}{1 + (1 - e^{-i\beta} \alpha_m) w(z)} \right| \\ &\leq \frac{|e^{i\beta} - \alpha_m|^2 n \rho^2}{1 + |e^{i\beta} - \alpha_m| \rho}, \quad z \in \mathbb{U}. \end{aligned} \quad (52)$$

Suppose that there exists a point z_0 , ($0 < |z_0| < 1$) such that

$$\max\{|w(z)|; |z| \leq |z_0|\} = |w(z_0)| = \rho > 1. \quad (53)$$

Then, applying Lemma 1, we write that $w(z_0) = \rho e^{i\theta}$, ($0 \leq \theta \leq 2\pi$) and $z_0 w'(z_0) = k w(z_0)$, ($k \geq n$). This shows us that

$$\begin{aligned} \left| \left(\frac{A_{-j-\lambda+1}f(z)}{A_{-j-\lambda}f(z)} - 1 \right) \left(\frac{A_{-j-\lambda}f(z)}{z} - 1 \right) \right| &= \left| \frac{(1 - e^{-i\beta} \alpha_m)^2 z_0 w(z_0) w'(z_0)}{1 + (1 - e^{-i\beta} \alpha_m) w(z_0)} \right| \\ &= \frac{|e^{i\beta} - \alpha_m|^2 \rho^2 k}{|1 + (1 - e^{-i\beta} \alpha_m) \rho e^{i\theta}|} \\ &\geq \frac{|e^{i\beta} - \alpha_m|^2 n \rho^2}{1 + |e^{i\beta} - \alpha_m| \rho} \end{aligned} \quad (54)$$

which contradicts our condition (50). Thus there is no z_0 , ($0 < |z_0| < 1$) such that $|w(z_0)| = \rho > 1$. This shows us that

$$\left| \left(\frac{A_{-j-\lambda}f(z)}{z} - 1 \right) \right| < \rho |e^{i\beta} - \alpha_m|, \quad z \in \mathbb{U}. \quad (55)$$

□

Example 2 Consider a function $f \in T_n$ given by

$$f(z) = z + a_{n+1}z^{n+1}, z \in \mathbb{U} \quad (56)$$

with $0 < |a_{n+1}| < \frac{1}{P(n, j, \lambda)}$, where $P(n, j, \lambda)$ is given by (35). It follows that

$$\begin{aligned} \left| \left(\frac{A_{-j-\lambda+1}f(z)}{A_{-j-\lambda}f(z)} - 1 \right) \left(\frac{A_{-j-\lambda}f(z)}{z} - 1 \right) \right| &= \left| \frac{nP(n, j, \lambda)^2 a_{n+1}^2 z^{2n}}{1 + P(n, j, \lambda) a_{n+1} z^n} \right| \\ &< \frac{nP(n, j, \lambda)^2 |a_{n+1}|^2}{1 - P(n, j, \lambda) |a_{n+1}|}, \quad z \in \mathbb{U}. \end{aligned} \quad (57)$$

Considering five boundary points z_1, z_2, z_3, z_4 and z_5 in Example 1, we see that

$$\left| e^{i\beta} - \alpha_5 \right| = \frac{\sqrt{2}(3 + \sqrt{2} + \sqrt{3})P(n, j, \lambda)|a_{n+1}|}{10} \quad (58)$$

with $\beta = 0$. If we consider $\rho > 1$ such that

$$\frac{nP(n, j, \lambda)^2 |a_{n+1}|^2 |z|}{1 - P(n, j, \lambda) |a_{n+1}|} \leq \frac{|e^{i\beta} - \alpha_5|^2 n \rho^2}{1 + |e^{i\beta} - \alpha_5| \rho}, \quad (59)$$

then ρ satisfies

$$\rho \geq \frac{10}{\sqrt{2}(3 + \sqrt{2} + \sqrt{3})P(n, j, \lambda)|a_{n+1}|} > 1. \quad (60)$$

For such α_5 and ρ , f satisfies

$$\left| \frac{A_{-j-\lambda}f(z)}{z} - 1 \right| < P(n, j, \lambda) |a_{n+1}| \leq \rho |e^{i\beta} - \alpha_5|, \quad z \in \mathbb{U}. \quad (61)$$

Our next result reads as follows.

Theorem 3 If $f \in T_n$ satisfies

$$\left| \frac{A_{-j-\lambda+p}f(z)}{z} - 1 \right| < \rho |e^{i\beta} - \alpha_m| (n+1), \quad z \in \mathbb{U}. \quad (62)$$

for some α_m defined by (12) with $\alpha_m \neq 1$ and for some real $\rho > 1$, then

$$\left| \frac{A_{-j-\lambda+p-1}f(z)}{z} - 1 \right| < \rho |e^{i\beta} - \alpha_m|, \quad z \in \mathbb{U} \quad (63)$$

where $p = 0, 1, 2, \dots, j$.

Proof. We consider the function w defined by

$$\begin{aligned} w(z) &= \frac{e^{i\beta} \frac{A_{-j-\lambda+p-1}f(z)}{z} - \alpha_m}{e^{i\beta} - \alpha_m} - 1 \\ &= \frac{e^{i\beta}}{e^{i\beta} - \alpha_m} \left\{ \sum_{k=n+1}^{\infty} \frac{\Gamma\left(\frac{3+\lambda}{2}\right) \Gamma\left(\frac{2k+1-\lambda}{2}\right)}{\Gamma\left(\frac{3-\lambda}{2}\right) \Gamma\left(\frac{2k+1+\lambda}{2}\right) k^{j-p+1}} a_k z^{k-1} \right\}. \end{aligned} \quad (64)$$

Thus w is analytic in \mathbb{U} , $w(0) = 0$, and

$$A_{-j-\lambda+p-1}f(z) = z + (1 - e^{-i\beta} \alpha_m)zw(z). \quad (65)$$

Noting that

$$\begin{aligned} A_{-j-\lambda+p}f(z) &= z(A_{-j-\lambda+p-1}f(z))' \\ &= z \left\{ 1 + (1 - e^{-i\beta} \alpha_m)w(z) \left(1 + \frac{zw'(z)}{w(z)} \right) \right\}, \end{aligned} \quad (66)$$

we have that

$$\begin{aligned} \left| \frac{A_{-j-\lambda+p}f(z)}{z} - 1 \right| &= \left| 1 - e^{-i\beta} \alpha_m \right| |w(z)| \left| 1 + \frac{zw'(z)}{w(z)} \right| \\ &< \rho \left| e^{i\beta} - \alpha_m \right| (n+1), \quad z \in \mathbb{U} \end{aligned} \quad (67)$$

by the condition (62). Suppose that there exists a point z_0 , ($0 < |z_0| < 1$) such that

$$\max\{|w(z)|; |z| \leq |z_0|\} = |w(z_0)| = \rho > 1. \quad (68)$$

Then, letting $w(z_0) = \rho e^{i\theta}$, ($0 \leq \theta \leq 2\pi$) and $z_0 w'(z_0) = kw(z_0)$, ($k \geq n$) with Lemma 1, we see that

$$\left| \frac{A_{-j-\lambda+p}f(z_0)}{z_0} - 1 \right| = \rho \left| e^{i\beta} - \alpha_m \right| (k+1) \geq \rho \left| e^{i\beta} - \alpha_m \right| (n+1). \quad (69)$$

This contradicts the inequality (67). Therefore, we don't have any $z_0 \in \mathbb{U}$ such that $|w(z_0)| = \rho > 1$. This shows us that

$$|w(z)| = \left| \frac{\alpha_m}{e^{i\beta} - \alpha_m} \left(\frac{A_{-j-\lambda+p-1}f(z)}{z} - 1 \right) \right| < \rho, \quad z \in \mathbb{U}, \quad (70)$$

that is, that

$$\left| \frac{A_{-j-\lambda+p-1}f(z)}{z} - 1 \right| < \rho |e^{i\beta} - \alpha_m|, \quad z \in \mathbb{U}. \quad (71)$$

This completes the proof of our theorem. \square

Corollary 1 If $f \in T_n$ satisfies

$$\left| \frac{A_{-j-\lambda+p}f(z)}{z} - 1 \right| < \rho |e^{i\beta} - \alpha_m| (n+1)^p, \quad z \in \mathbb{U} \quad (72)$$

for some α_m given by (12) with $\alpha_m \neq 1$, and for some real $\rho > 1$, then

$$\left| \frac{A_{-j-\lambda}f(z)}{z} - 1 \right| < \rho |e^{i\beta} - \alpha_m|, \quad z \in \mathbb{U} \quad (73)$$

where $p = 0, 1, 2, \dots, j$.

Proof. With Theorem 3, we say that if $f \in T_n$ satisfies

$$\left| \frac{A_{-j-\lambda+p}f(z)}{z} - 1 \right| < \rho |e^{i\beta} - \alpha_m| (n+1)^p, \quad z \in \mathbb{U}, \quad (74)$$

then

$$\left| \frac{A_{-j-\lambda+p-1}f(z)}{z} - 1 \right| < \rho |e^{i\beta} - \alpha_m| (n+1)^{p-1}, \quad z \in \mathbb{U}. \quad (75)$$

Further, we have that

$$\left| \frac{A_{-j-\lambda+p-2}f(z)}{z} - 1 \right| < \rho |e^{i\beta} - \alpha_m| (n+1)^{p-2}, \quad z \in \mathbb{U}, \quad (76)$$

from (75). Finally, we obtain that

$$\left| \frac{A_{-j-\lambda}f(z)}{z} - 1 \right| < \rho |e^{i\beta} - \alpha_m|, \quad z \in \mathbb{U}. \quad (77)$$

□

Example 3 Consider the function $f \in T_n$ given by

$$f(z) = z + a_{n+1}z^{n+1}, \quad z \in \mathbb{U}. \quad (78)$$

Since

$$A_{-j-\lambda+p}f(z) = z + \frac{\Gamma\left(\frac{3+\lambda}{2}\right) \Gamma\left(\frac{2n+3-\lambda}{2}\right)}{\Gamma\left(\frac{3-\lambda}{2}\right) \Gamma\left(\frac{2n+3+\lambda}{2}\right) (n+1)^{j-p+2}} a_{n+1}z^{n+1}, \quad (79)$$

we have

$$\left| \frac{A_{-j-\lambda+p}f(z)}{z} - 1 \right| = \left| P(n, j, \lambda)(n+1)^{p-2} a_{n+1}z^n \right| < P(n, j, \lambda)(n+1)^{p-2} |a_{n+1}| \quad (80)$$

where

$$0 < |a_{n+1}| < \frac{1}{P(n, j, \lambda)} \quad (81)$$

and $P(n, j, \lambda)$ is given by (35).

Consider five boundary points z_1, z_2, z_3, z_4 and z_5 in Example 1. Then α_5 satisfies (46) and $|1 - e^{-i\beta}\alpha_5|$ satisfies (47) for $\beta = 0$. For such α_5 and β , we consider $\rho > 1$ by

$$\left| \frac{A_{-j-\lambda+p}f(z)}{z} - 1 \right| < P(n, j, \lambda)(n+1)^{p-2}|a_{n+1}| \leq \rho \left| e^{i\beta} - \alpha_5 \right| (n+1)^{p-2}, \quad z \in \mathbb{U}, \quad (82)$$

Then ρ satisfies

$$\rho \geq \frac{P(n, j, \lambda)|a_{n+1}|}{|e^{i\beta} - \alpha_5|} = \frac{10}{\sqrt{2}(3 + \sqrt{2} + \sqrt{3})} > 1. \quad (83)$$

With the above α_5 and ρ , we have

$$\left| \frac{A_{-j-\lambda}f(z)}{z} - 1 \right| < P(n, j, \lambda)|a_{n+1}| \leq \rho \left| e^{i\beta} - \alpha_5 \right|, \quad z \in \mathbb{U}. \quad (84)$$

References

- [1] M. Acu, Some preserving properties of the generalized Alexander integral operator, *General Math.*, **10** (2002), 37–46.
- [2] J. W. Alexander, Functions which map the interior of the unit circle upon simple regions, *Ann. of Math.*, Second Series, **17** (1915), 12–22.
- [3] I. S. Jack, Functions starlike and convex of order α , *J. London Math. Soc.*, **2** (1971), 469–474.
- [4] K. Kugita, K. Kuroki and S. Owa, On (α, δ) -neighborhood defined by a new operator for certain analytic functions, *Far East J. Math. Sci.*, **47** (2010), 1–12.
- [5] S. S. Miller and P. T. Mocanu, Second order differential inequalities in the complex plane, *J. Math. Anal. Appl.*, **65** (1978), 289–305.
- [6] S. S. Miller and P. T. Mocanu, *Differential Subordinations: Theory and Applications*, Marcel Dekker Inc. New York, (2000).

- [7] S. Owa, On the distortion theorems I, *Kyungpook Math. J.*, **18** (1978), 53–59.
- [8] S. Owa and H. M. Srivastava, Univalent and starlike generalized hypergeometric functions, *Canad. J. Math.*, **39** (1987), 1057–1077.

Received: April 29, 2020



On $\lambda^D - R_0$ and $\lambda^D - R_1$ spaces

Sarhad F. Namiq

Department of Mathematics,
College of Education,
University of Garmian,
Kurdistan-Region, Iraq
email: sarhad1983@gmail.com

Ennis Rosas

Departamento de Ciencias
Naturales y Exactas,
Universidad de la Costa, Barranquilla,
Colombia & Departamento de
Matemáticas, Universidad de Oriente,
Cumaná, Venezuela
email: ennisrafael@gmail.com

Abstract. In this paper we introduce the new types of separation axioms called $\lambda^D - R_0$ and $\lambda^D - R_1$ spaces, by using λ^D -open set. The notion $\lambda^D - R_0$ and $\lambda^D - R_1$ spaces are introduced and some of their properties are investigated.

1 Introduction

In 1943, the notion of R_0 topological space was introduced by Shanin [6]. Later, Davis [3] rediscovered it and studied some properties of this weak separation axiom. In the same paper, Davis also introduced the notion of R_1 topological space which are independent of both T_0 and T_1 , but strictly weaker than T_2 . The notion of λ -open (λ^* -open) sets was introduced by Alais B. Khalaf and Sarhad F. Namiq [1]. The notion of λ^D -open sets was introduced by Sarhad F. Namiq [5]. In this paper, we continue the study of the above mentioned classes of topological spaces satisfying these axioms by introducing two more notions in terms of λ^D -open sets called $\lambda^D - R_0$ and $\lambda^D - R_1$.

2010 Mathematics Subject Classification: 54A05, 54B05, 54C05

Key words and phrases: λ^D -open set; $\lambda^D - R_0$ and $\lambda^D - R_1$

2 Preliminaries

Throughout, X denote a topological space. Let A be a subset of X , the closure and the interior of A are denoted by $\text{Cl}(A)$ and $\text{Int}(A)$ respectively. A subset A of a topological space (X, τ) is said to be dense set [7] if $\text{Cl}(A) = X$. A subset A of a topological space (X, τ) is said to be semi open [4] if $A \subseteq \text{Cl}(\text{Int}(A))$. The complement of a semi open set is said to be semi closed [4]. The family of all semi open (resp. semi closed) sets in a topological space (X, τ) is denoted by $\text{SO}(X, \tau)$ or $\text{SO}(X)$ (resp. $\text{SC}(X, \tau)$ or $\text{SC}(X)$). We consider λ as a function defined on $\text{SO}(X)$ into $\mathcal{P}(X)$ and $\lambda : \text{SO}(X) \rightarrow \mathcal{P}(X)$ is called an s-operation if $V \subseteq \lambda(V)$ for each non-empty semi open set V . It is assumed that $\lambda(\emptyset) = \emptyset$ and $\lambda(X) = X$ for any s-operation λ .

Definition 1 [1] *Let (X, τ) be a topological space and $\lambda : \text{SO}(X) \rightarrow \mathcal{P}(X)$ be an s-operation, then a subset A of X is called a λ^* -open set which is equivalent to λ -open set, if for each $x \in A$, there exists a semi open set U such that $x \in U$ and $\lambda(U) \subseteq A$. The complement of a λ^* -open set is said to be λ^* -closed set which is equivalent to λ -closed set. The family of all λ^* -open (resp., λ^* -closed) subsets of a topological space (X, τ) is denoted by or $\text{SO}_\lambda(X)$ (resp. $\text{SC}_\lambda(X, \tau)$ or $\text{SC}_\lambda(X)$).*

Definition 2 [5] *Let (X, τ) be a topological space and $\lambda : \text{SO}(X) \rightarrow \mathcal{P}(X)$ be an s-operation, then a λ^* -open subset A of X is called a λ^D -open set, if for each $x \in A$, there exists a dense set D such that $x \in D \subseteq A$. The complement of a λ^D -open set is said to be λ^D -closed. The family of all λ^D -open (resp., λ^D -closed) subsets of a topological space (X, τ) is denoted by or $\text{SO}_{\lambda^D}(X)$ (resp. $\text{SC}_{\lambda^D}(X, \tau)$ or $\text{SC}_{\lambda^D}(X)$).*

Example 1 *Let $X = \{a, b, c, d\}$ with topology $\tau = \{\emptyset, X, \{a\}, \{b\}, \{a, b\}, \{a, b, c\}\}$. The $\text{SO}(X) = \{\emptyset, X, \{a\}, \{b\}, \{a, b\}, \{a, c\}, \{a, d\}, \{b, c\}, \{b, d\}, \{a, b, c\}, \{a, b, d\}, \{b, c, d\}, \{a, c, d\}\}$. Define $\lambda : \text{SO}(X) \rightarrow \mathcal{P}(X)$ as:*

$$\lambda(A) = \begin{cases} A & \text{if } a \in A \\ X & \text{if } a \notin A \end{cases}$$

The $\text{SO}_{\lambda^D}(X) = \{\emptyset, X, \{a, b\}, \{a, b, c\}, \{a, b, d\}\}$.

Definition 3 [5] *Let (X, τ) be a topological space and let A be a subset of X . Then:*

1. *The λ -closure of A (denoted by $\lambda^D\text{Cl}(A)$) is the intersection of all λ^D -closed sets containing A .*

2. The λ -interior of A (denoted by $\lambda^D \text{Int}(A)$) is the union of all λ^D -open sets of X contained in A .

Proposition 1 [5] For each point $x \in X$, $x \in \lambda^D \text{Cl}(A)$ if and only if $V \cap A \neq \emptyset$, for every $V \in \text{SO}_{\lambda^D}(X)$ such that $x \in V$.

3 On $\lambda^D - R_0$ and $\lambda^D - R_1$ spaces

We introduce the following definitions.

Definition 4 For any s -operation $\lambda : \text{SO}(X) \rightarrow \mathcal{P}(X)$ and any subset A of a space (X, τ) the λ^D -kernel of A , denoted by $\lambda^D \text{Ker}(A)$ is defined as:

$$\lambda^D \text{Ker}(A) = \cap \{G \in \text{SO}_{\lambda^D}(X) : A \subseteq G\}.$$

Lemma 1 Let (X, τ) be a topological space, $A \subseteq X$ and $\lambda : \text{SO}(X) \rightarrow \mathcal{P}(X)$ be an s -operation. Then $\lambda^D \text{Ker}(A) = \{x \in X : \lambda^D \text{Cl}(\{x\}) \cap A \neq \emptyset\}$.

Proof. Let $x \in \lambda^D \text{Ker}(A)$ such that $\lambda^D \text{Cl}(\{x\}) \cap A = \emptyset$. Since $x \notin X \setminus \lambda^D \text{Cl}(\{x\})$ which is a λ^D -open set containing A . Thus $x \notin \lambda^D \text{Ker}(A)$ a contradiction.

Conversely, let $x \in X$ be such that $\lambda^D \text{Cl}(\{x\}) \cap A \neq \emptyset$. If possible, let $x \notin \lambda^D \text{Ker}(A)$. Then there exist a λ^D -open set G such that $x \notin G$ and $A \subseteq G$. Let $y \in \lambda^D \text{Cl}(\{x\}) \cap A$. This implies that $y \in \lambda^D \text{Cl}(\{x\})$ and $y \in G$, which gives $x \in G$, a contradiction. \square

Theorem 1 Let (X, τ) be a topological space, A and B be subsets of X . Then:

- (1) $x \in \lambda^D \text{Ker}(A)$ if and only if $A \cap F \neq \emptyset$; for any λ^D -closed set F containing x .
- (2) $A \subseteq \lambda^D \text{Ker}(A)$ and $A = \lambda^D \text{Ker}(A)$ if A is λ^D -open.
- (3) If $A \subseteq B$, then $\lambda^D \text{Ker}(A) \subseteq \lambda^D \text{Ker}(B)$.

Proof. Obvious. \square

Definition 5 Let $\lambda : \text{SO}(X) \rightarrow \mathcal{P}(X)$ be an s -operation, a topological space (X, τ) is called $\lambda^D - R_0$, if $U \in \text{SO}_{\lambda^D}(X)$ and $x \in U$ then $\lambda^D \text{Cl}(\{x\}) \subseteq U$.

Example 2 Let $X = \{a, b, c, d\}$, and $\tau = \mathcal{P}(X)$. We define an s -operation $\lambda : \text{SO}(X) \rightarrow \mathcal{P}(X)$ as:

$\lambda(A) = A$, for every subset A of X .

$\text{SO}(X) = \mathcal{P}(X) = \text{SO}_{\lambda^D}(X) = \text{SC}_{\lambda^D}(X)$.

Theorem 2 For any topological space X and any s -operation $\lambda : SO(X) \rightarrow \mathcal{P}(X)$, the following statements are equivalent:

- (1) X is $\lambda^D - R_0$.
- (2) $F \in SC_{\lambda^D}(X)$ and $x \notin F$ implies that $F \subseteq U$ and $x \notin U$ for some $U \in SO_{\lambda^D}(X)$.
- (3) $F \in SC_{\lambda}(X)$ and $x \notin F$ implies that $F \cap \lambda^D Cl(\{x\}) \neq \emptyset$.
- (4) For any two distinct points x, y of X , either $\lambda^D Cl(\{x\}) = \lambda^D Cl(\{y\})$ or $\lambda^D Cl(\{x\}) \cap \lambda^D Cl(\{y\}) = \emptyset$.

Proof.

(1) \Rightarrow (2): Let $F \in SC_{\lambda^D}(X)$ and $x \notin F$. This implies that $x \in X \setminus F \in SO_{\lambda^D}(X)$, then $\lambda^D Cl(\{x\}) \subseteq X \setminus F$ (by (1)). Put $U = X \setminus \lambda^D Cl(\{x\})$. Then $x \notin U \in SO_{\lambda^D}(X)$ and $F \subseteq U$.

(2) \Rightarrow (3): $F \in SC_{\lambda^D}(X)$ and $x \notin F$ then there exists $U \in SO_{\lambda^D}(X)$ such that $x \notin U$ and $F \subseteq U$ (by (2)), then $U \cap \lambda^D Cl(\{x\}) = \emptyset$ and $F \cap \lambda^D Cl(\{x\}) = \emptyset$.

(3) \Rightarrow (4): Suppose that for any two distinct points x, y of X , if $\lambda^D Cl(\{x\}) \neq \lambda^D Cl(\{y\})$ Then, without loss of generality, we suppose that there exists some $z \in \lambda^D Cl(\{x\})$ such that $z \notin \lambda^D Cl(\{y\})$. Thus, there exists a λ^D -open set V such that $z \in V$ and $y \notin V$ but $x \in V$. Thus $x \notin \lambda^D Cl(\{y\})$. Hence by (3), $\lambda^D Cl(\{x\}) \cap \lambda^D Cl(\{y\}) = \emptyset$.

(4) \Rightarrow (1): Let $U \in SO_{\lambda^D}(X)$ and $x \in U$. Then for each $y \notin U$, $x \notin \lambda^D Cl(\{y\})$. Thus $\lambda^D Cl(\{x\}) \neq \lambda^D Cl(\{y\})$. Hence by (4), $\lambda^D Cl(\{x\}) \cap \lambda^D Cl(\{y\}) = \emptyset$, for each $y \in X \setminus U$. So $\lambda^D Cl(\{x\}) \cap [\cup\{\lambda^D Cl(\{y\}) : y \in X \setminus U\}] = \emptyset$. Now, $U \in SO_{\lambda^D}(X)$ and $y \in X \setminus U$ then $\{y\} \subseteq \lambda^D Cl(\{y\}) \subseteq \lambda^D Cl(X \setminus U) = X \setminus U$. Thus $X \setminus U = \cup\{\lambda^D Cl(\{y\}) : y \in X \setminus U\}$. Hence, $\lambda^D Cl(\{y\}) \cap X \setminus U = \emptyset$ then $\lambda^D Cl(\{x\}) \subseteq U$. This showing that (X, τ) is $\lambda^D - R_0$. \square

Lemma 2 Let $\lambda : SO(X) \rightarrow \mathcal{P}(X)$ be an s -operation. Then $y \in \lambda^D \text{Ker}(\{x\})$ if and only if $x \in \lambda^D Cl(\{y\})$.

Proof. Let $y \notin \lambda^D \text{Ker}(\{x\})$. Then there exists $V \in SO_{\lambda^D}(X)$ containing x such that $y \notin V$. Therefore $x \notin \lambda^D Cl(\{y\})$. The converse part can be proved in a similar way. \square

Theorem 3 Let $\lambda : SO(X) \rightarrow \mathcal{P}(X)$ be an s -operation. Then for any two points x, y in X , $\lambda^D \text{Ker}(\{x\}) \neq \lambda^D \text{Ker}(\{y\})$ if and only if $\lambda^D Cl(\{y\}) \neq \lambda^D Cl(\{x\})$.

Proof. Suppose that $\lambda^D \text{Ker}(\{x\}) \neq \lambda^D \text{Ker}(\{y\})$. Then there exists $z \in \lambda^D \text{Ker}(\{x\})$ such that $z \notin \lambda^D \text{Ker}(\{y\})$. Now, $z \in \lambda^D \text{Ker}(x)$ if and only if $x \in \lambda^D \text{Ker}(\{z\})$ by Lemma 2 and $z \notin \lambda^D \text{Ker}(\{y\})$ if and only if $y \in \lambda^D \text{Cl}(\{x\})$ by Lemma 2. Hence $\lambda^D \text{Cl}(\{x\}) \neq \lambda^D \text{Cl}(\{y\})$.

Conversely, suppose that $\lambda^D \text{Cl}(\{x\}) \neq \lambda^D \text{Cl}(\{y\})$. Then there exists $z \in X$ such that $z \in \lambda^D \text{Cl}(\{x\})$ and $z \notin \lambda^D \text{Cl}(\{y\})$ so there exists $U \in \text{SO}_{\lambda^D}(X)$ such that $z \in U$, $y \notin U$ and $x \in U$. Then $y \notin \lambda^D \text{Ker}(\{x\})$. Thus $\lambda^D \text{Ker}(\{x\}) \neq \lambda^D \text{Ker}(\{y\})$. \square

Theorem 4 *Let $\lambda : \text{SO}(X) \rightarrow \mathcal{P}(X)$ be an s -operation. Then (X, τ) is $\lambda^D - R_0$ if and only if for any two points $x, y \in X$, $\lambda^D \text{Ker}(\{x\}) \not\subseteq \lambda^D \text{Ker}(\{y\})$, implies that $\lambda^D \text{Ker}(\{x\}) \cap \lambda^D \text{Ker}(\{y\}) = \emptyset$.*

Proof. Let x, y be any two points in a $\lambda^D - R_0$ space X such that $\lambda^D \text{Ker}(\{x\}) \neq \lambda^D \text{Ker}(\{y\})$. Hence by Theorem 3, $\lambda^D \text{Cl}(\{x\}) \neq \lambda^D \text{Cl}(\{y\})$. We show that $\lambda^D \text{Ker}(\{x\}) \cap \lambda^D \text{Ker}(\{y\}) = \emptyset$. In fact, if $z \in \lambda^D \text{Ker}(\{x\}) \cap \lambda^D \text{Ker}(\{y\})$, then by Lemma 2, we have $x, y \in \lambda^D \text{Cl}(z)$ and by Theorem 2, we obtain that $\lambda^D \text{Cl}(\{x\}) = \lambda^D \text{Cl}(\{z\}) = \lambda^D \text{Cl}(\{y\})$ which is impossible.

Conversely, suppose that for any points $x, y \in X$, $\lambda^D \text{Ker}(\{x\}) \neq \lambda^D \text{Ker}(\{y\})$. Thus $\lambda^D \text{Ker}(\{x\}) \cap \lambda^D \text{Ker}(\{y\}) = \emptyset$. Hence we get $\lambda^D \text{Cl}(\{x\}) \cap \lambda^D \text{Cl}(\{y\}) = \emptyset$. In fact $z \in \lambda^D \text{Cl}(\{x\}) \cap \lambda^D \text{Cl}(\{y\})$, this implies that $x, y \in \lambda^D \text{Ker}(\{z\})$. Thus $\lambda^D \text{Cl}(\{x\}) \cap \lambda^D \text{Cl}(\{z\}) \neq \emptyset$. Hence by hypothesis, we get $\lambda^D \text{Ker}(\{x\}) = \lambda^D \text{Ker}(\{z\})$. By similar way it follows that $\lambda^D \text{Ker}(\{x\}) = \lambda^D \text{Ker}(\{z\})$. Thus $\lambda^D \text{Ker}(\{x\}) \neq \lambda^D \text{Ker}(\{y\})$ which is a contradiction. Hence $\lambda^D \text{Cl}(\{x\}) \cap \lambda^D \text{Cl}(\{y\}) \neq \emptyset$ and then by Theorem 2, the space X is $\lambda^D - R_0$. \square

Theorem 5 *Let (X, τ) be a topological space and for any s -operation $\lambda : \text{SO}(X) \rightarrow \mathcal{P}(X)$ the following statements are equivalent:*

- (1) X is a $\lambda^D - R_0$ space.
- (2) For any non-empty set A in X and any $G \in \text{SO}_{\lambda^D}(X)$ such that $A \cap G \neq \emptyset$ there exists $F \in \text{SC}_{\lambda^D}(X)$ such that $A \cap F \neq \emptyset$ and $F \subseteq G$.
- (3) For any $G \in \text{SO}_{\lambda^D}(X)$, $G = \cup\{F \in \text{SC}_{\lambda^D}(X) : F \subseteq G\}$.
- (4) For any $F \in \text{SC}_{\lambda^D}(X)$, $F = \cap\{G \in \text{SO}_{\lambda^D}(X) : F \subseteq G\}$.
- (5) For any $x \in X$, $\lambda^D \text{Cl}(\{x\}) \subseteq \lambda^D \text{Ker}(\{x\})$.

Proof.

(1) \Rightarrow (2): Let A be a non-empty subset of X and $G \in \text{SO}_{\lambda^D}(X)$ such that $A \cap G \neq \emptyset$. Let $x \in A \cap G$. Then as $x \in G \in \text{SO}_{\lambda^D}(X)$, by (1), we get $\lambda^D \text{Cl}(\{x\}) \subseteq G$. Put $F = \lambda^D \text{Cl}(\{x\})$. Then $F \in \text{SC}_{\lambda^D}(X)$, $F \subseteq G$ and $A \cap F \neq \emptyset$.

(2) \Rightarrow (3): Let $G \in \text{SO}_{\lambda^D}(X)$. Then $\bigcup \{F \in \text{SC}_{\lambda^D}(X) : F \subseteq G\} \subseteq G$. Let $x \in G$. Then there exists $F \in \text{SC}_{\lambda^D}(X)$ such that $x \in F$ and $F \subseteq G$. Thus $x \in F \cup \{K \in \text{SC}_{\lambda^D}(X) : K \subseteq G\}$. Hence (3) follows.

(3) \Rightarrow (4): Straight forward.

(4) \Rightarrow (5): Let $x \in X$. Now, $y \notin \lambda^D \text{Ker}(\{x\})$ implies there exists $V \in \text{SO}_{\lambda^D}(X)$ such that $x \in V$ and $y \notin V$ then $\lambda^D \text{Cl}(\{y\}) \cap V = \emptyset$. This implies by (4) $[\bigcap \{G \in \text{SO}_{\lambda^D}(X) : \lambda^D \text{Cl}(\{y\}) \subseteq G\}] \cap V = \emptyset$. Then there exists $G \in \text{SO}_{\lambda^D}(X)$ such that $x \in G$ and $\lambda^D \text{Cl}(\{y\}) \subseteq G$, so $y \notin \lambda^D \text{Cl}(\{x\})$.

(5) \Rightarrow (1): Let $G \in \text{SO}_{\lambda^D}(X)$ and $x \in G$. Let $y \in \lambda^D \text{Ker}(\{x\})$. Then $x \in \lambda^D \text{Cl}(\{y\})$ and hence $y \in G$. This implies that $\lambda^D \text{Ker}(\{x\}) \subseteq G$. Thus $x \in \lambda^D \text{Cl}(\{x\}) \subseteq \lambda^D \text{Ker}(\{x\}) \subseteq G$. Hence X is $\lambda^D - R_0$. \square

Corollary 1 *Let $\lambda : \text{SO}(X) \rightarrow \mathcal{P}(X)$ be an s -operation. Then X is $\lambda^D - R_0$ if and only if $\lambda^D \text{Cl}(\{x\}) = \lambda^D \text{Ker}(\{x\})$, for all $x \in X$.*

Proof. Suppose that X is $\lambda^D - R_0$. By Theorem 5, $\lambda^D \text{Cl}(\{x\}) \subseteq \lambda^D \text{Ker}(\{x\})$. For each $x \in X$. Let $y \in \lambda^D \text{Ker}(\{x\})$. Then $x \in \lambda^D \text{Cl}(\{y\})$ (by Lemma 2), and hence by Theorem 2, $\lambda^D \text{Cl}(\{x\}) = \lambda^D \text{Cl}(\{y\})$. Thus $y \in \lambda^D \text{Cl}(\{x\})$ and hence $\lambda^D \text{Ker}(\{x\}) \subseteq \lambda^D \text{Cl}(\{x\})$. Thus $\lambda^D \text{Cl}(\{x\}) = \lambda^D \text{Ker}(\{x\})$. \square

The converse is obvious in view of Theorem 5.

Theorem 6 *Let (X, τ) be a topological space and $\lambda : \text{SO}(X) \rightarrow \mathcal{P}(X)$ be an s -operation. A space X is $\lambda^D - R_0$ if and only if for any $x, y \in X$, whenever $x \in \lambda^D \text{Cl}(\{y\})$ implies $y \in \lambda^D \text{Cl}(\{x\})$ and conversely.*

Proof. Suppose that a topological space (X, τ) is $\lambda^D - R_0$. Let $x \in \lambda^D \text{Cl}(\{y\})$. Then by Theorem 5, we have $\lambda^D \text{Cl}(\{y\}) \subseteq \lambda^D \text{Ker}(\{x\})$. Thus $x \in \lambda^D \text{Ker}(\{y\})$. Hence by Lemma 1, we have $y \in \lambda^D \text{Cl}(\{x\})$.

Conversely, let $U \in \text{SO}_{\lambda^D}(X)$ and $x \in U$. Let $y \in \lambda^D \text{Cl}(\{x\})$ hence by hypothesis, $x \in \lambda^D \text{Cl}(\{y\})$. Since $x \in U$, so $y \in U$. Hence $\lambda^D \text{Cl}(\{x\}) \subseteq U$. Thus X is $\lambda^D - R_0$. \square

Theorem 7 *Let X be a topological space and $\lambda : \text{SO}(X) \rightarrow \mathcal{P}(X)$ be an s -operation. Then the following statements are equivalent:*

- (1) X is $\lambda^D - R_0$.

- (2) If $F \in SC_{\lambda^D}(X)$ then $F = \lambda^D \text{Ker}(F)$.
- (3) If $F \in SC_{\lambda^D}(X)$ and $x \in F$, then $\lambda^D \text{Ker}(\{x\}) \subseteq F$.
- (4) If $x \in X$, then $\lambda^D \text{Ker}(\{x\}) \subseteq \lambda^D \text{Cl}(\{x\})$.

Proof.

(1) \Rightarrow (2): Follows from Theorem 5.

(2) \Rightarrow (3): Follows from the fact that $x \in F$ then $\lambda^D \text{Ker}(\{x\}) \subseteq \lambda^D \text{Ker}(F) = F$ by part 3 of Theorem 1.

(3) \Rightarrow (4): Since $x \in \lambda^D \text{Cl}(\{x\}) \in SC_{\lambda^D}(X)$ we have by (3), $\lambda^D \text{Ker}(\{x\}) \subseteq \lambda^D \text{Cl}(\{x\})$ and (4) follows.

(4) \Rightarrow (1): Let $U \in SO_{\lambda^D}(X)$ and $x \in U$. To show $\lambda^D \text{Cl}(\{x\}) \subseteq U$. If possible, suppose that, there exists $y \in \lambda^D \text{Cl}(\{x\})$ such that $y \notin U$. Then $y \in X \setminus U$. This by (4) implies that $\lambda^D \text{Ker}(\{y\}) \subseteq X \setminus U$. Therefore $U \subseteq X \setminus \lambda^D \text{Ker}(\{x\})$. So $x \notin \lambda^D \text{Ker}(\{y\})$. Then, there exists a λ^D -open set G such that $y \in G$ but $x \notin G$. This implies that $y \notin \lambda^D \text{Cl}(\{x\})$ which is impossible. Hence $\lambda^D \text{Cl}(\{x\}) \subseteq U$. Thus X is a $\lambda^D - R_0$ space. \square

Definition 6 Let (X, τ) be a topological space and $\lambda : SO(X) \rightarrow \mathcal{P}(X)$ be an s -operation. The space X is said to be $\lambda^D - R_1$ if for $x, y \in X$ with $\lambda^D \text{Cl}(\{x\}) \neq \lambda^D \text{Cl}(\{y\})$ there exist disjoint λ^D -open sets U and V such that $\lambda^D \text{Cl}(\{x\}) \subseteq U$ and $\lambda^D \text{Cl}(\{y\}) \subseteq V$.

Remark 1 A space X in Example 2 is $\lambda^D - R_1$.

Theorem 8 Every $\lambda^D - R_1$ space is a $\lambda^D - R_0$ space.

Proof. Let $U \in SO_{\lambda^D}(X)$ and $x \in U$. If $y \notin U$ then $\lambda^D \text{Cl}(\{x\}) \neq \lambda^D \text{Cl}(\{y\})$ (as $x \notin \lambda^D \text{Cl}(\{y\})$). Hence there exists $V \in SO_{\lambda^D}(X)$ such that $\lambda^D \text{Cl}(\{y\}) \subseteq V$ and $x \notin V$. This gives $y \notin \lambda^D \text{Cl}(\{y\})$, proving that $\lambda^D \text{Cl}(\{x\}) \subseteq U$. So X is a $\lambda^D - R_0$ space. \square

The converse of Theorem 8 is not true, we can show it by the following example:

Example 3 Let $X = \{a, b, c, d\}$, and $\tau = \mathcal{P}(X)$. We define an s -operation $\lambda : SO(X) \rightarrow \mathcal{P}(X)$ as:

$$\lambda(A) = \begin{cases} X & \text{Otherwise} \\ A & \text{if } A = \emptyset \text{ or } \{b, c\} \text{ or } \{a, c\} \text{ or } \{a, b\}. \end{cases}$$

Now:

$$SO(X) = \mathcal{P}(X).$$

$$SO_{\lambda^D}(X) = \{\emptyset, \{b, c\}, \{a, c\}, \{a, b\}, X\}.$$

$$SC_{\lambda^D}(X) = \{\emptyset, \{a\}, \{b\}, \{c\}, X\}.$$

Clearly X is $\lambda^D - R_0$ but it is not $\lambda^D - R_1$.

Theorem 9 Let (X, τ) be a topological space and $\lambda : SO(X) \rightarrow \mathcal{P}(X)$ be an s -operation. Then the following statements are equivalent:

- (1) X is $\lambda^D - R_1$.
- (2) For any $x, y \in X$, one of the following holds:
 - a) For $U \in SO_\lambda(X)$, $x \in U$ if and only if $y \in U$;
 - b) There exist disjoint λ^D -open sets U and V such that $x \in U$, $y \in V$.
- (3) If $x, y \in X$, such that $\lambda^D Cl(\{x\}) \neq \lambda^D Cl(\{y\})$ then there exist λ^D -closed sets F and H such that $x \in F$, $y \notin F$, $y \in H$, $x \notin H$ and $X = F \cup H$.

Proof.

(1) \Rightarrow (2): Let $x, y \in X$. Then $\lambda^D Cl(\{x\}) = \lambda^D Cl(\{y\})$ or $\lambda^D Cl(\{x\}) \neq \lambda^D Cl(\{y\})$. If $\lambda^D Cl(\{x\}) = \lambda^D Cl(\{y\})$ and $U \in SO_{\lambda^D}(X)$, then for any $U \in SO_{\lambda^D}(X)$, $x \in U$ then $y \in \lambda^D Cl(\{x\}) = \lambda^D Cl(\{y\}) \subseteq U$ then (as X is $\lambda^D - R_0$). If $\lambda^D Cl(\{x\}) \neq \lambda^D Cl(\{y\})$, then there exist $U, V \in SO_{\lambda^D}(X)$ such that $x \in \lambda^D Cl(\{x\}) \subseteq U$, $y \in \lambda^D Cl(\{y\}) \subseteq V$ and $U \cap V = \emptyset$.

(2) \Rightarrow (3): Let $x, y \in X$ such that $\lambda^D Cl(\{x\}) \neq \lambda^D Cl(\{y\})$. Then $x \notin \lambda^D Cl(\{y\})$, so that there exists $G \in SO_{\lambda^D}(X)$, such that $x \in G$ and $y \notin G$. Thus by (2), there exist disjoint λ^D -open sets U and V such that $x \in U$, $y \in V$. Put $F = X \setminus V$ and $H = X \setminus U$. Then $F, H \in SO_{\lambda^D}(X)$, $x \in F$, $y \notin F$, $y \in H$, $x \notin H$ and $X = F \cup H$.

(3) \Rightarrow (1): Let $U \in SO_\lambda(X)$ and $x \in U$. Then $\lambda^D Cl(\{x\}) \subseteq U$. In fact, otherwise there exists $y \in \lambda^D Cl(\{x\}) \cap X \setminus U$. Implies that $\lambda^D Cl(\{x\}) \neq \lambda^D Cl(\{y\})$ (as $x \notin \lambda^D Cl(\{y\})$) and so by (3), there exist $F, H \in SO_{\lambda^D}(X)$ such that $x \in F$, $y \notin F$, $y \in H$, $x \notin H$ and $X = F \cup H$. Then $y \in H \setminus F = X \setminus F$ and $x \notin X \setminus F$, where $X \setminus F \in SO_{\lambda^D}(X)$, which is a contradiction to the fact that $y \in \lambda^D Cl(\{x\})$. Hence $\lambda^D Cl(\{x\}) \subseteq U$. Thus X is $\lambda^D - R_0$. To show X to be $\lambda^D - R_1$. Assume that $a, b \in X$ with $\lambda^D Cl(\{a\}) \neq \lambda^D Cl(\{b\})$. Then as above, there exist $K, L \in SC_{\lambda^D}(X)$ such that $a \in K$, $b \notin K$, $b \in L$, $a \notin L$ and $X = K \cup L$. Thus $a \in K \setminus L \in SO_{\lambda^D}(X)$, $b \in L \setminus K \in SO_{\lambda^D}(X)$. So $\lambda^D Cl(\{a\}) \subseteq K \setminus L$, $\lambda^D Cl(\{b\}) \subseteq L \setminus K$. Thus X is $\lambda^D - R_1$. \square

Proposition 2 Let (X, τ) be a topological space and $\lambda : SO(X) \rightarrow \mathcal{P}(X)$ be an s-operation. Then X is $\lambda^D - R_1$, if and only if for $x, y \in X$, with $\lambda^D \text{Ker}(\{x\}) \neq \lambda^D \text{Ker}(\{y\})$ there exist disjoint λ^D -open sets U and V such that $\lambda^D \text{Cl}(\{x\}) \subseteq U$ and $\lambda^D \text{Cl}(\{y\}) \subseteq V$.

Proof. Follows from Theorem 3 and Definition 6. □

4 Conclusion

Introduced by Alais B. Khalaf and Sarhad F. Namiq [1]. The main results are the following:

- (1) Let (X, τ) be a topological space, and $\lambda : SO(X) \rightarrow \mathcal{P}(X)$ be an s-operation and $A \subseteq X$. Then $\lambda^D \text{Ker}(\{A\}) = \{x \in X : \lambda^D \text{Cl}(\{x\}) \cap A \neq \emptyset\}$.
- (2) For any topological space X and any s-operation $\lambda : SO(X) \rightarrow \mathcal{P}(X)$, the following statements are equivalent:
 - a) X is $\lambda^D - R_0$.
 - b) $F \in SC_{\lambda^D}(X)$ and $x \in F$ implies that $F \subseteq U$ and $x \in U$ for some $U \in SO_{\lambda^D}(X)$.
 - c) $F \in SC_{\lambda^D}(X)$ and $x \notin F$ implies that $\lambda^D \text{Cl}(\{x\}) \cap \lambda^D \text{Cl}(\{y\}) = \emptyset$.
 - d) For any two distinct points x, y of X , either $\lambda^D \text{Cl}(\{x\}) = \lambda^D \text{Cl}(\{y\})$ or $\lambda^D \text{Cl}(\{x\}) \cap \lambda^D \text{Cl}(\{y\}) = \emptyset$.
- (3) Every $\lambda^D - R_1$ space is a $\lambda^D - R_0$ space.

References

- [1] Alias B. Khalaf and Sarhad F. Namiq, *New types of continuity and separation axiom based operation in topological spaces*, M. Sc. Thesis, University of Sulaimani (2011).
- [2] Alias B. Khalaf, Sarhad F. Namiq, Generalized λ -Closed Sets and $(\lambda, \gamma)^D$ -Continuous Functions, *International Journal of Scientific & Engineering Research*, Volume 3, Issue 12, December-2012 1 ISSN 2229-5518.
- [3] A. S. Davis, Indexed systems of neighborhoods for general topological spaces, *Amer. Math. Monthly*, **68** (1961), 886-893.

- [4] N. Levine, Semi-open sets and semi-continuity in topological spaces, *Amer. Math. Monthly*, **70** (1) (1963), 36–41.
- [5] Sarhad F. Namiq, Some Properties of λ^D –Open Sets in Topological Spaces (submit).
- [6] N. A. Shanin, On separation in topological spaces, *Dokl. Akad. Nauk. SSSR*,. **38** (1943), 110–113.
- [7] J. N. Sharma and J. P. Chauhan, *Topology (General and Algebraic)*, Krishna Prakashna Media, India. (2011).

Received: January 23, 2020



Soft covered ideals in semigroups

Şerif Özlü

Department of Mathematics
Gaziantep University
27300 Gaziantep, Turkey
email: serif.ozlu@hotmail.com

Ashhan Sezgin

Department of Mathematics
and Science Education
Amasya University
05100 Amasya, Turkey
email: aslihan.sezgin@amasya.edu.tr

Abstract. In this work, soft covered ideals in semigroups are constructed and in this concept, soft covered semigroups, soft covered left (right) ideals, soft covered interior ideals, soft covered (generalized) bi-ideals and soft covered quasi ideals of a semigroup are defined. Various properties of these ideals are introduced and the interrelations of these soft covered ideals and the relations of soft anti covered ideals and soft covered ideals are investigated.

1 Introduction

Soft set theory introduced by Molodstov in 1999 [1] is applied uncertainty complicated problems especially in economy, medical, engineering and over classical Mathematics such as groups [2], semirings [3], rings [4], BCK/BCI-algebras [5, 6, 7], d-algebras [8], BL-algebras [9], BCH-algebras [10] and near-rings [11].

The works over soft set theory progressed rapidly and after then, this theory started to extend over fuzzy set, rough set and so on and a lot of mathematical structures were constructed over these sets. Moreover, soft set theory has

2010 Mathematics Subject Classification: Primary 03EE99; Secondary 03E75

Key words and phrases: soft set, soft covered left(right) ideal, soft covered ideal, soft covered bi-ideal, soft covered interior ideal, soft covered quasi-ideal

received abroad attention by several authors in game theory, intelligent system and especially decision making method. After decision making method was applied over uncertainty sets, it could solve complicated problems including medical diagnoses, finding the best alternative solution for a company for example; the best worker, the best job etc., logistic industry and so on. A relation has been developed between soft set and decision making method and it has overcome several problems [12, 13, 14, 15].

A lot of ideals were introduced over fuzzy sets. Mandal [16] introduced fuzzy ideals and definition of fuzzy interior ideals of ordered semirings. Dib and Galhum [17] defined fuzzy groupoid, a fuzzy semigroup and also fuzzy ideals and fuzzy bi-ideals of a semigroup. Moreover, Kazanci and Yamak [18] gave a kind of generalized fuzzy bi-ideals of semigroups. Kavikumar and Khamis [19] studied fuzzy ideals and fuzzy quasi ideals in ternary semirings. Then, Changphas and Summagrap [20] worked semigroups in covered ideals. Sezer and others wrote several papers [21, 22] over soft ideals in semigroups. In one of papers, they introduced soft union semigroup, ideals and bi-ideals [21]. After then, they defined soft intersection quasi ideals, generalized bi-ideals of a semigroup and surveyed regular, weakly regular, intra regular, completely regular, quasi regular semigroups with help of these ideals [22].

Fabrici [23, 24] investigated covered left, right ideal of a semigroup. By inspiring covered ideals, anti-covered (AC)-left (right), interior ideals were defined in [25] and Xie and Yan [25] obtained fuzzy anti-covered (AC)-left (right), bi-ideal, interior ideal of a semigroup and studied their basic properties. In [26], soft anti covered (AC)-ideals of semigroups were defined and studied in detailed.

In this manuscript, we first give the basic definitions of soft sets and covered ideals in semigroups. By using these basic definitions, we introduce covered ideals of semigroups by defining soft covered semigroups, soft covered left (right) ideals, soft covered bi-ideals, soft covered interior ideals, soft covered quasi-ideals, soft covered generalized bi-ideals. We discuss their properties and the interrelations with each others. Also, we study the relationship between soft anti covered ideals defined in [26] and soft covered ideals which is the most important point in this article.

2 Methodology

From now on, U refers to an initial universe, E is a set of parameters, $P(U)$ is the power set of U and $A, B, C \subseteq E$.

Definition 1 [1, 13] A soft set f_A over U is a set defined by

$$f_A : E \rightarrow P(U) \text{ such that } f_A(x) = \emptyset \text{ if } x \notin A.$$

Here f_A is also called an approximate function. A soft set over U can be represented by the set of ordered pairs

$$f_A = \{(x, f_A(x)) : x \in E, f_A(x) \in P(U)\}.$$

It is clear to see that a soft set is a parameterized family of subsets of the set U . Note that the set of all soft sets over U will be denoted by $S(U)$.

Definition 2 [13] Let $f_A, f_B \in S(U)$. Then, f_A is called a soft subset of f_B and denoted by $f_A \subseteq f_B$, if $f_A(x) \subseteq f_B(x)$ for all $x \in E$.

Definition 3 [13] Let $f_A, f_B \in S(U)$. Then, union of f_A and f_B , denoted by $f_A \widetilde{\cup} f_B$, is defined as $f_A \widetilde{\cup} f_B = f_{A \widetilde{\cup} B}$, where $f_{A \widetilde{\cup} B}(x) = f_A(x) \cup f_B(x)$, intersection of f_A and f_B , denoted by $f_A \widetilde{\cap} f_B$, is defined as $f_A \widetilde{\cap} f_B = f_{A \widetilde{\cap} B}$, where $f_{A \widetilde{\cap} B}(x) = f_A(x) \cap f_B(x)$ for all $x \in E$.

Definition 4 [13] Let $f_A, f_B \in S(U)$. Then, $\widetilde{\wedge}$ -product of f_A and f_B , denoted by $f_A \widetilde{\wedge} f_B$, is defined as $f_A \widetilde{\wedge} f_B = f_{A \widetilde{\wedge} B}$, where $f_{A \widetilde{\wedge} B}(x, y) = f_A(x) \cap f_B(y)$ for all $(x, y) \in E \times E$.

Definition 5 [27] Let f_A and f_B be soft sets over the common universe U and Ψ be a function from A to B . Then, soft image of f_A under Ψ , denoted by $\Psi(f_A)$, is a soft set over U by

$$(\Psi(f_A))(b) = \begin{cases} \bigcap \{f_A(a) \mid a \in A \text{ and } \Psi(a) = b\}, & \text{if } \Psi^{-1}(b) \neq \emptyset, \\ \emptyset, & \text{otherwise} \end{cases}$$

for all $b \in B$. And soft pre-image (or soft inverse image) of f_B under Ψ , denoted by $\Psi^{-1}(f_B)$, is a soft set over U by $(\Psi^{-1}(f_B))(a) = f_B(\Psi(a))$ for all $a \in A$.

From now on, S denotes a semigroup.

Definition 6 [21] Let f_S and g_S be soft sets over the common universe U . Then, soft union product $f_S * g_S$ is defined by

$$(f_S * g_S)(x) = \begin{cases} \bigcap_{x=yz} \{f_S(y) \cup g_S(z)\}, & \text{if } \exists y, z \in S \text{ such that } x = yz, \\ \emptyset, & \text{otherwise} \end{cases}$$

for all $x \in S$.

Theorem 1 [21] *Let $f_S, g_S, h_S \in S(\mathcal{U})$. Then,*

- i) $(f_S * g_S) * h_S = f_S * (g_S * h_S)$.
- ii) $f_S * g_S \neq g_S * f_S$, generally.
- iii) $f_S * (g_S \widetilde{\cup} h_S) = (f_S * g_S) \widetilde{\cup} (f_S * h_S)$ and $(f_S \widetilde{\cup} g_S) * h_S = (f_S * h_S) \widetilde{\cup} (g_S * h_S)$.
- iv) $f_S * (g_S \widetilde{\cap} h_S) = (f_S * g_S) \widetilde{\cap} (f_S * h_S)$ and $(f_S \widetilde{\cap} g_S) * h_S = (f_S * h_S) \widetilde{\cap} (g_S * h_S)$.
- v) If $f_S \subseteq g_S$, then $f_S * h_S \subseteq g_S * h_S$ and $h_S * f_S \subseteq h_S * g_S$.
- vi) If $t_S, l_S \in S(\mathcal{U})$ such that $t_S \subseteq f_S$ and $l_S \subseteq g_S$, then $t_S * l_S \subseteq f_S * g_S$.

From now on, if $f_S : S \rightarrow P(\mathcal{U})$ is a soft set satisfying $f_S(x) = \emptyset$ for all $x \in S$, then f_S is denoted by θ and if $f_S : S \rightarrow P(\mathcal{U})$ is a soft set satisfying $f_S(x) = \mathcal{U}$ for all $x \in S$, then f_S is denoted by \mathbb{S} .

Lemma 1 *Let f_S be any soft set over \mathcal{U} . Then, we have the followings.*

- i) $\theta * \theta \supseteq \theta$.
- ii) $(\mathbb{S} - f_S) * \theta \supseteq \theta$ and $\theta * (\mathbb{S} - f_S) \supseteq \theta$.
- iii) $(\mathbb{S} - f_S) \widetilde{\cup} \theta = (\mathbb{S} - f_S)$ and $(\mathbb{S} - f_S) \widetilde{\cap} \theta = \theta$.

Soft anti covered (AC)-ideals of a semigroup were defined in [26] as following:

Definition 7 [26] f_S is called as soft AC-semigroup over \mathcal{U} , if

$$f_S(xy) \supseteq (\mathbb{S} - f_S)(x) \cap (\mathbb{S} - f_S)(y)$$

for all $x, y \in S$.

Definition 8 [26] f_S is called a soft AC-left ideal over \mathcal{U} , if $f_S(xy) \supseteq (\mathbb{S} - f_S)(y)$, AC-right ideal of S over \mathcal{U} , if $f_S(xy) \supseteq (\mathbb{S} - f_S)(x)$, AC-ideal of S over \mathcal{U} , if $f_S(xy) \supseteq (\mathbb{S} - f_S)(y)$ and $f_S(xy) \supseteq (\mathbb{S} - f_S)(x)$ for all $x, y \in S$.

Definition 9 [26] A soft AC-semigroup f_S over \mathcal{U} is called a soft AC-bi-ideal over \mathcal{U} , if

$$f_S(xyz) \supseteq (\mathbb{S} - f_S)(x) \cap (\mathbb{S} - f_S)(z)$$

Definition 10 [26] f_S is called a soft AC-interior ideal over \mathcal{U} , if

$$f_S(xyz) \supseteq (\mathbb{S} - f_S)(y)$$

and is called soft AC-generalized bi-ideal of S , if

$$f_S(xyz) \supseteq (\mathbb{S} - f_S)(x) \cap (\mathbb{S} - f_S)(y)$$

for all $x, y, z \in S$.

From now on, all covered ideals are denoted by C-ideals for the sake of brevity. C-ideals of a semigroup (and fuzzy C-ideals of a semigroup) are defined in [23, 24] as following:

L is called C-left ideal of S , if $L \subseteq S(S - L)$ and R is called C-right ideal of S , if $R \subseteq (S - R)S$. By *C-two-sided ideal*, it is meant a subset of S , which is both C-left and C-right ideal of S . X is called a *C-interior* of S if $X \subseteq (S - X)S(S - X)$. X is called C-bi-ideal of S if $X \subseteq S(S - X)S$ and X is called C-quasi-ideal of S if $X \subseteq (S - X)S \cup S(S - X)$.

3 Results and discussion

3.1 Soft C-semigroup

In this section, we construct soft covered semigroup and study some properties of it.

Definition 11 Let S be semigroup and f_S be soft set over \mathcal{U} . f_S is called a soft covered semigroup, if

$$f_S(xy) \subseteq (\mathbb{S} - f_S)(x) \cup (\mathbb{S} - f_S)(y)$$

for all $x, y \in S$.

From now on, soft covered semigroup is denoted by soft C-semigroup for the sake of brevity.

Example 1 Consider the semigroup $S = \{a, b, c, d\}$ constructed by the following table:

.	a	b	c	d
a	a	a	a	a
b	a	a	a	a
c	a	a	b	a
d	a	a	b	b

Let $\mathbf{U} = \mathbf{D}_3 = \{< x, y >: x^3 = y^2 = e, xy = yx^2\} = \{e, x, x^2, y, yx, yx^2\}$ be the universal set and f_S be a soft set over \mathbf{U} such that $f_S(a) = \emptyset$, $f_S(b) = \{y\}$, $f_S(c) = \{e, yx^2\}$, $f_S(d) = \{e, x^2, yx, yx^2\}$ and so $(S - f_S)(a) = \{e, x, x^2, y, yx, yx^2\}$, $(S - f_S)(b) = \{e, x, x^2, yx, yx^2\}$, $(S - f_S)(c) = \{x, x^2, y, yx\}$, $(S - f_S)(d) = \{x, y\}$. One can show that f_S is a soft C -semigroup. However, if $f_S(b) = \{x, x^2, yx\}$, we can easily show that f_S is not a soft C -semigroup.

It is easy to see that if $f_S(x) = \emptyset$ for $x \in S$, then f_S is a soft C -semigroup over \mathbf{U} . We denote such kind of C -semigroup by θ .

Definition 12 [21] Let X be a subset of S . We denote by \mathcal{S}_X^c the soft anti characteristic function of X and define as

$$\mathcal{S}_X^c(x) = \begin{cases} \emptyset, & x \in X \\ \mathbf{U}, & x \in S - X \end{cases}$$

It is clear that soft anti characteristic function is a soft set over \mathbf{U} clearly,

$$\mathcal{S}_X^c : S \rightarrow P(\mathbf{U}).$$

Theorem 2 If X is a C -semigroup of S , then \mathcal{S}_X^c is a soft C -semigroup of S .

Proof. Let X be a C -semigroup and $x = pq \in X$. Since $X \subseteq (S - X)(S - X)$, then it follows that $x = pq \in (S - X)(S - X)$, and so $p, q \in (S - X)$. In this statement, $\mathcal{S}_X^c(pq) \subseteq (S - \mathcal{S}_X^c)(p) \cup (S - \mathcal{S}_X^c)(q)$, that is, \mathcal{S}_X^c is a soft C -semigroup of S . In fact,

$$\begin{aligned} \emptyset &= \mathcal{S}_X^c(pq) \\ &\subseteq (S - \mathcal{S}_X^c)(p) \cup (S - \mathcal{S}_X^c)(q) \\ &= (\mathbf{U} - \mathbf{U}) \cup (\mathbf{U} - \mathbf{U}) \\ &= \emptyset. \end{aligned}$$

□

Theorem 3 Let f_S be a soft set over \mathbf{U} . Then, f_S is a soft AC -semigroup over \mathbf{U} of S if and only if f_S^c is a soft C -semigroup over \mathbf{U} of S .

Proof. Let f_S be a soft AC -semigroup over \mathbf{U} of S . In this statement,

$$\begin{aligned} f_S^c(xy) &= (\mathbf{U} - f_S)(xy) \\ &\subseteq \mathbf{U} - ((S - f_S)(x) \cap (S - f_S)(y)) \\ &= (\mathbf{U} - (S - f_S)(x)) \cup (\mathbf{U} - (S - f_S)(y)) \\ &= (\mathbf{U} - (\mathbf{U} - f_S(x)) \cup (\mathbf{U} - (\mathbf{U} - f_S(y)) \\ &= (\mathbf{U} - f_S^c(x)) \cup (\mathbf{U} - f_S^c(y)) \\ &= (S - f_S^c)(x) \cup (S - f_S^c)(y) \end{aligned}$$

for all $x, y \in S$. Conversely, let f_S^c be a soft C-semigroup over \mathcal{U} of S . Then,

$$\begin{aligned}
 f_S(xy) &= (\mathcal{U} - f_S^c)(xy) \\
 &\supseteq \mathcal{U} - ((S - f_S^c)(x) \cup (S - f_S^c)(y)) \\
 &\supseteq (\mathcal{U} - (S - f_S^c)(x)) \cap (\mathcal{U} - (S - f_S^c)(y)) \\
 &= (\mathcal{U} - (\mathcal{U} - f_S^c(x))) \cap (\mathcal{U} - (\mathcal{U} - f_S^c(y))) \\
 &= (\mathcal{U} - f_S(x)) \cap (\mathcal{U} - f_S(y)) \\
 &= (S - f_S)(x) \cap (S - f_S)(y)
 \end{aligned}$$

for all $x, y \in S$. This completes the proof. \square

Theorem 4 Let f_S be a soft set over \mathcal{U} . Then, f_S is a soft C-semigroup over \mathcal{U} if and only if we have:

$$f_S \widetilde{\subseteq} (S - f_S) * (S - f_S).$$

Proof. To prove this, we assume that f_S is a soft C-semigroup over \mathcal{U} . If $f_S = \emptyset$, then it is trivial since

$$f_S(x) \subseteq ((S - f_S) * (S - f_S))(x)$$

thus, $f_S \subseteq (S - f_S) * (S - f_S)$. Otherwise, there exist elements $m, n \in S$ such that $x = mn$. Then, since f_S is a soft C-semigroup over \mathcal{U} , we have:

$$\begin{aligned}
 ((S - f_S) * (S - f_S))(x) &= \bigcap_{x=mn} (S - f_S)(m) \cup (S - f_S)(n) \\
 &\supseteq \bigcap_{x=mn} f_S(mn) \\
 &= f_S(x)
 \end{aligned}$$

thus, $f_S \widetilde{\subseteq} (S - f_S) * (S - f_S)$.

Conversely, suppose that $f_S \widetilde{\subseteq} (S - f_S) * (S - f_S)$. Let $m, n \in S$ and $x = mn$. Hence, we have:

$$\begin{aligned}
 f_S(mn) &= f_S(x) \\
 &\subseteq ((S - f_S) * (S - f_S))(x) \\
 &= \bigcap_{x=mn} (S - f_S)(m) \cup (S - f_S)(n) \\
 &\subseteq (S - f_S)(m) \cup (S - f_S)(n).
 \end{aligned}$$

Then, this means that f_S is a soft C-semigroup over \mathcal{U} . This completes the proof. \square

Proposition 1 Let f_S and f_T be soft C-semigroups over \mathcal{U} . Then, $f_S \widetilde{\wedge} f_T$ is a soft C-semigroup over \mathcal{U} .

Proof. Let $(x_1, y_1), (x_2, y_2)$ be any two elements of $S \times T$. Then,

$$\begin{aligned}
 f_{S \tilde{\wedge} T}((x_1, y_1), (x_2, y_2)) &= f_{S \wedge T}(x_1 x_2, y_1 y_2) \\
 &= f_S(x_1 x_2) \cap f_T(y_1 y_2) \\
 &\subseteq [(\mathbb{S} - f_S)(x_1) \cup (\mathbb{S} - f_S)(x_2)] \cap [(\mathbb{S} - f_T)(y_1) \\
 &\quad \cup (\mathbb{S} - f_T)(y_2)] \\
 &= [\mathbb{U} - (f_S(x_1) \cap f_S(x_2))] \cap [\mathbb{U} - (f_T(y_1) \cap f_T(y_2))] \\
 &= \mathbb{U} - [(f_S(x_1) \cap f_S(x_2)) \cup (f_T(y_1) \cap f_T(y_2))] \\
 &\subseteq \mathbb{U} - [(f_S(x_1) \cap f_S(x_2)) \cap (f_T(y_1) \cap f_T(y_2))] \\
 &= \mathbb{U} - [(f_S(x_1) \cap f_T(y_1)) \cap (f_S(x_2) \cap f_T(y_2))] \\
 &= (\mathbb{S} - f_{S \tilde{\wedge} T})(x_1, y_1) \cup (\mathbb{S} - f_{S \tilde{\wedge} T})(x_2, y_2).
 \end{aligned}$$

This implies that $f_S \tilde{\wedge} f_T$ is a soft C-semigroup over \mathbb{U} . □

Theorem 5 Let f_S and g_S be two soft C-semigroups over \mathbb{U} , then $f_S \tilde{\cap} g_S$ is also so.

Proof. Let f_S and g_S be two soft C-semigroups over \mathbb{U} for $x, y \in S$. Then,

$$\begin{aligned}
 (f_S \tilde{\cap} g_S)(xy) &= f_S(xy) \cap g_S(xy) \\
 &\subseteq [(\mathbb{S} - f_S)(x) \cup (\mathbb{S} - f_S)(y)] \cap [(\mathbb{S} - g_S)(x) \cup (\mathbb{S} - g_S)(y)] \\
 &= [\mathbb{U} - (f_S(x) \cap f_S(y))] \cap [\mathbb{U} - (g_S(x) \cap g_S(y))] \\
 &= \mathbb{U} - [(f_S(x) \cap f_S(y)) \cup (g_S(x) \cap g_S(y))] \\
 &\subseteq \mathbb{U} - [(f_S(x) \cap f_S(y)) \cap (g_S(x) \cap g_S(y))] \\
 &= \mathbb{U} - [(f_S(x) \cap g_S(x)) \cap ((f_S(y) \cap g_S(y)))] \\
 &= \mathbb{U} - [(f_S \tilde{\cap} g_S)(x) \cap (f_S \tilde{\cap} g_S)(y)] \\
 &= (\mathbb{S} - (f_S \tilde{\cap} g_S))(x) \cup (\mathbb{S} - (f_S \tilde{\cap} g_S))(y).
 \end{aligned}$$

This completes the proof. □

The union of two C-semigroups needs not to be a soft C-semigroup as shown in following example.

Example 2 Let f_S and g_S be two soft sets over $\mathbb{U} = D_3$ of semigroup $S = \{a, b, c, d\}$. We accept that f_S is the same soft set in Example 1 and define the soft set, g_S as following; $g_S(a) = \emptyset$, $g_S(b) = \{e, x\}$, $g_S(c) = \{x^2, yx, yx^2\}$ and $g_S(d) = \{y, yx, yx^2\}$. Since $(f_S \tilde{\cup} g_S)(dd) = (f_S \tilde{\cup} g_S)(b) \not\subseteq (\mathbb{S} - (f_S \tilde{\cup} g_S))(d) \cup (\mathbb{S} - (f_S \tilde{\cup} g_S))(d)$, $f_S \tilde{\cup} g_S$ is not a soft C-semigroup over \mathbb{U} .

Proposition 2 Let f_S and f_T be soft C-semigroups over \mathbb{U} and Ψ is a semi-group isomorphism from S to T . If f_S is a soft C-semigroup over \mathbb{U} , then $\Psi(f_S)$ is a soft C-semigroup.

Proof. Let $m_1, m_2 \in T$. Since Ψ is surjective, then there exist $k_1, k_2 \in S$ such that $\Psi(k_1) = m_1$ and $\Psi(k_2) = m_2$. Then,

$$\begin{aligned}
 (\Psi(f_S))(m_1 m_2) &= \bigcap \{f_S(k) : k \in S, \Psi(k) = m_1 m_2\} \\
 &= \bigcap \{f_S(k) : k \in S, k = \Psi^{-1}(m_1 m_2)\} \\
 &= \bigcap \{f_S(k) : k \in S, k = \Psi^{-1}(\Psi(k_1 k_2)) = k_1 k_2\} \\
 &= \bigcap \{f_S(k_1 k_2) : k_i \in S, \Psi(k_i) = m_i, i = 1, 2\} \\
 &\subseteq \bigcap \{(\mathbb{S} - f_S)(k_1) \cup (\mathbb{S} - f_S)(k_2) : m_i \in S, \Psi(k_i) = m_i, i = 1, 2\} \\
 &= (\bigcap \{(\mathbb{S} - f_S)(k_1) : k_1 \in S, \Psi(k_1) = m_1\}) \\
 &\quad \cup (\bigcap \{(\mathbb{S} - f_S)(k_2) : k_2 \in S, \Psi(k_2) = m_2\}) \\
 &\subseteq (\bigcup \{(\mathbb{S} - f_S)(k_1) : k_1 \in S, \Psi(k_1) = m_1\}) \\
 &\quad \cup (\bigcup \{(\mathbb{S} - f_S)(k_2) : k_2 \in S, \Psi(k_2) = m_2\}) \\
 &= \{\mathbb{U} - \bigcap (f_S(k_1)) : k_1 \in S, \Psi(k_1) = m_1\} \\
 &\quad \cup \{\mathbb{U} - \bigcap (f_S(k_2)) : k_2 \in S, \Psi(k_2) = m_2\} \\
 &= (\mathbb{S} - \Psi(f_S))(m_1) \cup (\mathbb{S} - \Psi(f_S))(m_2).
 \end{aligned}$$

Consequently, $\Psi(f_S)$ is a soft C-semigroup over \mathbb{U} . □

Proposition 3 Let f_S and f_T be soft C-semigroups over \mathbb{U} and Ψ be a semigroup homomorphism from S to T . If f_T is a soft C-semigroup over \mathbb{U} , then so $\Psi^{-1}(f_T)$ is.

Proof. Suppose $k_1, k_2 \in S$. Then,

$$\begin{aligned}
 (\Psi^{-1}(f_T))(k_1 k_2) &= f_T(\Psi(k_1 k_2)) \\
 &= f_T(\Psi(k_1) \Psi(k_2)) \\
 &\subseteq (\mathbb{S} - f_T)(\Psi(k_1)) \cup (\mathbb{S} - f_T)(\Psi(k_2)) \\
 &= \mathbb{U} - (f_T(\Psi(k_1)) \cap f_T(\Psi(k_2))) \\
 &= \mathbb{U} - (\Psi^{-1}(f_T(k_1)) \cap \Psi^{-1}(f_T(k_2))) \\
 &= (\mathbb{S} - \Psi^{-1}(f_T))(k_1) \cup (\mathbb{S} - \Psi^{-1}(f_T))(k_2).
 \end{aligned}$$

Hence, $\Psi^{-1}(f_T)$ is a soft C-semigroup over \mathbb{U} . □

3.2 Soft C-left (right), C-ideals of semigroups

In this subsection, soft covered left (right) ideal are introduced and also we survey some properties of these ideals.

Definition 13 Let S be semigroup and f_S be soft set over \mathbb{U} . f_S is called a soft covered left ideal over \mathbb{U} if $f_S(xy) \subseteq (\mathbb{S} - f_S)(y)$; covered right ideal of S over \mathbb{U} if $f_S(xy) \subseteq (\mathbb{S} - f_S)(x)$; covered ideal of S over \mathbb{U} if $f_S(xy) \subseteq (\mathbb{S} - f_S)(y)$ and $f_S(xy) \subseteq (\mathbb{S} - f_S)(x)$ for all $x, y \in S$.

From now on, soft covered left ideal is denoted by soft C-left ideal, soft covered right ideal by soft C-right ideal and soft covered ideal by soft C-ideal for the sake of brevity.

Example 3 Let $S = \{a, b, c, d\}$ in Example 1. Define the soft set f_S over $U = D_4$ as following, $f_S(a) = \emptyset$, $f_S(b) = \{x, y\}$, $f_S(c) = \{e, x^2, yx^2\}$, $f_S(d) = \{x^2, yx, yx^2\}$ and so $(S - f_S)(a) = \{e, x, x^2, y, yx, yx^2\}$, $(S - f_S)(b) = \{e, x^2, yx, yx^2\}$, $(S - f_S)(c) = \{x, y, yx\}$, $(S - f_S)(d) = \{e, x, y\}$. Then, f_S forms a soft C-left ideal of S over U but now suppose $f_S(b) = \{x, x^2, yx, yx^2\}$, then since $f_S(b) = f_S(dd) \not\subseteq (S - f_S)(d)$, f_S is not a soft C-left ideal.

It is easy to see that if $f_S(x) = \emptyset$ for $x \in S$, then f_S is a soft C-left (right) ideal over U . We denote such kind of C-left (right) by \emptyset .

Theorem 6 If X is a C-left ideal of S , then \mathcal{S}_X^c is a soft C-left ideal of S .

Proof. Assume that X is a C-left ideal and $x = mn \in X$. Then, since $X \subseteq S(S - X)$, $x = mn \in S(S - X)$, implying that $m \in S$ and $n \in S - X$. Hence, $\mathcal{S}_X^c(mn) \subseteq (S - \mathcal{S}_X^c)(n)$. In fact,

$$\emptyset = \mathcal{S}_X^c(mn) \subseteq (S - \mathcal{S}_X^c)(n) = U - U = \emptyset.$$

Thus, \mathcal{S}_X^c is a soft C-left ideal of S . □

Theorem 7 Let f_S be a soft set over U . Then, f_S is a soft AC-left (right, ideal) over U of S if and only if f_S^c is a soft C-left (right, ideal) ideal over U of S .

Proof. We give the proof for soft AC-left ideals. Let f_S be a soft AC-left ideal over U of S . In this statement,

$$\begin{aligned} f_S^c(xy) &= (U - f_S)(xy) \\ &\subseteq U - (S - f_S)(y) \\ &\subseteq U - (U - f_S(y)) \\ &= (U - f_S^c(y)) \\ &= (S - f_S^c)(y) \end{aligned}$$

for all $x, y \in S$. Conversely, let f_S^c be a soft C-left ideal over U of S . Then,

$$\begin{aligned} f_S(xy) &= (U - f_S^c)(xy) \\ &\supseteq U - ((S - f_S^c)(y)) \\ &\supseteq U - (U - f_S^c(y)) \\ &= (U - f_S(y)) \\ &= (S - f_S)(y) \end{aligned}$$

for all $x, y \in S$. This completes the proof. \square

Theorem 8 *Let f_S be soft set over U . Then, f_S is a soft C-left ideal over U if and only if*

$$f_S \widetilde{\subseteq} \theta * (S - f_S)$$

Proof. Assume that f_S is a soft C-left ideal over U . In this statement, if $f_S = \emptyset$, it is clear that $f_S \widetilde{\subseteq} (\theta * (S - f_S))$. According to assume, since f_S is a soft C-left ideal for $m, n \in S$ over U , we have:

$$\begin{aligned} (\theta * (S - f_S))(mn) &= \bigcap_{s=mn} \theta(m) \cup (S - f_S)(n) \\ &\supseteq \bigcap_{s=mn} \emptyset \cup (f_S)(mn) \\ &= \bigcap_{s=mn} (f_S)(mn) \\ &= f_S(s) \\ &= f_S(mn) \end{aligned}$$

hence, $f_S \widetilde{\subseteq} \theta * (S - f_S)$. Conversely, suppose that $f_S \widetilde{\subseteq} (\theta * (S - f_S))$. Let $m, n \in S$ and $s = mn$. Then, we have:

$$\begin{aligned} (f_S)(mn) &= (f_S)(s) \\ &\subseteq (\theta * (S - f_S))(mn) \\ &= \bigcap_{mn=xy} \theta(x) \cup (S - f_S)(y) \\ &\subseteq \emptyset \cup (S - f_S)(n) \\ &= (S - f_S)(n). \end{aligned}$$

\square

Theorem 9 *Let f_S be soft set over U . Then, f_S is a soft C-right ideal over U if and only if*

$$f_S \widetilde{\subseteq} (S - f_S) * \theta.$$

Theorem 10 *Let f_S be soft set over U . Then, f_S is a soft C-ideal over U if and only if $f_S \widetilde{\subseteq} \theta * (S - f_S)$ and $f_S \widetilde{\subseteq} (S - f_S) * \theta$.*

Theorem 11 *Let f_S and g_S be two soft subsets of S such that $g_S \widetilde{\subseteq} f_S$. If f_S is a soft C-left ideal, then g_S is also a soft C-left ideal.*

Proof. First assume that f_S is soft C-left ideal and $g_S \widetilde{\subseteq} f_S$. Let $p, q \in S$ and

$s = pq$. Then,

$$\begin{aligned}
 g_S(pq) \subseteq f_S(pq) &\subseteq (\theta * (\mathbb{S} - f_S))(pq) \\
 &= \bigcap_{s=pq} \theta(p) \cup (\mathbb{S} - f_S)(q) \\
 &= \bigcap_{s=pq} \emptyset \cup (\mathbb{S} - f_S)(q) \\
 &= \bigcap_{s=pq} (\mathbb{S} - f_S)(q) \\
 &\subseteq \bigcap_{s=pq} (\mathbb{S} - g_S)(q) \\
 &= \bigcap_{s=pq} \theta(p) \cup (\mathbb{S} - g_S)(q) \\
 &= (\theta * (\mathbb{S} - g_S))(pq).
 \end{aligned}$$

Consequently, $g_S(pq) \subseteq (\theta * (\mathbb{S} - g_S))(pq)$ meaning that g_S is a soft C-left ideal. \square

Definition 14 *Soft union left and soft C-left ideal f_S of S is called a completely soft C-left ideal of S .*

Theorem 12 *Let f_S be a soft subset of S . Then, the followings are equivalent.*

- i) f_S is a completely soft C-left ideal of S .
- ii) $(\forall x, y \in S) f_S(xy) \subseteq f_S(y) \cap (\mathbb{S} - f_S)(y)$.
- iii) $(\forall s \in S^2) f_S(s) \subseteq \bigcap_{s=xy} f_S(y) \cap \bigcap_{s=xy} (\mathbb{S} - f_S)(y)$.

Proof.

$i \Rightarrow ii$ Suppose that f_S is a completely soft C-left ideal of S . Then, since f_S is a soft C-left ideal $f_S(xy) \subseteq (\mathbb{S} - f_S)(y)$. Since f_S is a soft union left ideal of S , $f_S(xy) \subseteq f_S(y)$. This means that $f_S(xy) \subseteq f_S(y) \cap (\mathbb{S} - f_S)(y)$.

$ii \Rightarrow iii$ Accept that $(\forall x, y \in S) f_S(xy) \subseteq f_S(y) \cap (\mathbb{S} - f_S)(y)$. Then, by taking into account the sets, $f_S(xy) \subseteq f_S(y)$ and $f_S(xy) \subseteq (\mathbb{S} - f_S)(y)$. Let $s = xy$ for any $x, y \in S$. This implies that $f_S(s) = f_S(xy) \subseteq \bigcap_{s=xy} f_S(y) \cap \bigcap_{s=xy} (\mathbb{S} - f_S)(y)$.

$iii \Rightarrow i$ Assume that $\forall s = xy \in S^2$, $f_S(s) \subseteq \bigcap_{s=xy} f_S(y) \cap \bigcap_{s=xy} (\mathbb{S} - f_S)(y)$. Thus, $f_S(xy) \subseteq \bigcap_{s=xy} f_S(y)$ and $f_S(xy) \subseteq \bigcap_{s=xy} (\mathbb{S} - f_S)(y)$. Then,

$$\begin{aligned}
 f_S(s) = f_S(xy) &\subseteq \bigcap_{s=xy} (\mathbb{S} - f_S)(y) \\
 &= \bigcap_{s=xy} \emptyset \cup (\mathbb{S} - f_S)(y) \\
 &= \bigcap_{s=xy} \theta(x) \cup (\mathbb{S} - f_S)(y) \\
 &= (\theta * (\mathbb{S} - f_S))(s)
 \end{aligned}$$

and also,

$$\begin{aligned}
 f_S(xy) &\subseteq \bigcap_{s=xy} f_S(y) \\
 &\subseteq f_S(y).
 \end{aligned}$$

Clearly, f_S is a completely soft C-left ideal of S . \square

Theorem 13 *If f_S and g_S are two soft C-left ideals of S over U , then $f_S \widetilde{\cap} g_S$ is a soft C-left ideal over U .*

Proof. Assume that f_S and g_S are two soft C-left ideals. Then,

$$\begin{aligned} (f_S \widetilde{\cap} g_S)(mn) &= f_S(mn) \cap g_S(mn) \\ &\subseteq (\mathbb{S} - f_S)(n) \cap (\mathbb{S} - g_S)(n) \\ &= U - (f_S(n) \cup g_S(n)) \\ &\subseteq U - (f_S(n) \cap g_S(n)) \\ &= (\mathbb{S} - (f_S \widetilde{\cap} g_S))(n) \end{aligned}$$

for all $m, n \in S$. This means that $f_S \widetilde{\cap} g_S$ is a soft C-left ideal over U .

Now, we show that if f_S and g_S are two soft C-left ideals of S over U , then $f_S \widetilde{\cup} g_S$ is not a soft C-left ideal of S with the following example. \square

Example 4 *Let consider the semigroup $S = \{a, b, c, d\}$ over $U = D_4$ in Example 1 and introduce the soft set f_S over U as following $f_S(a) = \emptyset$, $f_S(b) = \{e, x\}$, $f_S(c) = \{x^2, yx, yx^2\}$, $f_S(d) = \{y, x^2, yx^2\}$ so $(\mathbb{S} - f_S)(a) = \{e, x, x^2, y, yx, yx^2\}$, $(\mathbb{S} - f_S)(b) = \{y, x^2, yx, yx^2\}$, $(\mathbb{S} - f_S)(c) = \{e, x, y\}$, $(\mathbb{S} - f_S)(d) = \{e, x, yx\}$. Also let $g_S(a) = \emptyset$, $g_S(b) = \{e\}$, $g_S(c) = \{y, x^2, yx, yx^2\}$, $g_S(d) = \{x, x^2, yx, yx^2\}$ so $(\mathbb{S} - g_S)(a) = \{e, x, x^2, y, yx, yx^2\}$, $(\mathbb{S} - g_S)(b) = \{x, y, x^2, yx^2\}$, $(\mathbb{S} - g_S)(c) = \{e, x\}$, $(\mathbb{S} - g_S)(d) = \{e, y\}$. This shows that $(f_S \widetilde{\cup} g_S)(dd) = (f_S \widetilde{\cup} g_S)(b) \not\subseteq (\mathbb{S} - (f_S \widetilde{\cup} g_S))(d)$.*

Proposition 4 *f_S is a soft C-ideal over U of S if and only if*

$$f_S(mn) \subseteq (\mathbb{S} - f_S)(m) \cap (\mathbb{S} - f_S)(n)$$

for all $m, n \in S$.

Proof. Let f_S be a soft C-ideal of S over U and $m, n \in S$. Then, $f_S(mn) \subseteq (\mathbb{S} - f_S)(m)$ and $f_S(mn) \subseteq (\mathbb{S} - f_S)(n)$. Then, $f_S(mn) \subseteq (\mathbb{S} - f_S)(m) \cap (\mathbb{S} - f_S)(n)$.

Conversely, suppose that $f_S(mn) \subseteq (\mathbb{S} - f_S)(m) \cap (\mathbb{S} - f_S)(n)$ for all $m, n \in S$. It follows that

$$f_S(mn) \subseteq (\mathbb{S} - f_S)(m) \cap (\mathbb{S} - f_S)(n) \subseteq (\mathbb{S} - f_S)(m)$$

and

$$f_S(mn) \subseteq (\mathbb{S} - f_S)(m) \cap (\mathbb{S} - f_S)(n) \subseteq (\mathbb{S} - f_S)(n)$$

so f_S is a soft C-ideal of S over U . \square

Theorem 14 *Let f_S be soft set over \mathcal{U} . If f_S is a soft C-left (right) ideal over \mathcal{U} , f_S is a soft C-semigroup over \mathcal{U} .*

Proof. We give the proof for soft C-left ideals. Similarly, it can be indicated for soft C-right ideals. Let f_S be a soft C-left ideal of S over \mathcal{U} . Then, $f_S(pq) \subseteq (S - f_S)(q)$ for all $p, q \in S$. Thus, $f_S(pq) \subseteq (S - f_S)(q) \subseteq (S - f_S)(p) \cup (S - f_S)(q)$, therefore f_S is a soft C-semigroup. \square

Proposition 5 *Let f_S be soft set over \mathcal{U} . Then, $(S - f_S) \widetilde{\cap} (\theta * (S - f_S))$ is a soft C-left ideal over \mathcal{U} and $(S - f_S) \widetilde{\cap} ((S - f_S) * \theta)$ is a soft C-right ideal of S over \mathcal{U} .*

Proof. Assume that f_S is a soft C-left ideal of S . Then,

$$\begin{aligned} \theta * [(S - f_S) \widetilde{\cap} (\theta * (S - f_S))] &= [(\theta * (S - f_S)) \widetilde{\cap} ((\theta * (\theta * (S - f_S))))] \\ &= (\theta * (S - f_S)) \widetilde{\cap} ((\theta * \theta) * (S - f_S)) \\ &\supseteq (\theta * (S - f_S)) \widetilde{\cap} ((\theta * (S - f_S))) \\ &= (\theta * (S - f_S)) \\ &\supseteq (S - f_S) \widetilde{\cap} (\theta * (S - f_S)). \end{aligned}$$

Clearly, $(S - f_S) \widetilde{\cap} (\theta * (S - f_S))$ is a soft C-left ideal of S over \mathcal{U} . Also,

$$\begin{aligned} [((S - f_S) \widetilde{\cap} ((S - f_S) * \theta))] * \theta &= [((S - f_S) * \theta) \widetilde{\cap} (((S - f_S) * \theta) * \theta)] \\ &= ((S - f_S) * \theta) \widetilde{\cap} (((S - f_S) * \theta * \theta)) \\ &\supseteq ((S - f_S) * \theta) \widetilde{\cap} ((S - f_S) * \theta) \\ &= ((S - f_S) * \theta) \\ &\supseteq (S - f_S) \widetilde{\cap} ((S - f_S) * \theta). \end{aligned}$$

Hence, $(S - f_S) \widetilde{\cap} ((S - f_S) * \theta)$ is a soft C-right ideal of S over \mathcal{U} . \square

Theorem 15 *Let f_S and g_S be a soft C-right ideal and soft C-left ideal of S over \mathcal{U} , respectively. Then,*

$$f_S \widetilde{\cup} g_S \widetilde{\subseteq} (S - f_S) * (S - g_S).$$

Proof. We know that f_S is a soft C-R-right ideal of S over \mathcal{U} and g_S is a soft C-left ideal of S over \mathcal{U} and also $\theta \widetilde{\subseteq} (S - f_S)$, $\theta \widetilde{\subseteq} (S - g_S)$. Thus,

$$g_S \widetilde{\subseteq} \theta * (S - g_S) \widetilde{\subseteq} (S - f_S) * (S - g_S)$$

and

$$f_S \widetilde{\subseteq} (S - f_S) * \theta \widetilde{\subseteq} (S - f_S) * (S - g_S)$$

from here

$$f_S \widetilde{\cup} g_S \subseteq (\mathbb{S} - f_S) * (\mathbb{S} - g_S).$$

Now, we survey that again, let f_S and g_S be a soft C-right ideal a soft C-left ideal of S over U , respectively. In this statement,

$$f_S \widetilde{\cap} g_S \not\subseteq (\mathbb{S} - f_S) * (\mathbb{S} - g_S).$$

with the following example. □

Example 5 Think the semigroup $S = \{a, b, c, d\}$ over $U = D_4$ in Example 1 and let the soft set f_S and g_S be $f_S(b) = \{x, yx\}$, $(\mathbb{S} - f_S)(c) = \{e, x, yx\}$, $(\mathbb{S} - f_S)(d) = \{x, y, yx\}$ and $g_S(b) = \{yx, yx^2\}$, $(\mathbb{S} - g_S)(c) = \{x, yx, yx^2\}$, $(\mathbb{S} - g_S)(d) = \{yx, yx^2\}$. Then, f_S is a soft C-right ideal and g_S is a soft C-left ideal.

$$\begin{aligned} ((\mathbb{S} - f_S) * (\mathbb{S} - g_S))(b) &= \{(\mathbb{S} - f_S)(d) \cup (\mathbb{S} - g_S)(d)\} \cap \{(\mathbb{S} - f_S)(c) \cup (\mathbb{S} - g_S)(c)\} \\ &\cap \{(\mathbb{S} - f_S)(d) \cup (\mathbb{S} - g_S)(c)\} \\ &= \{x, yx, yx^2\} \\ &\not\subseteq (f_S \widetilde{\cap} g_S)(b) \\ &= \{yx\}. \end{aligned}$$

Proposition 6 Let f_S and f_T be soft C-left (right) ideals of S over U . Again, $f_S \widetilde{\wedge} f_T$ is a soft C-left (right) ideal of $S \times T$ over U .

Proof. We accept that f_S and f_T are soft C-left ideals of S over U and $(x_1, y_1), (x_2, y_2) \in S \times T$,

$$\begin{aligned} f_{S \widetilde{\wedge} T}((x_1, y_1), (x_2, y_2)) &= f_{S \widetilde{\wedge} T}(x_1 x_2, y_1 y_2) \\ &= f_S(x_1 x_2) \cap f_T(y_1 y_2) \\ &\subseteq (\mathbb{S} - f_S)(x_2) \cap (\mathbb{S} - f_T)(y_2) \\ &= U - (f_S(x_2) \cup f_T(y_2)) \\ &\subseteq U - (f_S(x_2) \cap f_T(y_2)) \\ &= (\mathbb{S} - f_{S \widetilde{\wedge} T})(x_2, y_2). \end{aligned}$$

Therefore, $f_S \widetilde{\wedge} f_T$ is a soft C-left ideal over U . □

We give following propositions without proof. The proofs are similar to those in section 2.

Proposition 7 Let f_S and h_S be soft sets over U and Ψ is a semigroup isomorphism from S to T . If f_S is a soft C-left (right) ideal of S over U , then so is $\Psi(f_S)$ of T over U .

Proposition 8 Let f_S and h_S be soft sets over \mathcal{U} and Ψ is a semigroup homomorphism from S to T . If f_T is a soft C -left (right) ideal of T over \mathcal{U} , then so is $\Psi^{-1}(f_T)$ of S over \mathcal{U} .

3.3 Soft C -bi-ideals of semigroups

In this subsection, we define soft covered bi-ideals and provide their basic properties by using soft set operations and soft intersection products and also support them with examples.

Definition 15 A soft C -semigroup f_S over \mathcal{U} is called a soft covered bi-ideal over \mathcal{U} if

$$f_S(xyz) \subseteq (\mathbb{S} - f_S)(x) \cup (\mathbb{S} - f_S)(z).$$

For the sake of brevity, soft covered bi-ideal is denoted by soft C -bi-ideal.

Example 6 Define operation over $S = \mathbb{Z}_4 = \{0, 1, 2, 3\}$ by the following table:

.	0	1	2	3
0	0	0	0	0
1	0	0	0	0
2	0	0	0	1
3	0	0	1	2

Now let $\mathcal{U} = D_2 = \{e, x, y, yx\}$ be universal set and f_S be a soft set over \mathcal{U} defined by $f_S(0) = \emptyset$, $f_S(1) = \{x\}$, $f_S(2) = \{x, y\}$, $f_S(3) = \{yx\}$ and so $(\mathbb{S} - f_S)(0) = \mathcal{U}$, $(\mathbb{S} - f_S)(1) = \{e, y, yx\}$, $(\mathbb{S} - f_S)(2) = \{e, yx\}$, $(\mathbb{S} - f_S)(3) = \{e, x, y\}$. Then, f_S is a soft C -bi-ideal, but if $f_S(1) = \{e, yx\}$, then f_S is not a soft C -bi-ideal.

It is easy to see that if $f_S(x) = \emptyset$ for $x \in S$, then f_S is a soft C -bi-ideal over \mathcal{U} . We denote such kind of C -bi-ideal by \emptyset .

Theorem 16 If X is a C -bi-ideal of S , then \mathcal{S}_X is a soft C -bi-ideal of S .

Proof. We accept that X is a C -bi-ideal of S . Let $x = mnp \in X$, then it is clear that $x = mnp \in (S - X)S(S - X)$, implying that $m, p \in S - X$ and $n \in S$. In this statement, $\mathcal{S}_X(mnp) \subseteq (\mathbb{S} - \mathcal{S}_X)(m) \cup (\mathbb{S} - \mathcal{S}_X)(p)$. In fact,

$$\begin{aligned} \emptyset &= \mathcal{S}_X^c(mnp) \\ &\subseteq (\mathbb{S} - \mathcal{S}_X^c)(m) \cup (\mathbb{S} - \mathcal{S}_X^c)(p) \\ &= (\mathcal{U} - \mathcal{U}) \cup (\mathcal{U} - \mathcal{U}) \\ &= \emptyset. \end{aligned}$$

□

Theorem 17 Let f_S be a soft set over U . Then, f_S is a soft AC-bi-ideal over U of S if and only if f_S^c is a soft C-bi-ideal over U of S .

Proof. Let f_S be a soft AC-bi-ideal over U of S . In this statement,

$$\begin{aligned}
 f_S^c(xyz) &= (U - f_S)(xyz) \\
 &\subseteq U - ((S - f_S)(x) \cap (S - f_S)(z)) \\
 &= U - ((U - f_S(x)) \cap (U - f_S(z))) \\
 &= (U - (U - f_S(x))) \cup (U - (U - f_S(z))) \\
 &= (U - f_S^c(x)) \cup (U - f_S^c(z)) \\
 &= (S - f_S^c)(x) \cup (S - f_S^c)(z)
 \end{aligned}$$

for all $x, y, z \in S$. Conversely, let f_S^c be a soft C-bi-ideal over U of S . Then,

$$\begin{aligned}
 f_S(xyz) &= (U - f_S^c)(xyz) \\
 &\supseteq U - ((S - f_S^c)(x) \cup (S - f_S^c)(z)) \\
 &= (U - (U - f_S^c(x))) \cap (U - (U - f_S^c(z))) \\
 &= (U - f_S(x)) \cap (U - f_S(z)) \\
 &= (S - f_S)(x) \cap (S - f_S)(z)
 \end{aligned}$$

for all $x, y, z \in S$. This completes the proof. \square

Theorem 18 A soft subset f_S of S is a soft C-bi-ideal if and only if

$$f_S \widetilde{\subseteq} (S - f_S) * \theta * (S - f_S)$$

Proof. Let f_S be a soft C-bi-ideal of S . Assume that $f_S = \emptyset$, then it is clear $f_S \widetilde{\subseteq} ((S - f_S) * \theta) * (S - f_S)$. Otherwise, let $a, b, t, p, z \in S$. Since f_S is a soft C-bi-ideal of S over U , then,

$$\begin{aligned}
 ((S - f_S) * \theta * (S - f_S))(s) &= \bigcap_{s=ab} [((S - f_S) * \theta)(a) \cup (S - f_S)(b)] \\
 &= \bigcap_{s=ab} [\bigcap_{a=tp} (S - f_S)(t) \cup \theta(p) \cup (S - f_S)(b)] \\
 &= \bigcap_{s=ab} [\bigcap_{a=tp} (S - f_S)(t) \cup \emptyset \cup (S - f_S)(b)] \\
 &= \bigcap_{s=ab} \bigcap_{a=tp} (S - f_S)(t) \cup (S - f_S)(b) \\
 &\supseteq \bigcap_{s=ab} \bigcap_{a=tp} f_S(tpb) \\
 &= \bigcap_{s=ab} f_S(ab) \\
 &= f_S(s)
 \end{aligned}$$

hence, $f_S \widetilde{\subseteq} ((S - f_S) * \theta * (S - f_S))$.

Conversely, let us assume that $f_S \widetilde{\subseteq} ((S - f_S) * \theta * (S - f_S))$. Let $s = abz \in S$,

$$\begin{aligned}
 f_S(abz) &= f_S(s) \\
 &\subseteq ((S - f_S) * \theta * (S - f_S))(s) \\
 &= \bigcap_{s=abz=mn} ((S - f_S) * \theta)(m) \cup (S - f_S)(n) \\
 &\subseteq ((S - f_S) * \theta)(ab) \cup (S - f_S)(z) \\
 &= \bigcap_{ab=tp} (S - f_S)(t) \cup \theta(p) \cup (S - f_S)(z) \\
 &\subseteq (S - f_S)(a) \cup \emptyset \cup (S - f_S)(z) \\
 &= (S - f_S)(a) \cup (S - f_S)(z).
 \end{aligned}$$

Consequently, f_S is a soft C-bi-ideal of S over U . This completes the proof. \square

Theorem 19 *The intersection of two soft C-bi-ideals over U is a soft C-bi-ideal over U .*

Proof. Let f_S and g_S be soft C-bi-ideals over U . Then,

$$\begin{aligned}
 (f_S \widetilde{\cap} g_S)(mnp) &= f_S(mnp) \cap g_S(mnp) \\
 &\subseteq [(S - f_S)(m) \cup (S - f_S)(p)] \cap [(S - g_S)(m) \cup (S - g_S)(p)] \\
 &= [U - (f_S(m) \cap f_S(p))] \cap [U - (g_S(m) \cap g_S(p))] \\
 &= U - [(f_S(m) \cap f_S(p)) \cup (g_S(m) \cap g_S(p))] \\
 &\subseteq U - [(f_S(m) \cap f_S(p) \cap (g_S(m) \cap g_S(p)))] \\
 &= U - ((f_S(m) \cap g_S(m)) \cap ((f_S(p) \cap g_S(p))) \\
 &= U - ((f_S \widetilde{\cap} g_S)(m) \cap (f_S \widetilde{\cap} g_S)(p)) \\
 &= (U - (f_S \widetilde{\cap} g_S)(m)) \cup (U - (f_S \widetilde{\cap} g_S)(p)) \\
 &= (S - (f_S \widetilde{\cap} g_S))(m) \cup (S - (f_S \widetilde{\cap} g_S))(p)
 \end{aligned}$$

for $m, n, p \in S$. This completes the proof. Now, we show that if f_S and g_S are two C-bi-ideals of S over U , then $f_S \widetilde{\cup} g_S$ is not a soft C-bi-ideal of S with the following example. \square

Example 7 *Consider the semigroup $S = Z_4 = \{0, 1, 2, 3\}$ and define the soft set f_S and g_S over $U = D_2$ in Example 6 as following, respectively. $f_S(0) = \emptyset$, $f_S(1) = \{x\}$, $f_S(2) = \{e\}$, $f_S(3) = \{y, yx\}$ so $(S - f_S)(0) = U$, $(S - f_S)(1) = \{e, y, yx\}$, $(S - f_S)(2) = \{x, y, yx\}$, $(S - f_S)(3) = \{e, x\}$ and $g_S(0) = \emptyset$, $g_S(1) = \{e\}$, $g_S(2) = \{e\}$, $g_S(3) = \{x, yx\}$ so $(S - g_S)(0) = U$, $(S - g_S)(1) = \{x, y, yx\}$, $(S - g_S)(2) = \{x, y, yx\}$, $(S - g_S)(3) = \{e, y\}$. Then, f_S and g_S are two C-bi-ideals of S over U , since $(f_S \widetilde{\cup} g_S)(333) = (f_S \widetilde{\cup} g_S)(1) \not\subseteq (S - (f_S \widetilde{\cup} g_S)(3)) \cup (S - (f_S \widetilde{\cup} g_S)(3))$, $(f_S \widetilde{\cup} g_S)$ is not a soft C-bi-ideal.*

Theorem 20 *Every soft C-left (right) ideal of semigroup S over \mathcal{U} is a soft C-bi-ideal of S over \mathcal{U} .*

Proof. Assume that f_S is a soft C-Left ideal of S over \mathcal{U} for $m, p, q \in S$. Then,

$$f_S(mpq) = f_S((mp)q) \subseteq (\mathbb{S} - f_S)(q) \subseteq (\mathbb{S} - f_S)(m) \cup (\mathbb{S} - f_S)(q).$$

Hence, f_S is a soft C-bi-ideal of S . \square

Proposition 9 *Let f_S and f_T be soft C-bi-ideals of S over \mathcal{U} . Then, $f_S \widetilde{\wedge} f_T$ is a soft C-bi-ideal of $S \times T$ over \mathcal{U} .*

Proof. We know that f_S and f_T are soft C-bi-ideals and there exists $(x_1, y_1), (x_2, y_2), (x_3, y_3) \in S \times T$,

$$\begin{aligned} f_{S \widetilde{\wedge} T}((x_1, y_1), (x_2, y_2), (x_3, y_3)) &= f_{S \wedge T}(x_1 x_2 x_3, y_1 y_2 y_3) \\ &= f_S(x_1 x_2 x_3) \cap f_T(y_1 y_2 y_3) \\ &\subseteq [(\mathbb{S} - f_S)(x_1) \cup (\mathbb{S} - f_S)(x_3)] \cap [(\mathbb{S} - f_T)(y_1) \\ &\quad \cup (\mathbb{S} - f_T)(y_3)] \\ &= [\mathcal{U} - (f_S(x_1) \cap f_S(x_3))] \cap [\mathcal{U} - (f_T(y_1) \cap f_T(y_3))] \\ &= \mathcal{U} - [(f_S(x_1) \cap f_S(x_3)) \cup (f_T(y_1) \cap f_T(y_3))] \\ &\subseteq \mathcal{U} - [(f_S(x_1) \cap f_S(x_3)) \cap (f_T(y_1) \cap f_T(y_3))] \\ &= \mathcal{U} - [(f_S(x_1) \cap f_T(x_3)) \cap (f_S(y_1) \cap f_T(y_3))] \\ &= (\mathbb{S} - f_{S \widetilde{\wedge} T})(x_1, y_1) \cup (\mathbb{S} - f_{S \widetilde{\wedge} T})(x_3, y_3). \end{aligned}$$

This shows that $f_S \widetilde{\wedge} f_T$ is a soft C-bi-ideal over \mathcal{U} . \square

Proposition 10 *Let f_S and h_S be soft sets over \mathcal{U} and Ψ is a semigroup isomorphism from S to T . If f_S is a soft C-bi-ideal of S over \mathcal{U} , then so is $\Psi(f_S)$ of T over \mathcal{U} .*

Proposition 11 *Let f_S and h_S be soft sets over \mathcal{U} and Ψ is a semigroup homomorphism from S to T . If f_T is a soft C-bi-ideal of T over \mathcal{U} , then so is $\Psi^{-1}(f_T)$ of S over \mathcal{U} .*

3.4 Soft C-interior ideal of semigroups

In this section, we introduce soft covered interior ideals of semigroups, obtain their basic properties with respect to soft operations and soft intersection product.

Definition 16 Let f_S be soft set over \mathcal{U} and $x, y, z \in S$. If

$$f_S(xyz) \subseteq (\mathbb{S} - f_S)(y)$$

f_S called a soft covered interior ideal over \mathcal{U} .

For the sake of brevity, soft covered interior ideal is denoted by soft C-interior ideal over \mathcal{U} .

Example 8 Let think the semigroup \mathbb{Z}_4 and the soft set f_S over $\mathcal{U} = D_2 = \{e, x, y, yx\}$ in Example 6. Then, one can easily show that f_S is a soft C-interior ideal of S over \mathcal{U} . But we accept that $f_S(1) = \{e, y\}$, then $f_S(333) = f_S(1) \not\subseteq (\mathbb{S} - f_S)(3)$ which implies f_S is not a soft C-interior ideal.

It is easy to see that if $f_S(x) = \emptyset$ for $x \in S$, then f_S is a soft C-interior-ideal over \mathcal{U} . We denote such kind of C-interior-ideal by \emptyset .

Theorem 21 If X is a C-interior ideal of S , then \mathcal{S}_X^c is a soft C-interior ideal of S .

Proof. Let X be C-interior ideal and $x = mnp \in X$. Since $X \subseteq S(S - X)S$, then $x = mnp \in S(S - X)S$, implying that $m, p \in S$ and $n \in S - X$. In this statement, $\mathcal{S}_X^c(mnp) \subseteq (\mathbb{S} - \mathcal{S}_X^c)(n)$. In fact,

$$\begin{aligned} \emptyset &= \mathcal{S}_X^c(mnp) \\ &\subseteq (\mathbb{S} - \mathcal{S}_X^c)(n) \\ &= \mathcal{U} - \mathcal{U} \\ &= \emptyset. \end{aligned}$$

□

Theorem 22 Let f_S be a soft set over \mathcal{U} . Then, f_S is a soft AC-interior ideal over \mathcal{U} of S if and only if f_S^c is a soft C-interior ideal over \mathcal{U} of S .

Proof. Let f_S is a soft AC-interior ideal over \mathcal{U} of S . In this statement,

$$\begin{aligned} f_S^c(xyz) &= (\mathcal{U} - f_S)(xyz) \\ &\subseteq \mathcal{U} - (\mathbb{S} - f_S)(y) \\ &= \mathcal{U} - (\mathcal{U} - f_S(y)) \\ &= \mathcal{U} - f_S^c(y) \\ &= (\mathbb{S} - f_S^c)(y) \end{aligned}$$

for all $x, y, z \in S$. Conversely, let f_S^c be a soft C-interior ideal over \mathcal{U} of S . Then,

$$\begin{aligned} f_S(xyz) &= (\mathcal{U} - f_S^c)(xyz) \\ &\supseteq \mathcal{U} - ((S - f_S^c)(y)) \\ &= \mathcal{U} - (\mathcal{U} - f_S^c(y)) \\ &= \mathcal{U} - f_S(y) \\ &= (S - f_S)(y) \end{aligned}$$

for all $x, y, z \in S$. This completes the proof. \square

Theorem 23 Let f_S be soft set over \mathcal{U} . Then, f_S is a soft C-interior ideal over \mathcal{U} if and only if

$$f_S \widetilde{\subseteq} \theta * (S - f_S) * \theta$$

Proof. Let f_S be a soft C-interior ideal over \mathcal{U} and $x \in S$. If $f_S = \emptyset$, it is clear that $f_S(x) \widetilde{\subseteq} (\theta * (S - f_S) * \theta)(x)$, thus $f_S \widetilde{\subseteq} \theta * (S - f_S) * \theta$. If there exist elements y, z, u, v of S such that $x = yz$ and $y = mp$, we can write:

$$f_S(x) = f_S(yz) = f_S(mpz) \subseteq (S - f_S)(p).$$

Then,

$$\begin{aligned} (\theta * (S - f_S) * \theta)(x) &= ((\theta * (S - f_S)) * \theta)(x) \\ &= \bigcap_{x=yz} (\theta * (S - f_S))(y) \cup \theta(z) \\ &= \bigcap_{x=yz} [\bigcap_{y=mp} (\theta(m) \cup (S - f_S)(p))] \cup \theta(z) \\ &= \bigcap_{x=yz} \bigcap_{y=mp} (\emptyset \cup (S - f_S)(p)) \cup \emptyset \\ &\supseteq \bigcap_{x=yz} \bigcap_{y=mp} \emptyset \cup f_S(mpz) \cup \emptyset \\ &= f_S(x). \end{aligned}$$

Thus, $f_S \widetilde{\subseteq} \theta * (S - f_S) * \theta$.

Conversely, accept that $f_S \widetilde{\subseteq} \theta * (S - f_S) * \theta$ for $x, a, y, m, n, p, q \in S$. Then,

$$\begin{aligned} f_S(xay) &\subseteq (\theta * (S - f_S) * \theta)(xay) \\ &= \bigcap_{xay=mn} \{(\theta * (S - f_S))(m) * \theta(n)\} \\ &\subseteq (\theta * (S - f_S))(xa) \cup \theta(y) \\ &= (\theta * (S - f_S))(xa) \cup \emptyset \\ &= \bigcap_{xa=pq} \theta(p) \cup (S - f_S)(q) \\ &\subseteq \theta(x) \cup (S - f_S)(a) \\ &= (S - f_S)(a) \end{aligned}$$

hence, f_S is a soft C-interior ideal. This completes the proof. \square

Theorem 24 *Every soft union right C-Left ideal of S is a soft C-interior ideal of S .*

Proof. We accept that f_S is a soft union right C-Left ideal. Then,

$$\begin{aligned}
 (\theta * (S - f_S) * \theta)(mnp) &= [(\theta * (S - f_S)) * \theta](mnp) \\
 &= \bigcap_{mnp=uv} [(\theta * (S - f_S))(u) \cup \theta(v)] \\
 &= \bigcap_{mnp=uv} (\theta * (S - f_S))(u) \\
 &\supseteq \bigcap_{mnp=uv} f_S(u) \\
 &\supseteq \bigcap_{mnp=uv} f_S(uv) \\
 &= f_S(mnp)
 \end{aligned}$$

f_S is a C-interior ideal of S . □

Theorem 25 *Let f_S and g_S be two soft C-interior ideals of S over U . Then, $f_S \widetilde{\cap} g_S$ is a soft C-interior ideal over U .*

Proof. Let $m, n, p \in S$. Then,

$$\begin{aligned}
 (f_S \widetilde{\cap} g_S)(mnp) &= f_S(mnp) \cap g_S(mnp) \\
 &\subseteq (S - f_S)(n) \cap (S - g_S)(n) \\
 &\subseteq (S - f_S)(n) \cup (S - g_S)(n) \\
 &= (S - (f_S \widetilde{\cap} g_S))(n)
 \end{aligned}$$

thus, $f_S \widetilde{\cap} g_S$ is a soft C-interior ideal over U . Now, we show that if f_S and g_S are two soft C-interior ideals of S over U , $f_S \widetilde{\cup} g_S$ is not a soft C-interior ideal with the following example. □

Example 9 *Let \mathbb{Z}_4 be the semigroup over $U = D_2 = \{e, x, y, yx\}$ in Example 6. Let the soft set f_S and g_S over U be as following: $f_S(0) = \emptyset$, $f_S(1) = \{e, x\}$, $f_S(2) = \{x\}$, $f_S(3) = \{y, yx\}$ and $g_S(0) = \emptyset$, $g_S(1) = \{e\}$, $g_S(2) = \{y\}$, $g_S(3) = \{x, y, yx\}$. We see that $(f_S \widetilde{\cup} g_S)(333) = (f_S \widetilde{\cup} g_S)(1) \not\subseteq S - (f_S \widetilde{\cup} g_S)(3)$.*

Definition 17 *A soft set f_S over U is defined soft semi prime, if for all $a \in S$,*

$$f_S(a) \subseteq f_S(a^2)$$

Proposition 12 *Let f_S be soft semi prime C-interior ideal of a semigroup S . Then, $f_S(a^n) \subseteq (S - f_S)(a^{n+1})$ for all positive integers n .*

Proof. Let n be any positive integer. Then,

$$f_S(a^n) \subseteq f_S(a^{2n}) \subseteq f_S(a^{4n}) = f_S(a^{3n-2}a^{n+1}a) \subseteq (\mathbb{S} - f_S)(a^{n+1}). \quad \square$$

Proposition 13 *Let f_S and h_S be soft sets over \mathbb{U} and Ψ is a semigroup isomorphism from S to T . If f_S is a soft C -interior ideal of S over \mathbb{U} , then so is $\Psi(f_S)$ of T over \mathbb{U} .*

Proposition 14 *Let f_S and h_S be soft sets over \mathbb{U} and Ψ is a semigroup homomorphism from S to T . If f_S is a soft C -interior ideal of T over \mathbb{U} , then so is $\Psi^{-1}(f_T)$ of S over \mathbb{U} .*

3.5 Soft C -quasi ideals of semigroups

In this subsection, we introduce soft covered quasi-ideals of semigroups, define their basic properties with respect to soft set operations, soft intersection product and certain kinds of soft C -ideals.

Definition 18 *A soft set f_S over \mathbb{U} is called a soft covered quasi-ideal of S over \mathbb{U} if*

$$f_S \widetilde{\subseteq} ((\mathbb{S} - f_S) * \theta) \widetilde{\cup} (\theta * (\mathbb{S} - f_S)).$$

For the sake of brevity, soft covered quasi-ideal of S is denoted by soft C -quasi-ideal.

Proposition 15 *Every soft C -quasi-ideal of S is a soft C -semigroup of S .*

Proof. We accept that f_S is a soft C -quasi-ideal of S . Then, since $\theta \subseteq (\mathbb{S} - f_S)$,

$$\theta * (\mathbb{S} - f_S) \widetilde{\subseteq} (\mathbb{S} - f_S) * (\mathbb{S} - f_S)$$

and

$$(\mathbb{S} - f_S) * \theta \widetilde{\subseteq} (\mathbb{S} - f_S) * (\mathbb{S} - f_S)$$

from here

$$f_S \widetilde{\subseteq} (\theta * (\mathbb{S} - f_S)) \widetilde{\cup} ((\mathbb{S} - f_S) * \theta) \widetilde{\subseteq} (\mathbb{S} - f_S) * (\mathbb{S} - f_S)$$

since f_S is a soft C -quasi-ideal of S . Hence, f_S is a soft C -semigroup of S . \square

Theorem 26 *If X is a C -quasi-ideal of S , then S_X^c is a soft C -quasi-ideal of S .*

Proof. Let X be a C -quasi-ideal and $x = mn \in X$. Since $X \subseteq ((S - X)S) \cup (S(S - X))$, then it is clear that $x = mn \in ((S - X)S) \cup (S(S - X))$, implying that $mn \in (S - X)S$ or $mn \in S(S - X)$ and so $m \in (S - X)$ and $n \in S$ or $m \in S$ and $n \in (S - X)$. Hence since $\mathcal{S}_X^c(x) = \mathcal{S}_X^c(mn) = \emptyset$, in any case

$$\mathcal{S}_X^c \widetilde{\subseteq} ((S - \mathcal{S}_X^c) * \theta) \widetilde{\cup} (\theta * (S - \mathcal{S}_X^c)).$$

In fact, if we consider $[(S - \mathcal{S}_X^c) * \theta] \widetilde{\cup} (\theta * (S - \mathcal{S}_X^c))(x)$ and if $m \in (S - X)$, then

$$\begin{aligned} ((S - \mathcal{S}_X^c) * \theta)(x) &= \bigcap_{x=mn} (S - \mathcal{S}_X^c)(m) \cup \theta(n) \\ &= \emptyset \end{aligned}$$

and

$$\begin{aligned} (\theta * (S - \mathcal{S}_X^c))(x) &= \bigcap_{x=mn} \theta(m) \cup (S - \mathcal{S}_X^c)(n) \\ &= \emptyset \end{aligned}$$

and so $[(S - \mathcal{S}_X^c) * \theta] \widetilde{\cup} (\theta * (S - \mathcal{S}_X^c))(x) = \emptyset \cup \emptyset = \emptyset$. Hence,

$$\mathcal{S}_X^c \widetilde{\subseteq} ((S - \mathcal{S}_X^c) * \theta) \widetilde{\cup} (\theta * (S - \mathcal{S}_X^c)).$$

Also if $m \in X$, then

$$\begin{aligned} ((S - \mathcal{S}_X^c) * \theta)(x) &= \bigcap_{x=mn} (S - \mathcal{S}_X^c)(m) \cup \theta(n) \\ &= \mathcal{U} \end{aligned}$$

and

$$\begin{aligned} (\theta * (S - \mathcal{S}_X^c))(x) &= \bigcap_{x=mn} \theta(m) \cup (S - \mathcal{S}_X^c)(n) \\ &= \mathcal{U} \end{aligned}$$

and so $[(S - \mathcal{S}_X^c) * \theta] \widetilde{\cup} (\theta * (S - \mathcal{S}_X^c))(x) = \mathcal{U} \cup \mathcal{U} = \mathcal{U}$. Hence,

$$\mathcal{S}_X^c \widetilde{\subseteq} ((S - \mathcal{S}_X^c) * \theta) \widetilde{\cup} (\theta * (S - \mathcal{S}_X^c)).$$

□

Proposition 16 *Every soft C -left (right) ideal of S is a soft C -quasi-ideal of S .*

Proof. We accept that f_S is a soft C -Left ideal of S over \mathcal{U} which is defined $f_S \subseteq (\theta * (S - f_S))$. In this statement, we have:

$$f_S \widetilde{\subseteq} (\theta * (S - f_S)) \widetilde{\subseteq} ((S - f_S) * \theta) \widetilde{\cup} (\theta * (S - f_S))$$

hence, f_S is a soft C -quasi-ideal. □

The converse of the above proposition does not hold in general as shown in the table.

Example 10 Think $S = \mathbb{Z}_4 = \{0, 1, 2, 3\}$ defined by the following table:

.	0	1	2	3
0	0	0	0	0
1	0	0	0	0
2	0	0	0	0
3	0	0	1	1

Let f_S be a soft set over $\mathcal{U} = D_2 = \{e, x, y, yx\}$ and is defined as following, $f_S(0) = \emptyset$, $f_S(1) = \{e\}$, $f_S(2) = \{e\}$, $f_S(3) = \{y, yx\}$ and so $(S - f_S)(0) = \mathcal{U}$, $(S - f_S)(1) = \{x, y, yx\}$, $(S - f_S)(2) = \{x, y, yx\}$, $(S - f_S)(3) = \{e, x\}$. Then, one can show that $f_S(1) \subseteq ((S - f_S) * \theta)(1) \cup (\theta * (S - f_S))(1) = (S - f_S)(3) \cup ((S - f_S)(3) \cap (S - f_S)(2)) = (S - f_S)(3) = \{e, x\}$ is a soft C -quasi-ideal but since $f_S(3.2) = f_S(1) \not\subseteq (S - f_S)(2)$, f_S is not a soft C -Left ideal.

Proposition 17 Every soft C -quasi-ideal is a soft C -bi-ideal of S .

Proof. Let f_S be a soft C -quasi-ideal of S . Then,

$$\begin{aligned}
 f_S &\subseteq ((S - f_S) * \theta) \widetilde{\cup} (\theta * (S - f_S)) \\
 &\subseteq ((S - f_S) * (\theta * \theta)) \widetilde{\cup} ((\theta * \theta) * (S - f_S)) \\
 &\subseteq ((S - f_S) * \theta * (S - f_S)) \widetilde{\cup} ((S - f_S) * \theta) * (S - f_S) \\
 &= ((S - f_S) * \theta * (S - f_S)) \widetilde{\cup} ((S - f_S) * \theta * (S - f_S)) \\
 &= (S - f_S) * \theta * (S - f_S)
 \end{aligned}$$

Hence, f_S is a soft C -bi-ideal of S together with Proposition 15. The converse of this proposition does not hold in general as shown in the following example. \square

Example 11 Let $S = \{a, b, c, d\}$ be the semigroup in Example 6 and $\mathcal{U} = D_2 = \{e, x, y, yx\}$. Let f_S be a soft set over \mathcal{U} defined by $f_S(0) = \emptyset$, $f_S(1) = \{x\}$, $f_S(2) = \{x, y\}$, $f_S(3) = \{yx\}$ and so $(S - f_S)(0) = \mathcal{U}$, $(S - f_S)(1) = \{e, y, yx\}$, $(S - f_S)(2) = \{e, yx\}$, $(S - f_S)(3) = \{e, x, y\}$. Then, f_S is a soft C -bi-ideals. However, f_S is not a soft C -quasi-ideal since $f_S(1) = \{x\} \not\subseteq ((S - f_S) * \theta)(1) \cup (\theta * (S - f_S))(1) = (S - f_S)(2) \cap (S - f_S)(3) = \{e\}$.

Proposition 18 Let f_S and g_S be any soft C -Right ideal and soft C -Left ideal of S over \mathcal{U} , respectively. Then, $f_S \widetilde{\cap} g_S$ is a soft C -quasi-ideal.

Proof. Let f_S be any soft C-Right ideal of S and g_S be any soft C-Left ideal of S . Then,

$$\begin{aligned} (((S - (f_S \widetilde{\cap} g_S)) * \theta) \widetilde{\cup} (\theta * (S - (f_S \widetilde{\cap} g_S)))) &\supseteq (((S - (f_S \widetilde{\cap} g_S)) * \theta) \\ &\widetilde{\cap} (\theta * (S - (f_S \widetilde{\cap} g_S)))) \\ &\supseteq ((S - f_S) * \theta) \widetilde{\cap} (\theta * (S - g_S)) \\ &\supseteq f_S \widetilde{\cap} g_S. \end{aligned}$$

Thus, $f_S \widetilde{\cap} g_S$ is a soft C-quasi-ideal of S over \mathcal{U} . \square

Proposition 19 *Let f_S and g_S be any soft C-quasi-ideals of S over \mathcal{U} . Then, $f_S \widetilde{\cap} g_S$ is a soft C-quasi-ideal.*

Proof. Let f_S and g_S be any soft C-quasi-ideals of S . Then,

$$\begin{aligned} (((S - (f_S \widetilde{\cap} g_S)) * \theta) \widetilde{\cup} (\theta * (S - (f_S \widetilde{\cap} g_S)))) &\supseteq ((S - f_S) * \theta) \widetilde{\cup} (\theta * (S - f_S)) \\ &\supseteq f_S \end{aligned}$$

and

$$\begin{aligned} (((S - (f_S \widetilde{\cap} g_S)) * \theta) \widetilde{\cup} (\theta * (S - (f_S \widetilde{\cap} g_S)))) &\supseteq ((S - g_S) * \theta) \widetilde{\cup} (\theta * (S - g_S)) \\ &\supseteq g_S. \end{aligned}$$

Thus, $((S - (f_S \widetilde{\cap} g_S)) * \theta) \widetilde{\cup} (\theta * (S - (f_S \widetilde{\cap} g_S))) \supseteq (f_S \widetilde{\cap} g_S)$. Then, $f_S \widetilde{\cap} g_S$ is a soft C-quasi-ideal. \square

Proposition 20 *Let f_S be any soft C-quasi-ideal of a commutative semigroup S and $a \in A$. Then,*

$$f_S(a^{n+1}) \subseteq (S - f_S)(a^n)$$

for every positive integer n .

Proof. For any positive integer n , we have:

$$\begin{aligned} ((S - f_S) * \theta)(a^{n+1}) &= \bigcap_{a^{n+1}=xy} (S - f_S)(x) \cup \theta(y) \\ &\subseteq (S - f_S)(a^n) \cup \emptyset \\ &= (S - f_S)(a^n) \end{aligned}$$

from here $((S - f_S) * \theta)(a^{n+1}) \subseteq (S - f_S)(a^n)$.

Moreover, since f_S is a soft C-quasi-ideal of S , we have:

$$\begin{aligned} f_S(a^{n+1}) &\subseteq [((S - f_S) * \theta) \widetilde{\cup} (\theta * (S - f_S))](a^{n+1}) \\ &= ((S - f_S) * \theta)(a^{n+1}) \cup (\theta * (S - f_S))(a^{n+1}) \\ &\subseteq (S - f_S)(a^n) \cup (S - f_S)(a^n) \\ &= (S - f_S)(a^n). \end{aligned}$$

This completes the proof. \square

3.6 Soft C-generalized Bi-ideals of semigroups

In this subsection, we study soft covered generalized bi-ideals of semigroups, introduce their basic prosperities as regards soft set operations, soft intersection product and certain kinds of soft C-ideals.

Definition 19 *A soft set over \mathcal{U} is called a soft covered generalized bi-ideal of S , if*

$$f_S(xyz) \subseteq (\mathbb{S} - f_S)(x) \cup (\mathbb{S} - f_S)(z)$$

for all $x, y, z \in S$.

For the sake of brevity, soft covered generalized bi-ideal is denoted by soft C-generalized bi-ideal of S . Clearly, every soft C-bi-ideal of S is a soft C-generalized bi-ideal of S but converse of this statement is not true. This is indicated by following example.

Example 12 *Let think the semigroup $S = \{0, 1, 2, 3\}$ in Example 6 and define the soft set f_S over $\mathcal{U} = D_4$ such that $f_S(0) = \emptyset$, $f_S(1) = \{x, x^2, y\}$, $f_S(2) = \{x, yx\}$, $f_S(3) = \{e, yx, yx^2\}$ and so $(\mathbb{S} - f_S)(0) = \{e, x, x^2, y, yx, yx^2\}$, $(\mathbb{S} - f_S)(1) = \{e, yx, yx^2\}$, $(\mathbb{S} - f_S)(2) = \{e, y, x^2, yx^2\}$, $(\mathbb{S} - f_S)(3) = \{x, y, x^2\}$.*

Then, one can easily show that f_S is a soft C-generalized bi-ideal of S over \mathcal{U} . However, since $f_S(33) = f_S(2) \not\subseteq (\mathbb{S} - f_S)(3) \cup (\mathbb{S} - f_S)(3)$, f_S is not a soft C-bi-ideal of S .

Theorem 27 *If X is a C-generalized bi-ideal of S , then S_X^c is a soft C-generalized bi-ideal of S .*

Theorem 28 *Let f_S be a soft set over \mathcal{U} . Then, f_S is a soft C-generalized bi-ideal over \mathcal{U} of S if and only if f_S^c is a soft C-generalized bi-ideal over \mathcal{U} of S .*

Theorem 29 *Let f_S be a soft set over \mathcal{U} . Then, f_S is a soft C-generalized bi-ideal of S over \mathcal{U} if and only if*

$$f_S \widetilde{\subseteq} (\mathbb{S} - f_S) * \theta * (\mathbb{S} - f_S).$$

Theorem 30 *Every soft C-left (right) ideal of a semigroup S over \mathcal{U} is a soft C-generalized bi-ideal of S over \mathcal{U} .*

Proof. Let f_S be a soft C-left (right) ideal of S over \mathcal{U} and $x, y, z \in S$. Then,

$$f_S(xyz) = f_S((xy)z) \subseteq (\mathbb{S} - f_S)(z) \subseteq (\mathbb{S} - f_S)(x) \cup (\mathbb{S} - f_S)(z)$$

Thus, f_S is a soft C-generalized bi-ideal of S . □

Proposition 21 Let f_S and f_T be soft C -generalized bi-ideals of S over \mathcal{U} . Then, $f_S \tilde{\wedge} f_T$ is a soft C -generalized bi-ideal of $S \times T$ over \mathcal{U} .

Proposition 22 Let f_S and h_S be soft sets over \mathcal{U} and Ψ is a semigroup isomorphism from S to T . If f_S is a soft C -generalized bi-ideal of S over \mathcal{U} , then so is $\Psi(f_S)$ of T over \mathcal{U} .

Proposition 23 Let f_S and h_S be soft sets over \mathcal{U} and Ψ is a semigroup homomorphism from S to T . If f_T is a soft C -generalized bi-ideal of T over \mathcal{U} , then so is $\Psi^{-1}(f_T)$ of S over \mathcal{U} .

4 Conclusion

In this manuscript, we have introduced soft C -semigroups, C -left (right) ideals, C -bi-ideals, C -interior ideals, C -quasi-ideals and C -generalized bi-ideals. Moreover, we survey the relation between soft AC -ideals and soft C -ideals. Addition to, we obtain the interrelations of various soft C -ideals as in the following figure.




Figure 1

References

- [1] D. Molodtsov, Soft set theory-first results, *Comput Math Appl.*, **37** (1999), 19–31.
- [2] H. Aktas and N. Çağman, Soft sets and soft groups, *Inform Sci.*, **177** (2007), 2726–2735.
- [3] F. Feng, Y. B. Jun and X. Zhao, Soft semirings, *Comput Math Appl.*, **56** (2008), 2621–2628.
- [4] U. Acar, F. Koyuncu and B. Tanay, Soft sets and soft rings, *Comput Math Appl.*, **59** (2010), 3458–3463.
- [5] Y. B. Jun, Soft BCK/BCI-algebras, *Comput Math Appl.*, **56** (2008), 1408–1413.
- [6] Y. B. Jun and C. H. Park, Applications of soft sets in ideal theory of BCK/BCI-algebras, *Inform Sci.*, **178** (2008), 2466–2475.
- [7] Y. B. Jun, K. J. Lee and J. Zhan, Soft \mathfrak{p} -ideals of soft BCI-algebras, *Comput Math Appl.*, **58** 2009, 2060–2068.
- [8] Y. B. Jun, K. J. Lee, C. H. Park, Soft set theory applied to ideals in d-algebras, *Comput. Math. Appl.*, **57** (3) (2009), 367–378.
- [9] J. Zhan, Y. B. Jun, Soft BL-algebras based on fuzzy sets, *Comput Math Appl.*, **59** (6) (2010), 2037–2046.
- [10] O. Kazancı, Ş. Yılmaz and S. Yamak, Soft sets and soft BCH-algebras, *Hacet J Math Stat.*, **39** (2) (2010), 205–217.
- [11] A. Sezgin and A. O. Atagün and E. Aygün, A note on soft N-groups and idealistic soft N-groups, *Filomat*, **25** (1), 53–68.
- [12] N. Çağman and S. Enginoğlu, Soft matrix theory and its decision making, *Comput Math Appl.*, **59** (2010), 3308–3314.
- [13] N. Çağman and S. Enginoğlu, Soft set theory and uni-int decision making, *Eur J Oper Res.*, **207** (2010), 848–855.
- [14] P. K. Maji, A. R. Roy and R. Biswas, An application of soft sets in a decision making problem, *Comput Math Appl.*, **44** (2002), 1077–1083.

- [15] Y. Zou and Z. Xiao, Data analysis approaches of soft sets under incomplete information, *Knowled-Based Syst.*, **21** (2008), 941–945.
- [16] D. Mandal, Fuzzy ideals and fuzzy interior idelas in ordered semigroups, In *Fuzzy Information and Engineering*, (2014), 101–114.
- [17] K. A. Dib and N. Galhum, Fuzzy ideals and fuzzy bi-ideals in fuzzy semigroups, *Fuzzy Sets and Systems*, **92** (1992), 103–111.
- [18] O. Kazancı and S. Yamak, Generalized fuzzy bi-ideals of semigroup, *Soft Computing*, **12** (2008), 1119–1124.
- [19] J. Kavikumar, A. B. Khamis, Fuzzy Ideals and Fuzzy Quasi-ideals in Ternary Semirings, *IAENG International Journal of Applied Mathematics*, **37** (2) (2007).
- [20] T. Changphas and P. Summaprab, On Ordered Semigroups Containing Covered Ideals, *Communications in Algebra*, **44** (9) (2016), 4104–4113, DOI: 10.1080/00927872.2015.1087015.
- [21] A. S. Sezgin, N. Çağman, A. O. Atagün, M. I. Ali, Türkmen E. Soft intersection semigroups, ideals and bi-ideals; a new application on semigroup theory I, *Filomat*, **29** (5) (2015), 917–946.
- [22] A. S. Sezgin, N. Çağman, A. O. Atagün, Soft intersection interior ideals, quasi-ideals and generalized bi-ideals; a new approach to semigroup theory II, *Journal of Multiple-Valued Logic and Soft Computing*, **23** (1–2) (2014), 161–207.
- [23] I. Fabrici, Semigroups containing covered one-sided ideals, *Mahematica Slovaca*, **31** (1981), 225–231.
- [24] I. Fabrici, Semigroups containing covered two-sided ideals, *Mahematica Slovaca*, **34** (1984), 355–363.
- [25] X., Xie, F. Yan, Fuzzy Anti-Covered Left Ideals in Semigroups, submitted.
- [26] A. Sezgin, Ş Özlü, Soft Anti-Covered Ideals in Semigroups, submitted.
- [27] N. Çağman, F. Çitak and H. Aktaş, Soft int-groups and its applications to group theory, *Neural Comput. Appl.*, **21** (1) (2012), 151–158.

Received: November 20, 2019



Some fixed point results on S-metric spaces

Maryam Shahraki

Department of Mathematics,
Qaemshahr Branch, Islamic Azad
University, Qaemshahr, Iran
email: m.shahraki@gu.ac.ir

Shaban Sedghi*

Department of Mathematics,
Qaemshahr Branch, Islamic Azad
University, Qaemshahr, Iran
email: sedghi.gh@qaemiau.ac.ir

S. M. A. Aleomraninejad

Department of Mathematics,
Faculty of Science, Qom University
of Technology, Qom, Iran
email: aleomran@qut.ac.ir

Zoran D. Mitrović

University of Banja Luka, Faculty of
Electrical Engineering, 78000 Banja
Luka, Bosnia and Herzegovina
email: zoran.mitrovic@etf.unibl.org

Abstract. In this paper, a general form of the Suzuki type function is considered on S- metric space, to get a fixed point. Then we show that our results generalize some old results.

1 Introduction and preliminaries

In 1922, Banach [1] proposed a theorem, which is well-known as Banach's Fixed Point Theorem (or Banach's Contraction Principle, BCP for short) to establish the existence of solutions for nonlinear operator equations and integral equations. Since then, because of simplicity and usefulness, it has become a very popular tool in solving a variety of problems such as control theory, economic theory, nonlinear analysis and global analysis. Later, a huge amount

2010 Mathematics Subject Classification: 47H10, 54H25

Key words and phrases: fixed point, S-metric space, Suzuki methods

*Corresponding author

of literature is witnessed on applications, generalizations and extensions of this theorem. They are carried out by several authors in different directions, e.g., by weakening the hypothesis, using different setups.

Many mathematics problems require one to find a distance between two or more objects which is not easy to measure precisely in general. There exist different approaches to obtaining the appropriate concept of a metric structure. Due to the need to construct a suitable framework to model several distinguished problems of practical nature, the study of metric spaces has attracted and continues to attract the interest of many authors. Over last few decades, a numbers of generalizations of metric spaces have thus appeared in several papers, such as 2-metric spaces, G-metric spaces, D*-metric spaces, partial metric spaces and cone metric spaces. These generalizations were then used to extend the scope of the study of fixed point theory. For more discussions of such generalizations, we refer to [3, 4, 5, 6, 7, 8, 9, 10, 12, 13, 20, 21, 22, 23]. Sedghi et al [17] have introduced the notion of an S-metric space and proved that this notion is a generalization of a G-metric space and a D*-metric space. Also, they have proved properties of S-metric spaces and some fixed point theorems for a self-map on an S-metric space.

The Banach contraction principle is the most powerful tool in the history of fixed point theory. Boyd and Wong [2] extended the Banach contraction principle to the nonlinear contraction mappings. We begin by briefly recalling some basic definitions and results for S-metric spaces that will be needed in the sequel. For more details please see [1, 14, 18].

Definition 1 [17] *Let X be a (nonempty) set, an S-metric on X is a function $S : X^3 \rightarrow [0, +\infty)$ that satisfies the following conditions, for each $x, y, z, a \in X$,*

- (1). $S(x, y, z) \geq 0$,
- (2). $S(x, y, z) = 0$ if and only if $x = y = z$,
- (3). $S(x, y, z) \leq S(x, x, a) + S(y, y, a) + S(z, z, a)$,

for all $x, y, z, a \in X$.

The pair (X, S) is called an S-metric space.

Immediate examples of such S-metric spaces are:

Example 1 [15, 18] *Let $X = \mathbb{R}^n$ and $\| \cdot \|$ a norm on X , then*

$$S(x, y, z) = \|y + z - 2x\| + \|y - z\|$$

is an S -metric on X .

Let X be a nonempty set, d is ordinary metric on X , then

$$S(x, y, z) = d(x, z) + d(y, z)$$

is an S -metric on X . This S -metric is called the usual S -metric on X .

Definition 2 [16] Let (X, S) be an S -metric space.

- (i) A sequence $\{x_n\} \subset X$ converges to $x \in X$ if $S(x_n, x_n, x) \rightarrow 0$ as $n \rightarrow +\infty$. That is, for each $\varepsilon > 0$, there exists $n_0 \in \mathbb{N}$ such that for all $n \geq n_0$ we have $S(x_n, x_n, x) < \varepsilon$. We write $x_n \rightarrow x$ for brevity.
- (ii) A sequence $\{x_n\} \subset X$ is a Cauchy sequence if $S(x_n, x_n, x_m) \rightarrow 0$ as $n, m \rightarrow +\infty$. That is, for each $\varepsilon > 0$, there exists $n_0 \in \mathbb{N}$ such that for all $n, m \geq n_0$ we have $S(x_n, x_n, x_m) < \varepsilon$.
- (iii) The S -metric space (X, S) is complete if every Cauchy sequence is a convergent sequence.

Definition 3 [15] Let (X, S) be an S -metric space. For $r > 0$ and $x \in X$ we define the open ball $B_s(x, r)$ and closed ball $B_s[x, r]$ with center x and radius r as follows respectively:

$$\begin{aligned} B_s(x, r) &= \{y \in X : S(y, y, x) < r\}, \\ B_s[x, r] &= \{y \in X : S(x, x, y) \leq r\}. \end{aligned}$$

Example 2 [15] Let $X = \mathbb{R}$ and $S(x, y, z) = |y + z - 2x| + |y - z|$ for all $x, y, z \in \mathbb{R}$. Then

$$\begin{aligned} B_s(1, 2) &= \{y \in \mathbb{R} : S(y, y, 1) < 2\} = \{y \in \mathbb{R} : |y - 1| < 1\} \\ &= \{y \in \mathbb{R} : 0 < y < 2\} = (0, 2). \end{aligned}$$

Lemma 1 [16] Let (X, S) be an S -metric space. If $r > 0$ and $x \in X$, then the ball $B_s(x, r)$ is open subset of X .

Lemma 2 [15, 16, 18] In an S -metric space, we have $S(x, x, y) = S(y, y, x)$.

Proof. By third condition of S -metric, we have

$$\begin{aligned} S(x, x, y) &\leq S(x, x, x) + S(x, x, x) + S(y, y, x) \\ &= S(y, y, x) \end{aligned} \tag{1}$$

$$\begin{aligned} S(y, y, x) &\leq S(y, y, y) + S(y, y, y) + S(x, x, y) \\ &= S(x, x, y), \end{aligned} \quad (2)$$

hence by (1) and (2), we get $S(x, x, y) = S(y, y, x)$. \square

Lemma 3 [18] *Let (X, S) be an S -metric space. If sequence $\{x_n\}$ in X converges to x , then x is unique.*

Lemma 4 [18] *Let (X, S) be an S -metric space. If sequence $\{x_n\}$ in X converges to x , then $\{x_n\}$ is a Cauchy sequence.*

Lemma 5 [15, 16, 18] *Let (X, S) be an S -metric space. If there exist sequences $\{x_n\}$ and $\{y_n\}$ such that $\lim_{n \rightarrow +\infty} x_n = x$ and $\lim_{n \rightarrow +\infty} y_n = y$, then $\lim_{n \rightarrow +\infty} S(x_n, x_n, y_n) = S(x, x, y)$.*

Definition 4 [15, 19] *Let X be a (nonempty) set, a b -metric on X is a function $d : X^2 \rightarrow [0, +\infty)$ if there exists a real number $b \geq 1$ such that the following conditions hold for all $x, y, z \in X$,*

- (1) $d(x, y) = 0$ if and only if $x = y$,
- (2) $d(x, y) = d(y, x)$,
- (3) $d(x, z) \leq b[d(x, y) + d(y, z)]$.

The pair (X, d) is called a b -metric space.

Proposition 1 [16] *Let (X, S) be an S -metric space and let*

$$d(x, y) = S(x, x, y),$$

for all $x, y \in X$. Then we have

- (1) d is a b -metric on X ;
- (2) $x_n \rightarrow x$ in (X, S) if and only if $x_n \rightarrow x$ in (X, d) ;
- (3) $\{x_n\}$ is a Cauchy sequence in (X, S) if and only if $\{x_n\}$ is a Cauchy sequence in (X, d) .

Definition 5 *Let \mathcal{L} be the set of all continuous functions $g : [0, \infty)^4 \rightarrow [0, +\infty)$, satisfying the conditions:*

- (i) $g(1, 1, 1, 1) < 1$,
- (ii) g is subhomogeneous, i.e., $g(\alpha x_1, \alpha x_2, \alpha x_3, \alpha x_4) \leq \alpha g(x_1, x_2, x_3, x_4)$, for all $\alpha \geq 0$,

- (iii) if $x_i, y_i \in [0, +\infty)$, $x_i \leq y_i$ for $i = 1, \dots, 4$ we have $g(x_1, x_2, x_3, x_4) \leq g(y_1, y_2, y_3, y_4)$.

Example 3 The function $g(x_1, x_2, x_3, x_4) = k \max\{x_i\}_{i=0}^4$ for $k \in (0, 1)$ is in class \mathcal{L} .

Example 4 The function $g(x_1, x_2, x_3, x_4) = k \max\{x_1, x_2, \frac{x_3+x_4}{2}\}$ for $k \in (0, 1)$ is in class \mathcal{L} .

Proposition 2 If $g \in \mathcal{L}$ and $u, v \in [0, +\infty]$ are such that $u \leq g(v, v, v, u)$, then $u \leq hv$, where $h = g(1, 1, 1, 1)$.

Proof. If $v < u$, then

$$u \leq g(v, v, v, u) \leq g(u, u, u, u) < ug(1, 1, 1, 1) = hu < u,$$

which is a contradiction. Thus $u \leq v$, which implies

$$u \leq g(v, v, v, u) \leq g(v, v, v, v) < vg(1, 1, 1, 1) = hv. \quad \square$$

Corollary 1 [15] Let (X, S) be a complete S -metric space and $T : X \rightarrow X$ a function such that for, all $x, y, z, a \in X$,

$$S(Tx, Ty, Tz) \leq LS(x, y, z),$$

where $L \in (0, 1/2)$. Then there exists a unique point $u \in X$ such that $Tu = u$.

2 Results

Now, we give our main result.

Theorem 1 Let (X, S) be a S -metric space and $T : X \rightarrow X$ be a function. Suppose that there exist $g \in \mathcal{L}$ and $\alpha \in (0, 1)$, such that $\alpha(h+2) \leq 1$ where $h = g(1, 1, 1, 1)$. Suppose also that $\alpha S(x, x, Tx) \leq S(x, y, z)$ implies

$$S(Tx, Ty, Tz) \leq g(S(x, y, z), S(x, x, Tx), S(y, y, Ty), S(z, z, Tz)),$$

for all $x, y, z \in X$. Then $F(T)$ is non-empty set.

Proof. Fix arbitrary $x_0 \in X$ and let $Tx_0 = x_1$. Since

$$\alpha S(x_0, x_0, Tx_0) < S(x_0, x_0, x_1),$$

then by the hypothesis of the theorem and condition (iii) Definition 5, respectively, we have

$$\begin{aligned} S(x_1, x_1, Tx_1) &= S(Tx_0, Tx_0, Tx_1) \\ &\leq g(S(x_0, x_0, x_1), S(x_0, x_0, Tx_0), S(x_0, x_0, Tx_0), S(x_1, x_1, Tx_1)) \\ &= g(S(x_0, x_0, x_1), S(x_0, x_0, x_1), S(x_0, x_0, x_1), S(x_1, x_1, Tx_1)) \end{aligned}$$

Then, by Proposition 2, we have $S(x_1, x_1, Tx_1) \leq hS(x_0, x_0, x_1)$.

Now let $Tx_1 = x_2$. Since $\alpha S(x_1, x_1, Tx_1) < S(x_1, x_1, x_2)$, by using and the properties of the function g we have

$$\begin{aligned} S(x_2, x_2, Tx_2) &= S(Tx_1, Tx_1, Tx_2) \\ &\leq g(S(x_1, x_1, x_2), S(x_1, x_1, Tx_1), S(x_1, x_1, Tx_1), S(x_2, x_2, Tx_2)) \\ &= g(S(x_1, x_1, x_2), S(x_1, x_1, x_2), S(x_1, x_1, x_2), S(x_2, x_2, Tx_2)). \end{aligned}$$

Then, by Proposition 2, we have $S(x_2, x_2, Tx_2) \leq hS(x_1, x_1, x_2)$.

In a similar way, we can let $Tx_2 = x_3$. So we have

$$S(x_2, x_2, x_3) < hS(x_1, x_1, x_2) < h^2S(x_0, x_0, x_1).$$

By continuing this process, we obtain a sequence $\{x_n\}_{n \geq 1}$ in X such that $x_{n+1} = Tx_n$, which satisfies $S(x_n, x_n, Tx_n) \leq hS(x_{n-1}, x_{n-1}, x_n)$ and

$$S(x_n, x_n, x_{n+1}) \leq h^n S(x_0, x_0, x_1).$$

If $x_m = x_{m+1}$ for some $m \geq 1$, then

Then T has a fixed point.

Suppose that $x_n \neq x_{n+1}$, for all $n \geq 1$. Repeated application of the triangle inequality implies

$$\begin{aligned} S(x_n, x_n, x_{n+m}) &\leq 2S(x_n, x_n, x_{n+1}) + S(x_{n+m}, x_{n+m}, x_{n+1}) \\ &= 2S(x_n, x_n, x_{n+1}) + S(x_{n+1}, x_{n+1}, x_{n+m}) \\ &\leq 2S(x_n, x_n, x_{n+1}) + 2S(x_{n+1}, x_{n+1}, x_{n+2}) \\ &\quad + S(x_{n+m}, x_{n+m}, x_{n+2}) \\ &\leq 2[S(x_n, x_n, x_{n+1}) + S(x_{n+1}, x_{n+1}, x_{n+2}) \\ &\quad + \cdots + S(x_{n+m-1}, x_{n+m-1}, x_{n+m})] \end{aligned}$$

$$\leq 2 \sum_{k=0}^{k=m-1} h^{k+n} S(x_0, x_0, x_1) \leq \frac{2h^n}{1-h} S(x_0, x_0, x_1).$$

So we get

$$\lim_{n \rightarrow +\infty} S(x_n, x_n, x_{n+m}) \rightarrow 0$$

and hence $\{x_n\}_{n \geq 1}$ is a Cauchy sequence in (X, S) . Regarding Definition 2, $\{x_n\}_{n \geq 1}$ is also a Cauchy sequence in (X, S) .

Since (X, S) is a complete S -metric space, by Definition 2, (X, S) is also complete.

Thus $\{x_n\}_{n \geq 1}$ converges to a limit, say, $x \in X$, that is,

$$\lim_{n \rightarrow +\infty} S(x_n, x_n, x) = 0.$$

It is easy to see that $\lim_{n \rightarrow \infty} S(x_n, x_{n+1}, x) = 0$. Now, we claim that for each $n \geq 1$ one of the relations

$$\alpha S(x_n, x_n, Tx_n) \leq S(x_n, x_n, x)$$

or

$$\alpha S(x_{n+1}, x_{n+1}, Tx_{n+1}) \leq S(x_n, x_n, x)$$

holds. If for some $n \geq 1$ we have

$$\alpha S(x_n, x_n, Tx_n) > S(x_n, x_n, x) \text{ and } \alpha S(x_{n+1}, x_{n+1}, Tx_{n+1}) > S(x_{n+1}, x_{n+1}, x),$$

then

$$\begin{aligned} S(x_n, x_n, x_{n+1}) &\leq 2S(x_n, x_n, x) + S(x_{n+1}, x_{n+1}, x) \\ &< 2\alpha S(x_n, x_n, Tx_n) + \alpha S(x_{n+1}, x_{n+1}, Tx_{n+1}) \\ &= 2\alpha S(x_n, x_n, x_{n+1}) + \alpha h S(x_n, x_n, x_{n+1}). \end{aligned}$$

This results in $\alpha(h+2) > 1$, which contradicts the initial assumption. Hence, our claim is proved.

Observe that by the assumption of the theorem, we have either

$$S(Tx_n, Tx_n, Tx) \leq g(S(x_n, x_n, x), S(Tx_n, x_n, x), S(Tx_n, x_n, x), S(Tx, x_n, x_n)),$$

or

$$\begin{aligned} S(Tx_{n+1}, Tx_{n+1}, Tx) &\leq g(S(x_{n+1}, x_{n+1}, x), S(Tx_{n+1}, x_{n+1}, x), \\ &\quad S(Tx_{n+1}, x_n, x), S(Tx, x_{n+1}, x_{n+1})). \end{aligned}$$

Therefore, one of the following cases holds.

Case (i). There exists an infinite subset $I \subseteq \mathbb{N}$ such that

$$\begin{aligned} S(x_{n+1}, x_{n+1}, Tx) &= S(Tx_n, Tx_n, Tx) \\ &\leq g(S(x_n, x_n, x), S(Tx_n, x_n, x), S(Tx_n, x_n, x), S(Tx, x_n, x_n)) \\ &= g(S(x_n, x_n, x), S(x_{n+1}, x_n, x), S(x_{n+1}, x_n, x), S(Tx, x_n, x_n)). \end{aligned}$$

for all $n \in I$.

Case (ii). There exists an infinite subset $J \subseteq \mathbb{N}$ such that

$$\begin{aligned} S(x_{n+2}, x_{n+2}, Tx) &= S(Tx_{n+1}, Tx_{n+1}, Tx) \\ &\leq g(S(x_{n+1}, x_{n+1}, x), S(Tx_{n+1}, x_{n+1}, x), \\ &\quad S(Tx_{n+1}, x_{n+1}, x), S(Tx, x_{n+1}, x_{n+1})) \\ &= g(S(x_{n+1}, x_{n+1}, x), S(x_{n+2}, x_{n+1}, x), \\ &\quad S(x_{n+2}, x_{n+1}, x), S(Tx, x_{n+1}, x_{n+1})). \end{aligned}$$

for all $n \in I$. In case (i), taking the limit as $n \rightarrow +\infty$ we obtain

$$S(x, x, Tx) \leq g(0, 0, 0, S(x, x, Tx))$$

Now by using Definition 5, Proposition 2, we have $S(x, x, Tx) = 0$, and thus $x = Tx$.

In case(ii), taking the limit as $n \rightarrow \infty$ we obtain

$$S(x, x, Tx) \leq g(0, 0, 0, S(x, x, Tx))$$

Now by using definition 5, propositions 2, we have $S(x, x, Tx) = 0$, and thus $x = Tx$. This completes the proof. \square

Corollary 2 *Let (X, S) be a S - metric space and $T : X \rightarrow X$ be a function. Suppose that there exist $g \in \mathcal{L}$ and $\alpha \in (0, 1)$, such that $\alpha(h + 2) \leq 1$ where $h = g(1, 1, 1, 1)$. Suppose also that $\alpha S(y, y, Ty) \leq S(x, y, z)$ implies*

$$S(Tx, Ty, Tz) \leq g(S(x, y, z), S(x, x, Tx), S(y, y, Ty), S(z, z, Tz))$$

for all $x, y, z \in X$. Then $F(T)$ is non-empty.

Corollary 3 *Let (X, S) be a complete S -metric space and $T : X \rightarrow X$ a function such that for all $x, y, z \in X$,*

$$S(Tx, Ty, Tz) \leq LS(x, y, z),$$

where $L \in (0, 1)$. Then there exists a unique point $u \in X$ such that $Tu = u$.

Proof. Let $g(x_1, x_2, x_3, x_4) = Lx_1$. □

Corollary 4 Let (X, S) be a complete S -metric space and $T : X \rightarrow X$ a function such that for all $x, y, z \in X$,

$$S(Tx, Ty, Tz) \leq L \max\{S(x, y, z), S(x, x, Tx), S(y, y, Ty), S(z, z, Tz)\}$$

where $L \in (0, 1)$. Then there exists a unique point $u \in X$ such that $Tu = u$.

Proof. Let $g(x_1, x_2, x_3, x_4) = L \max\{x_1, x_2, x_3, x_4\}$. □

References

- [1] S. Banach, Sur les operations dans les ensembles abstraits el leur appli-
cation aux equations integrals, *Fund. Math.*, **3** (1992), 133–181.
- [2] D. W. Boyd, S. W. Wong, On nonlinear contractions, *Proc. Am. Math.*
Soc., **20** (1969), 458–464.
- [3] M. Bukatin, R. Kopperman, S. Matthews, M. Pajoohesh, Partial metric
spaces, *Am. Math. Mon.*, **116** (2009), 708–718.
- [4] Lj. Ćirić, *Some recent results in metrical fixed point theory*, University of
Belgrade, Beograd 2003, Serbia.
- [5] B. C. Dhage, Generalized metric space and mapping with fixed point,
Bull. Calcutta. Math. Soc., **84** (1992), 329–336.
- [6] B. C. Dhage, Generalized metric space and topological structure I, *Analele*
Științifice ale Universității “Al. I. Cuza” din Iași. Serie Noua. Mathemat-
ica, **46** (1) (2000), 3–24.
- [7] T. Došenović, S. Radenović, S. Sedghi, Generalized metric spaces: Survey,
TWMS. J. Pure Appl. Math., **9** (1) (2018), 3–17.
- [8] J. Esfahani, Z. D. Mitrović, S. Radenović, S. Sedghi, Suzuki-type point
results in S -metric type spaces, *Comm. Appl. Nonlinear Anal.*, **25** (3)
(2018), 27–36.

- [9] S. Gahler, 2-metrische Raume und ihre topologische Struktur, *Math. Nachr.*, **26** (1963), 115–148.
- [10] L. G. Huang, X. Zhang, Cone metric spaces and fixed point theorems of contractive mappings, *J. Math. Anal. Appl.*, **332** (2012), 258–266.
- [11] J. K. Kim, S. Sedghi, N. Shobkolaei, Common Fixed point Theorems for the R-weakly commuting Mappings in S-metric spaces, *J. Comput. Anal. Appl.*, **19**(4) (2015), 751–759.
- [12] E. Malkowski, V. Rakočević, *Advanced Functional Analysis*, CRS Press, Taylor and Francis Group, Boca Raton, FL, 2019.
- [13] Z. Mustafa, B. Sims, A new approach to generalized metric spaces, *J. Nonlinear Convex Anal.*, **7** (2006), 289–297.
- [14] N. Y. Ozgur, N. Tas, Some Fixed Point Theorems on S-metric Spaces, *Mat. Vesnik*, **69** (1) (2017), 39–52.
- [15] M. M. Rezaee, M. Shahraki, S. Sedghi, I. Altun, Fixed Point Theorems For Weakly Contractive Mappings On S-Metric Spaces And a Homotopy Result, *Appl. Math. E-Notes*, **17** (2017), 1607–2510.
- [16] S. Sedghi, N. V. Dung, Fixed Point Theorems on S-Metric Spaces, *Mat. Vesnik*, **66** (1) (2014), 113–124.
- [17] S. Sedghi, N. Shobe, A. Aliouche, A generalization of fixed point theorems in S-metric spaces, *Mat. Vesnik*, **64** (2012), 258–266.
- [18] S. Sedghi, N. Shobe, T. Došenović, Fixed Point Results In S-Metric Spaces, *Nonlinear Funct. Anal. Appl.*, **20** (1) (2015), 55–67.
- [19] S. Sedghi, N. Shobe, M. Shahraki, T. Došenović, Common fixed Point of four maps In S-Metric Spaces, *Math. Sci.*, **12** (2018), 137–143.
- [20] S. Sedghi, N. Shobe, M. Zhou, A common fixed Point theorem in D^* -metric spaces, *Fixed point Theory Appl.*, (2007), Article ID27906.
- [21] S. Sedghi, A. Gholidahneh, T. Došenović, J. Esfahani, S. Radenović, Common fixed point of four maps in S_b -metric spaces, *J. Linear Topol. Algebra*, **05** (02) (2016), 93–104.

- [22] S. Sedghi, M. M. Rezaee, T. Došenović, S. Radenović, Common fixed point theorems for contractive mappings satisfying Φ -maps in S -metric spaces, *Acta Univ. Sapientiae, Mathematica*, **8** (2) (2016), 298–311.
- [23] V. Todorčević, *Harmonic Quasiconformal Mappings and Hyperbolic Type Metrics*, Springer Nature Switzerland AG 2019.

Received: April 15, 2020



On the metric dimension of strongly annihilating-ideal graphs of commutative rings

V. Soleymanivarniab

Department of Mathematics,
Science and Research Branch,
Islamic Azad University (IAU),
Tehran, Iran
email: soleymani.vali@yahoo.com

R. Nikandish*

Department of Mathematics,
Jundi-Shapur University of Technology,
Dezful, Iran
email: r.nikandish@ipm.ir

A. Tehranian

Department of Mathematics,
Science and Research Branch,
Islamic Azad University (IAU), Tehran, Iran
email: tehranian@srbiau.ac.ir

Abstract. Let R be a commutative ring with identity and $A(R)$ be the set of ideals with non-zero annihilator. The strongly annihilating-ideal graph of R is defined as the graph $SAG(R)$ with the vertex set $A(R)^* = A(R) \setminus \{0\}$ and two distinct vertices I and J are adjacent if and only if $I \cap \text{Ann}(J) \neq (0)$ and $J \cap \text{Ann}(I) \neq (0)$. In this paper, we study the metric dimension of $SAG(R)$ and some metric dimension formulae for strongly annihilating-ideal graphs are given.

2010 Mathematics Subject Classification: 13A99; 05C78; 05C12

Key words and phrases: metric dimension, strongly annihilating-ideal graph, commutative ring

*Corresponding author

1 Introduction

The problem of finding the metric dimension of a graph was first studied by Harary and Melter [7]. Determining the metric dimension of a graph as an NP-complete problem has attracted many graph theorists and it has appeared in various applications of graph theory, for example pharmaceutical chemistry [5], robot navigation [8], combinatorial optimization [14] and so on. Recently, there was much work done in computing the metric dimension of graphs associated with algebraic structures. Calculating the metric dimension for the commuting graph of a dihedral group was done in [1], for the zero-divisor graphs of commutative rings in [9, 10, 12], for the compressed zero-divisor graphs of commutative rings in [13], for total graphs of finite commutative rings in [6], for some graphs of modules in [11] and for annihilator graphs of commutative rings in [15]. Motivated by these papers, we study the metric dimension of another graph associated with a commutative ring.

Throughout this paper, all rings are assumed to be commutative with identity. The sets of all zero-divisors, nilpotent elements and maximal ideals are denoted by $Z(R)$, $Nil(R)$ and $Max(R)$, respectively. For a subset T of a ring R we let $T^* = T \setminus \{0\}$. An ideal with non-zero annihilator is called an *annihilating-ideal*. The set of annihilating-ideals of R is denoted by $A(R)$. For every subset I of R , we denote the *annihilator* of I by $Ann(I)$. A non-zero ideal I of R is called *essential* if I has a non-zero intersection with every other non-zero ideal of R . The set of essential annihilating-ideal ideals of R is denoted by $Ess(R)$. The ring R is said to be *reduced* if it has no non-zero nilpotent element. Some more definitions about commutative rings can be found in [2, 4].

We use the standard terminology of graphs following [18]. Let $G = (V, E)$ be a graph, where $V = V(G)$ is the set of vertices and $E = E(G)$ is the set of edges. We recall that a graph is *connected* if there exists a path connecting any two distinct vertices. The *distance* between two distinct vertices x and y , denoted by $d(x, y)$, is the length of the shortest path connecting them (if such a path does not exist, then we set $d(x, y) = \infty$). The *diameter* of a connected graph G , denoted by $diam(G)$, is the maximum distance between any pair of vertices of G . For a vertex x in G , we denote the set of all vertices adjacent to x by $N(x)$ and $N[x] = N(x) \cup \{x\}$. A *k-partite* graph is one whose vertex set can be partitioned into k subsets so that an edge has both ends in no subset. A *complete k-partite* graph is a k -partite graph in which each vertex is adjacent to every vertex that is not in the same subset. The complete bipartite (i.e., 2-partite) graph with part sizes m and n is denoted by $K_{m,n}$. If $m = 1$, then the bipartite graph is called star. A graph in which each

pair of vertices is joined by an edge is called a *complete* graph and use K_n to denote it with n vertices and its complement is denoted by \bar{K}_n (possibly n is zero). Also, a cycle of order n is denoted by C_n . A subset of vertices $S \subseteq V(G)$ resolves a graph G , and S is a *resolving set* of G , if every vertex is uniquely determined by its vector of distances to the vertices of S . In general, for an ordered subset $S = \{v_1, v_2, \dots, v_k\}$ of vertices in a connected graph G and a vertex $v \in V(G) \setminus S$ of G , the metric representation of v with respect to S is the k -vector $D(v|S) = (d(v, v_1), d(v, v_2), \dots, d(v, v_k))$. The set S is a resolving set for G if $D(u|S) = D(v|S)$ implies that $u = v$, for all pair of vertices, $v, u \in V(G) \setminus S$. A resolving set S of minimum cardinality is the *metric basis* for G , and the number of elements in the resolving set of minimum cardinality is the *metric dimension* of G . We denote the metric dimension of a graph G by $\dim_M(G)$. Let G be a connected graph such that $|V(G)| \geq 2$. Two distinct vertices u and v are *distance similar*, if $d(u, x) = d(v, x)$, for all $x \in V(G) \setminus \{u, v\}$. It can be easily checked that two distinct vertices u and v are distance similar if either $u - v \notin E(G)$ and $N(u) = N(v)$ or $u - v \in E(G)$ and $N[u] = N[v]$.

Let R be a commutative ring with identity and $A(R)$ be the set of ideals with non-zero annihilator. The *strongly annihilating-ideal graph* of R is defined as the graph $SAG(R)$ with the vertex set $A(R)^* = A(R) \setminus \{0\}$ and two distinct vertices I and J are adjacent if and only if $I \cap \text{Ann}(J) \neq (0)$ and $J \cap \text{Ann}(I) \neq (0)$. This graph was first introduced and studied in [16, 17]. It is worthy to mention that strongly annihilating-ideal graph is a generalization of annihilating-ideal graph. The *annihilating-ideal graph* of R , denoted by $AG(R)$, is a graph with the vertex set $A(R)^*$ and two distinct vertices I and J are adjacent if and only if $IJ = 0$ (see [3] for more details). In this paper, we study the metric dimension of $SAG(R)$ and we provide some metric dimension formulas for $SAG(R)$.

2 Metric dimension of a strongly annihilating-ideal graph of a reduced ring

Let R be a commutative ring. In this section, we provide a metric dimension formula for a strongly annihilating-ideal graph when R is reduced.

Lemma 1 *Let R be a ring which is not an integral domain. Then $\dim_M(SAG(R))$ is finite if and only if R has only finitely many ideals.*

Proof. One side is clear. To prove the other side, suppose that $\dim_M(SAG(R))$ is finite and let $W = \{I_1, I_2, \dots, I_n\}$ be the metric basis for $SAG(R)$, where n

is a non-negative. By [16, Theorem 2.1], $\text{diam}(\text{SAG}(\mathbf{R})) \leq 2$ and so for every $I \in \mathbf{A}(\mathbf{R})^* \setminus W$, there are $(2+1)^n$ possibilities for $D(I|W)$. Thus $|\mathbf{A}(\mathbf{R})^*| \leq 3^n + n$ and hence \mathbf{R} has only finitely many ideals. \square

If \mathbf{R} is a reduced ring with finitely many ideals, then by [2, Theorem 8.7], \mathbf{R} is a direct product of finitely many fields. Using this fact, we prove the following result.


Theorem 1 *Let \mathbf{R} be a reduced ring which is not an integral domain. If $\dim_{\mathbf{M}}(\text{SAG}(\mathbf{R}))$ is finite, then:*

- (1) *If $|\text{Max}(\mathbf{R})| \leq 3$, then $\dim_{\mathbf{M}}(\text{SAG}(\mathbf{R})) = |\text{Max}(\mathbf{R})| - 1$.*
- (2) *If $|\text{Max}(\mathbf{R})| \geq 4$, then $\dim_{\mathbf{M}}(\text{SAG}(\mathbf{R})) = |\text{Max}(\mathbf{R})|$.*

Proof. (1) Since $\dim_{\mathbf{M}}(\text{SAG}(\mathbf{R}))$ is finite, \mathbf{R} has only finitely many ideals, by Lemma 1. Also, since \mathbf{R} is not an integral domain, $|\text{Max}(\mathbf{R})| \neq 1$. Hence $|\text{Max}(\mathbf{R})| = 2$ or 3 . If $|\text{Max}(\mathbf{R})| = 2$, then $\mathbf{R} \cong F_1 \times F_2$, where F_i is a field. Thus $\text{SAG}(\mathbf{R}) = K_2$ and so $\dim_{\mathbf{M}}(\text{SAG}(\mathbf{R})) = 1$. If $|\text{Max}(\mathbf{R})| = 3$, then $\mathbf{R} \cong F_1 \times F_2 \times F_3$, where F_i is a field for every $1 \leq i \leq 3$. Let $W = \{F_1 \times (0) \times F_3, F_1 \times F_2 \times (0)\}$. By the following figure, one may easily get

$$\begin{aligned} D((0) \times F_2 \times (0)|W) &= (1, 2), \\ D(F_1 \times (0) \times (0)|W) &= (2, 2), \\ D((0) \times (0) \times F_3|W) &= (2, 1), \\ D((0) \times F_2 \times F_3|W) &= (1, 1). \end{aligned}$$

So for every $x, y \in V(\text{SAG}(\mathbf{R})) \setminus W$, $D(x|W) \neq D(y|W)$ and hence $\dim_{\mathbf{M}}(\text{SAG}(\mathbf{R})) = 2$.



- (2) Assume that $|\text{Max}(\mathbf{R})| = n \geq 4$. By Lemma 1, $\mathbf{R} \cong F_1 \times \cdots \times F_n$, where

F_i is a field for every $1 \leq i \leq n$. We show that $\dim_M(\text{SAG}(\mathbf{R})) = n$. Indeed, we have the following claims:

Claim 1. $\dim_M(\text{SAG}(\mathbf{R})) \geq n$.

Since $\mathbf{R} \cong F_1 \times \cdots \times F_n$, by Lemma 1, $\dim_M(\text{SAG}(\mathbf{R}))$ is finite. Let $W = \{I_1, I_2, \dots, I_k\}$ be the metric basis for $\text{SAG}(\mathbf{R})$, where k is a positive integer. On the other hand, by [16, Theorem 2.1], $\text{diam}(\text{SAG}(\mathbf{R})) \in \{1, 2\}$, and so for every $I \in A(\mathbf{R})^* \setminus W$, there are 2^k possibilities for $D(I|W)$. This implies that $|A(\mathbf{R})^*| - k \leq 2^k$. Since $|A(\mathbf{R})^*| = 2^n - 2$, $2^n - 2 - k \leq 2^k$ and hence $2^n \leq 2^k + 2 + k$. Since $n \geq 4$, we deduce that $k \geq n$. Therefore $\dim_M(\text{SAG}(\mathbf{R})) \geq n$.

Claim 2. $\dim_M(\text{SAG}(\mathbf{R})) \leq n$.

For every $1 \leq i \leq n$, let $(F_1, \dots, F_{i-1}, 0, F_{i+1}, \dots, F_n) = \mathbf{m}_i \in A(\mathbf{R})^*$. Put $W = \{\mathbf{m}_1, \mathbf{m}_2, \dots, \mathbf{m}_n\}$ (in fact $W = \text{Max}(\mathbf{R})$). We show that W is the resolving set for $\text{SAG}(\mathbf{R})$. To see this, let $I, J \in V(\text{SAG}(\mathbf{R})) \setminus W$ and $I \neq J$. We need only to show that $D(I|W) \neq D(J|W)$. Let $I = (I_1, I_2, \dots, I_n)$ and $J = (J_1, J_2, \dots, J_n)$. Since $I \neq J$, $I_i = 0$ and $J_i = F_i$ or $I_i = F_i$ and $J_i = 0$, for some $1 \leq i \leq n$. Without loss of generality, assume that $I_1 = 0$ and $J_1 = F_1$. It is easy to see that $d(I, \mathbf{m}_1) = 1$ and $d(J, \mathbf{m}_1) = 2$. This clearly shows that $D(I|W) \neq D(J|W)$. Therefore $\dim_M(\text{SAG}(\mathbf{R})) \leq n$.

Now, by Claims 1, 2, $\dim_M(\text{SAG}(\mathbf{R})) = n$, for $n \geq 4$. □

3 Metric dimension of a strongly annihilating-ideal graph of a non-reduced ring

In this section, we discuss the metric dimension of strongly annihilating-ideal graphs for non-reduced rings. First we need to recall two lemmas from [16].

Lemma 2 [16, Lemma 2.1] *Let \mathbf{R} be a ring and $I, J \in A(\mathbf{R})^*$. Then the following statements hold.*

- (1) *If $I - J$ is not an edge of $\text{SAG}(\mathbf{R})$, then $\text{Ann}(IJ) = \text{Ann}(I)$ or $\text{Ann}(IJ) = \text{Ann}(J)$. Moreover, if \mathbf{R} is a reduced ring, then the converse is also true.*
- (2) *If $I - J$ is an edge of $\mathbb{A}\mathbb{G}(\mathbf{R})$, then $I - J$ is an edge of $\text{SAG}(\mathbf{R})$.*
- (3) *If $\text{Ann}(I) \not\subseteq \text{Ann}(J)$ and $\text{Ann}(J) \not\subseteq \text{Ann}(I)$, then $I - J$ is an edge of $\text{SAG}(\mathbf{R})$. Moreover if \mathbf{R} is a reduced ring, then the converse is also true.*

- (4) Let $n \geq 1$ be a positive integer. Suppose that $R \cong R_1 \times \cdots \times R_n$, where R_i is a ring, for every $1 \leq i \leq n$, and $I = (I_1, \dots, I_n)$ and $J = (J_1, \dots, J_n)$ are two vertices of $\text{SAG}(R)$. If $I_i \cap \text{Ann}(J_i) \neq (0)$ and $J_j \cap \text{Ann}(I_j) \neq (0)$, for some $1 \leq i, j \leq n$, then $I - J$ is an edge of $\text{SAG}(R)$. In particular, if $I_i - J_i$ is an edge of $\text{SAG}(R_i)$ or $I_i = J_i$ and $I_i \cap \text{Ann}(I_i) \neq (0)$, for some $1 \leq i \leq n$, then $I - J$ is an edge of $\text{SAG}(R)$.
- (5) If $I, J \in \text{Ess}(R)$ or $\text{Ann}(I), \text{Ann}(J) \in \text{Ess}(R)$, then I is adjacent to J .
- (6) If $d_{\text{AG}(R)}(I, J) = 3$ for some distinct $I, J \in A(R)^*$, then $I - J$ is an edge of $\text{SAG}(R)$.
- (7) If $I - J$ is not an edge of $\text{SAG}(R)$ for some distinct $I, J \in A(R)^*$, then $d_{\text{AG}(R)}(I, J) = 2$.

Lemma 3 [16, Lemma 2.2] *Let R be a non-reduced ring and I be an ideal of R such that $I^n = (0)$, for some positive integer n . Then $\text{Ann}(I)$ is an essential ideal of R .*

Remark 1 *Let G be a connected graph and V_1, V_2, \dots, V_k be a partition of $V(G)$ such that for every $1 \leq i \leq k$, if $x, y \in V_i$, then $N(x) = N(y)$. Then $\dim_M(G) \geq |V(G)| - k$.*

Next, we provide some formulas for the metric dimension of strongly annihilating-ideal graphs for non-reduced rings.

Theorem 2 *Suppose that $R \cong R_1 \times \cdots \times R_n$, where R_i is an Artinian local ring such that for every $1 \leq i \leq n$, $|A(R_i)^*| = 1$. Then $\dim_M(\text{SAG}(R)) = 2n$.*

Proof. Assume that $X = (R_1, 0, \dots, 0)$ and $Y = (I_1, 0, \dots, 0)$, where $I_1 \in A(R_1)^*$. By Part 4 of Lemma 2, it is easy to see that $N(X) = N(Y)$. This implies that if W is the metric basis for $\text{SAG}(R)$, then $X \in W$ or $Y \in W$. Without loss of generality, we may assume that $X \in W$. Similarly, we may assume that $W_1 \subseteq W$, where $W_1 = \{(R_1, 0, \dots, 0), (0, R_2, 0, \dots, 0), \dots, (0, \dots, 0, R_n)\}$.

Now, assume that $X = (0, R_2, \dots, R_n)$ and $Y = (I_1, R_2, \dots, R_n)$, where $I_1 \in A(R_1)^*$. It is easy to see that $N(X) = N(Y)$ and so if W is the metric basis for $\text{SAG}(R)$, then $X \in W$ or $Y \in W$. Without loss of generality, we may assume that $X \in W$. Similarly, we may assume that $W_2 \subseteq W$, where

$$W_2 = \{(0, R_2, \dots, R_n), (R_1, 0, R_3, \dots, R_n), \dots, (R_1, \dots, R_{n-1}, 0)\}.$$

Since $|W_1| = |W_2| = n$ and $W_1 \cup W_2 \subseteq W$, $|W| \geq 2n$. We show that $|W| \leq 2n$. For this, it is enough to show that W is a resolving set and consequently it is

the metric basis for the graph $\text{SAG}(\mathbf{R})$. Let $X, Y \notin W$, $X \neq Y$, $X = (I_1, \dots, I_n)$ and $Y = (J_1, \dots, J_n)$. We show that $D(X|W) \neq D(Y|W)$. Since $X \neq Y$, for some $1 \leq i \leq n$, we conclude that $I_i \neq J_i$. Without loss of generality, one may assume that $I_1 \supset J_1$. We have the following cases:

Case 1. $I_1 = R_1$.

Subcase 1. For some $2 \leq j \leq n$, $J_j \neq 0$. In this case, $Z - Y$ is an edge of $\text{SAG}(\mathbf{R})$ but $Z - X$ is not an edge of $\text{SAG}(\mathbf{R})$, where $Z = (R_1, 0, \dots, 0)$. Since $Z \in W$, we deduce that $D(X|W) \neq D(Y|W)$.

Subcase 2. For every $2 \leq j \leq n$, $J_j = 0$. Since $I_1 = R_1$ and $(R_1, 0, \dots, 0) \in W$, for some $2 \leq i \leq n$, $I_i \neq 0$. If $I_i = R_i$, for some $2 \leq i \leq n$, then $Z - Y$ is an edge of $\text{SAG}(\mathbf{R})$ but $Z - X$ is not an edge of $\text{SAG}(\mathbf{R})$, where $Z = (0, \dots, 0, R_i, 0, \dots, 0)$. So we can let for every $2 \leq i \leq n$, $I_i \neq R_i$. Now, without loss of generality, we may assume that $I_2 \neq 0$. Obviously, $Z - X$ is an edge of $\text{SAG}(\mathbf{R})$ but $Z - Y$ is not an edge of $\text{SAG}(\mathbf{R})$, where $Z = (R_1, 0, R_3, \dots, R_n)$. Since $Z \in W$, $D(X|W) \neq D(Y|W)$.

Case 2. $I_1 \neq R_1$. Since $I_1 \neq R_1$, $J_1 \neq R_1$. Also, since $X \neq Y$, we may let $I_1 \in A(R_1)^*$ and $J_1 = 0$. If $I_i \neq R_i$, for some $2 \leq i \leq n$, then $Z - X$ is an edge of $\text{SAG}(\mathbf{R})$ but $Z - Y$ is not an edge of $\text{SAG}(\mathbf{R})$, where $Z = (0, R_2, R_3, \dots, R_n)$. Since $Z \in W$, $D(X|W) \neq D(Y|W)$. So let $X = (I_1, R_2, \dots, R_n)$. Since $J_1 = 0$ and $Y \notin W$, for some $2 \leq i \leq n$, $J_i \in A(R_1)^*$. Without loss of generality, we may assume that $J_2 \in A(R_2)^*$. If $J_i \neq 0$, for some $3 \leq i \leq n$, then we put $Z = (0, R_2, \dots, R_{i-1}, 0, R_{i+1}, \dots, R_n)$. It is not hard to check that $Z - Y$ is an edge of $\text{SAG}(\mathbf{R})$ but $Z - X$ is not an edge of $\text{SAG}(\mathbf{R})$. If for every $3 \leq i \leq n$, $J_i = 0$, then we put $Z = (R_1, R_2, \dots, 0, \dots, 0)$. In both cases we have that $D(X|W) \neq D(Y|W)$. Therefore, $|W| \leq 2n$. \square

Theorem 3 Suppose that $\mathbf{R} \cong R_1 \times \dots \times R_n$, where R_i is an Artinian local ring such that for every $1 \leq i \leq n$, $|A(R_i)^*| \geq 2$. Then $\dim_M(\text{SAG}(\mathbf{R})) = |A(\mathbf{R})^*| - 3^n + 2$.

Proof. If \mathbf{R} is local, then Lemma 3 implies that $\text{SAG}(\mathbf{R})$ is complete and hence $\dim_M(\text{SAG}(\mathbf{R})) = |A(\mathbf{R})^*| - 1$. So let $\mathbf{R} \cong R_1 \times \dots \times R_n$ and $n \geq 2$. Assume that

$X = (I_1, \dots, I_n)$, $Y = (J_1, \dots, J_n)$ are vertices of $\text{SAG}(\mathbf{R})$. Define the relation \sim on $V(\text{SAG}(\mathbf{R}))$ as follows: $X \sim Y$, whenever, the following two conditions hold.

- (1) " $I_i = 0$ if and only if $J_i = 0$ " for every $1 \leq i \leq n$.

(2) " $0 \neq I_i \subseteq \text{Nil}(R_i)$ if and only if $0 \neq J_i \subseteq \text{Nil}(R_i)$ " for every $1 \leq i \leq n$.

It is easily seen that \sim is an equivalence relation on $V(\text{SAG}(R))$. By $[X]$, we mean the equivalence class of X . Let X_1 and X_2 be two elements of $[X]$. Since $X_1 \sim X_2$, by Part 4 of Lemma 2, $N(X_1) = N(X_2)$. This, together with the fact that the number of equivalence classes is $3^n - 2$ and Remark 1, implies that

$$\dim_M(\text{SAG}(R)) \geq |A(R)^*| - (3^n - 2) = |A(R)^*| - 3^n + 2.$$

We show that

$$\dim_M(\text{SAG}(R)) \leq |A(R)^*| - 3^n + 2.$$

Let

$A = \{(I_1, \dots, I_n) \in V(\text{SAG}(R)) \mid I_i \in \{0, \text{Nil}(R_i), \dots, R_i\} \text{ for every } 1 \leq i \leq n\}$ and $W = A(R)^* \setminus A$.

It is shown that W is a resolving set and consequently it is the metric basis for the graph $\text{SAG}(R)$. To see this, let $X, Y \in A$ and $X \neq Y$. We show that $D(X|W) \neq D(Y|W)$. Let $X = (I_1, \dots, I_n)$ and $Y = (J_1, \dots, J_n)$. Since $X \neq Y$, for some $1 \leq i \leq n$, $I_i \neq J_i$. Without loss of generality, we may assume that $I_1 \supset J_1$. We have the following cases:

Case 1. $I_1 = R_1$.

Subcase 1. $J_1 = 0$. In this case $Z - X$ is an edge of $\text{SAG}(R)$ but $Z - Y$ is not an edge of $\text{SAG}(R)$, where $Z = (I'_1, R_2, \dots, R_n)$ and $I'_1 \in A(R_1)^* \setminus \{\text{Nil}(R_1)\}$. Since $Z \in W$, $D(X|W) \neq D(Y|W)$. **Subcase 2.** $J_1 = \text{Nil}(R_1)$. In this case $Z - Y$ is an edge of $\text{SAG}(R)$ but $Z - X$ is not an edge of $\text{SAG}(R)$, where $Z = (J'_1, 0, \dots, 0)$, $J'_1 \in A(R_1)^*$ and $J'_1 \neq \text{Nil}(R_1)$. Since $Z \in W$, $D(X|W) \neq D(Y|W)$.

Case 2. $I_1 = \text{Nil}(R_1)$.

Since $I_1 \neq J_1$ and $I_1 \supseteq J_1$, $J_1 = 0$. Hence $Z - X$ is an edge of $\text{SAG}(R)$ but $Z - Y$ is not an edge of $\text{SAG}(R)$, where $Z = (J'_1, R_2, \dots, R_n)$ and $J'_1 \in A(R_1)^*$ and $J'_1 \neq \text{Nil}(R_1)$. Since $Z \in W$, $D(X|W) \neq D(Y|W)$. Therefore,

$$\dim_M(\text{SAG}(R)) \leq |W|.$$

Since $|A| = 3^n - 2$, $|W| = |A(R)^*| - (3^n - 2) = |A(R)^*| - 3^n + 2$. Therefore,

$$\dim_M(\text{SAG}(R)) \leq |A(R)^*| - 3^n + 2.$$

□

Next, we provide some upper and lower bounds for the metric dimension of strongly annihilating-ideal graphs for some other classes of non-reduced rings.

Theorem 4 Suppose that $R \cong R_1 \times \cdots \times R_n \times F_{n+1} \times \cdots \times F_{n+m}$, where R_i is an Artinian local ring such that $|A(R_i)| = 2$ for every $1 \leq i \leq n$ and F_i is a field for every $1+n \leq i \leq n+m$. Then $n+m \leq \dim_M(\text{SAG}(R)) \leq 2^{n+m} - 2$.

Proof. Suppose that $W = \{I_1, I_2, \dots, I_k\}$ be the metric basis for $\text{SAG}(R)$, for some non-negative integer k . Since $\text{diam}(\text{SAG}(R)) \leq 2$, there are exactly $(2)^k$ possibilities for $D(I|W)$, for every $I \in A(R)^* \setminus W$. On the other hand, since $|A(R)^*| = 3^{n+2m} - 2$, we must have $3^{n+2m} - 2 - k \leq 2^k$. This implies that $n+m \leq k$. Hence $n+m \leq \dim_M(\text{SAG}(R))$. It is shown that $\dim_M(\text{SAG}(R)) \leq 2^{n+m} - 2$. Let

$W = \{(I_1, \dots, I_{n+m}) \in V(\text{SAG}(R)) \mid I_i \in \{0, R_1, \dots, R_n, F_1, \dots, F_m\} \text{ for every } 1 \leq i \leq n+m\}$.

We show that W is a resolving set for $\text{SAG}(R)$. For this, let $X, Y \in A(R)^* \setminus W$ and $X \neq Y$. We show that $D(X|W) \neq D(Y|W)$. Let $X = (I_1, \dots, I_{n+m})$ and $Y = (J_1, \dots, J_{n+m})$. Since $X \neq Y$, $I_i \neq J_i$, for some $1 \leq i \leq n+m$.

We have the following cases:

Case 1. For some $n+1 \leq i \leq n+m$, $I_i \neq J_i$.

Without loss of generality, we may assume that $i = n+m$, $I_{n+m} = F_{n+m}$ and $J_{n+m} = 0$. Now, put $Z = (R_1, \dots, R_n, F_{n+1}, \dots, F_{n+m-1}, 0)$. Since for some $1 \leq i \leq n$, $I_i \in A(R_i)^*$, one may easily see that $Z - X$ is an edge of $\text{SAG}(R)$ but $Z - Y$ is not an edge of $\text{SAG}(R)$. Since $Z \in W$, $D(X|W) \neq D(Y|W)$.

Case 2. For every $n+1 \leq i \leq n+m$, $I_i = J_i$.

Since $I_i \neq J_i$, for some $1 \leq i \leq n$, one can let $J_1 \subset I_1$. Thus we have the following subcases:

Subcase 1. $J_1 = 0$ and $I_1 \in A(R_1)^*$.

Since $J_1 = 0$, for some $2 \leq i \leq n$, $J_i \in A(R_i)^*$. Hence one can let $J_2 \in A(R_2)^*$. If for some $2 \leq i \leq n$, $I_i \neq R_i$ or for some $1+m \leq i \leq n+m$, $I_i \neq F_i$, then put $Z = (0, R_2, R_3, \dots, R_n, F_{n+1}, \dots, F_{n+m})$. $Z - X$ is an edge of $\text{SAG}(R)$ but $Z - Y$ is not an edge of $\text{SAG}(R)$. Since $Z \in W$, $D(X|W) \neq D(Y|W)$. So we let $X = (I_1, R_2, \dots, R_n, F_{n+1}, \dots, F_{n+m})$. Similarly, if for some $3 \leq i \leq n$, $J_i \neq R_i$ or for some $1+m \leq j \leq n+m$, $J_i \neq F_i$, then without loss of generality, we may assume that $J_3 \neq R_3$. Then put $Z = (0, 0, R_3, \dots, R_n, F_{n+1}, \dots, F_{n+m})$. Thus $Z - Y$ is an edge of $\text{SAG}(R)$ but $Z - X$ is not an edge of $\text{SAG}(R)$. Since $Z \in W$, $D(X|W) \neq D(Y|W)$. Now, let $X = (I_1, R_2, \dots, R_n, F_{n+1}, \dots, F_{n+m})$ and $Y = (0, J_2, R_3, \dots, R_n, F_{n+1}, \dots, F_{n+m})$. Put $Z = (0, R_2, 0, \dots, 0, 0, \dots, 0)$. Therefore, $Z - Y$ is an edge of $\text{SAG}(R)$ but $Z - X$ is not an edge of $\text{SAG}(R)$. Since $Z \in W$, $D(X|W) \neq D(Y|W)$.

Subcase 2. $J_1 = 0$ and $I_1 = R_1$.

Since $J_1 = 0$, for some $2 \leq i \leq n$, $J_i \in A(R_i)^*$. Hence one may let $J_2 \in A(R_2)^*$. Assume that $Z = (R_1, 0, \dots, 0)$. Thus $Z - Y$ is an edge of $SAG(R)$ but $Z - X$ is not an edge of $SAG(R)$ (note that since $Z \in W$, $Z \neq X$). This implies that $D(X|W) \neq D(Y|W)$.

Subcase 3. $J_1 \in A(R_1)^*$ and $I_1 = R_1$. If $J_i \neq 0$, for some $2 \leq i \leq n$, then one

may assume that $J_2 \neq 0$. Suppose that $Z = (R_1, 0, \dots, 0)$. Then $Z - Y$ is an edge of $SAG(R)$ but $Z - X$ is not an edge of $SAG(R)$. Hence $D(X|W) \neq D(Y|W)$. Let $Y = (J_1, 0, \dots, 0)$. Since $X \notin W$, for some $2 \leq i \leq n$, $I_i \in A(R_i)^*$. So, we can let $I_2 \in A(R_2)^*$. If $I_i \neq 0$, for some $3 \leq i \leq n$, then we can assume that $I_3 \neq 0$. If we put $Z = (R_1, R_2, 0, \dots, 0)$, then we easily get $D(X|W) \neq D(Y|W)$. Finally, if $X = (R_1, I_2, 0, \dots, 0)$ and $Y = (J_1, 0, \dots, 0)$, then $D(X|W) \neq D(Y|W)$. Since $Z - X$ is an edge of $SAG(R)$ but $Z - Y$ is not an edge of $SAG(R)$, where $Z = (R_1, 0, R_3, 0, \dots, 0)$. Therefore, $\dim_M(SAG(R)) \leq |W|$. Since $|W| = 2^{n+m} - 2$, $\dim_M(SAG(R)) \leq 2^{n+m} - 2$. \square

We end this paper with the following example.

Example 1 (1) Let $R = \mathbb{Z}_4 \times \mathbb{Z}_2$. Then $SAG(R) = C_4$ and hence $\dim_M(SAG(R)) = 2$. Also, in Theorem 4, $n = m = 1$, and so $\dim_M(SAG(R)) = 2$.

(2) Let $R = \mathbb{Z}_4 \times \mathbb{Z}_2 \times \mathbb{Z}_2$ and $\dim_M(SAG(R)) = k$. We show that $3 \leq k \leq 6$. Since $\text{diam}(SAG(R)) \leq 2$ and $|A(R)^*| = 10$, $10 - k \leq 2^k$. Thus $k \geq 3$. Let $W = \{((2), \mathbb{Z}_2, \mathbb{Z}_2), ((2), 0, \mathbb{Z}_2), ((2), \mathbb{Z}_2, 0), ((2), 0, 0)\}$. Then

$$\begin{aligned} D((\mathbb{Z}_4, 0, 0)|W) &= (1, 1, 1, 2), \\ D((\mathbb{Z}_4, \mathbb{Z}_2, 0)|W) &= (1, 1, 2, 2), \\ D((\mathbb{Z}_4, 0, \mathbb{Z}_2)|W) &= (1, 2, 1, 2), \\ D((0, \mathbb{Z}_2, \mathbb{Z}_2)|W) &= (2, 1, 1, 1), \\ D((0, \mathbb{Z}_2, 0)|W) &= (2, 1, 2, 1), \\ D((0, 0, \mathbb{Z}_2)|W) &= (2, 2, 1, 1). \end{aligned}$$

Therefore, W is a resolving set for $SAG(R)$ and hence $k \leq 6$.

Acknowledgements

The authors express their deep gratitude to the referees for their valuable suggestions which have definitely improved the paper.

References

- [1] F. Ali, M. Salman, S. Huang, On the commuting graph of dihedral group, *Comm. Algebra.*, **44** (2016), 2389–2401.
- [2] M. F. Atiyah, I. G. Macdonald, *Introduction to Commutative Algebra*, Addison-Wesley Publishing Company, (1969).
- [3] M. Behboodi, The annihilating-ideal graph of a commutative ring I, *J. Algebra Appl.*, **10** (2011), 727–739.
- [4] W. Bruns and J. Herzog, *Cohen-Macaulay Rings*, Cambridge University Press (1997).
- [5] G. Chartrand, L. Eroh, M. A. Johnson, O. R. Oellermann, Resolvability in graphs and the metric dimension of a graph, *Disc. Appl. Math.*, **105** (2000), 99–113.
- [6] D. Dolžan, The metric dimension of the total graph of a finite commutative ring, *Canad. Math. Bull.*, **59** (2016), 748–759.
- [7] F. Harary and R. A. Melter, On the metric dimension of a graph, *Ars Combin.*, **2** (1976), 191–195.
- [8] S. Khuller, B. Raghavachari, A. Rosenfeld, *Localization in graphs*, Technical report CS-TR-3326, University of Maryland at College Park, 1994.
- [9] S. Pirzada, R. Raja and S. P. Redmond, Locating sets and numbers of graphs associated to commutative rings, *J. Algebra Appl.* **13:7** (2014): 1450047 18 pp.
- [10] S. Pirzada, R. Raja, On the metric dimension of a zero-divisor graph, *Communications in Algebra*, **45:4** (2017), 1399–1408.
- [11] S. Pirzada, Rameez Raja, On graphs associated with modules over commutative rings, *J. Korean. Math. Soc.*, **53** (2016), 1167–1182.
- [12] R. Raja, S. Pirzada and S. P. Redmond, On Locating numbers and codes of zero-divisor graphs associated with commutative rings, *J. Algebra Appl.*, **15:1** (2016): 1650014 22 pp.
- [13] S. Pirzada, M. Imran Bhat, Computing metric dimension of compressed zero divisor graphs associated to rings, *Acta Univ. Sapientiae, Mathematica*, **10** (2) (2018), 298–318.

- [14] A. Sebö, E. Tannier, On metric generators of graphs, *Math. Oper. Res.*, **29** (2004), 383–393.
- [15] V. Soleymanivarniab, A. Tehranian, R. Nikandish, The metric dimension of annihilator graphs of commutative rings, *J. Algebra Appl.*, to appear.
- [16] N. KH. Tohidi, M. J. Nikmehr, R. Nikandish, On the strongly annihilating-ideal graph of a commutative ring, *Discrete Math. Algorithm. Appl.*, 09, 1750028 (2017) [13 pages].
- [17] N. KH. Tohidi, M. J. Nikmehr, R. Nikandish, Some results on the strongly annihilating-ideal graph of a commutative ring, *Bol. Soc. Mat. Mex.*, **24**, (2018), 307–318.
- [18] D. B. West, *Introduction to Graph Theory*, 2nd ed., Prentice Hall, Upper Saddle River (2001).

Received: February 23, 2020



On application of differential subordination for Carathéodory functions

V. Agnes Sagaya Judy Lavanya

Dr. MGR Janaki College of Arts and Science
Chennai 600028, TamilNadu, India
email: lavanyaravi06@gmail.com

M. P. Jeyaraman

L. N. Government College
Ponneri, Chennai 601 204,
Tamil Nadu, India
email: jeyaraman_mp@yahoo.co.in

H. Aaisha Farzana

A. M. Jain College
Meenambakkam, Chennai 600 114,
Tamil Nadu, India
email: h.aaisha@gmail.com

Abstract. New sufficient conditions involving the properties of analytic functions to belong to the class of Carathéodory functions are investigated. Certain univalence and starlikeness conditions are deduced as special cases of main results.

1 Introduction

Let \mathcal{H} be the class of analytic functions in the open unit disk $\mathbb{D} := \{z \in \mathbb{C} : |z| < 1\}$. Let \mathcal{A} denote the class of all the functions $f \in \mathcal{H}$ that satisfy the normalization $f(0) = 0$, $f'(0) = 1$. Let \mathcal{S} denote the subclass of \mathcal{A} consisting of univalent functions. The function $f \in \mathcal{A}$ satisfying the conditions $\operatorname{Re}\{zf'(z)/f(z)\} > 0$, $\operatorname{Re}\{1 + zf''(z)/f'(z)\} > 0$ belong to the familiar classes of starlike and convex functions denoted by \mathcal{S}^* and \mathcal{C} respectively. Let f and g be analytic in \mathbb{D} , then we say that f is subordinate to g in \mathbb{D} (written $f \prec g$)

2010 Mathematics Subject Classification: 30C45, 30C80

Key words and phrases: analytic functions, differential subordination, Carathéodory function, starlike function, univalent function

if there exists a Schwarz function $w(z)$, analytic in \mathbb{D} with $w(0) = 0$ such that $f(z) = g(w(z))$, ($z \in \mathbb{D}$). In particular, if the function g is univalent in \mathbb{D} , then the subordination is equivalent to $f(0) = g(0)$ or $f(\mathbb{D}) \subset g(\mathbb{D})$. Let us denote by \mathcal{Q} the set of functions q that are analytic and injective on $\bar{\mathbb{D}} \setminus E(q)$, where $E(q) = \{\zeta \in \partial\mathbb{D} : \lim_{z \rightarrow \zeta} q(z) = \infty\}$, and are such that $q'(\zeta) \neq 0$ for $\zeta \in \partial\mathbb{D} \setminus E(q)$. Further, the subclass of \mathcal{Q} for which $q(0) = a$ be denoted by $\mathcal{Q}(a)$. Let $\mathcal{P}(\alpha)$ be a class of functions of the form $p(z) = 1 + \sum_{n=1}^{\infty} p_n z^n$, which are analytic in \mathbb{D} , we say that $p(z) \in \mathcal{P}(\alpha)$ if $\operatorname{Re}\{p(z)\} > \alpha$. We note that for $\mathcal{P}(0) := \mathcal{P}$ is the class of Carathéodory functions in \mathbb{D} .

The function $q_c(z) = \sqrt{1 + cz}$, maps \mathbb{D} onto a set which is bounded by the lemniscate of Bernoulli. That is, $q_c(\mathbb{D}) = \{w \in \mathbb{C} : |w^2 - 1| < c\}$, and the class $\mathcal{S}^*(q_c)$ given by $\mathcal{S}^*(q_c) = \{f \in \mathcal{A} : |(zf'(z)/f(z))^2 - 1| < c\}$ ($0 < c \leq 1$), has been briefly discussed in [17]. We consider the class $\mathcal{U}(\lambda)$ of analytic functions satisfying the following condition, $\mathcal{U}(\lambda) := \{f \in \mathcal{A} : |(z/f(z))^2 f'(z) - 1| < \lambda, \quad 0 < \lambda \leq 1\}$. From [16] it is known that the functions in $\mathcal{U}(\lambda)$ are univalent if $0 < \lambda \leq 1$, but not necessarily univalent if $\lambda > 1$.

Various sufficient conditions for Carathéodory functions were studied by authors in [5, 6, 11, 12, 13, 14]. Using differential subordination as a tool, authors in [13] and [14] obtained sufficient conditions for Carathéodory functions. Recently, Kim et al. [5] obtained sufficient conditions involving the argument of the function such that the function is Carathéodory. Motivated by the aforementioned works, in this paper various results involving analytic function to be Carathéodory are obtained and as a consequence, sufficient conditions for functions to belong to the classes $\mathcal{S}^*(q_c)$ and $\mathcal{U}(\lambda)$ are provided. The results thus obtained generalize and extend certain recent results.

2 Main results

To prove the main results we need the following Lemma.

Lemma 1 [4] *Let w be a non constant regular function in \mathbb{D} . If $|w|$ attains its maximum value on the circle $|z| = r < 1$ at z_0 , then*

$$z_0 w'(z_0) = k w(z_0),$$

where $k \geq 1$ is a real number.

Lemma 2 [1, 3] *Let $q \in \mathcal{Q}(a)$, and let $p(z) = a + a_n z^n + \dots$ be analytic in \mathbb{D} with $p(z) \not\equiv a$ and $n \geq 2$, if p is not subordinate to q then there exist points $z_0 = r_0 e^{i\theta_0} \in \mathbb{D}$ and $\zeta_0 \in \partial\mathbb{D} \setminus E(q)$ and an $m \geq n$ for which $p(\mathbb{D}_{r_0}) \subset q(\mathbb{D})$,*

1. $p(z_0) = q(\zeta_0)$
2. $\operatorname{Re} \left\{ \frac{\zeta_0 q''(\zeta_0)}{q'(\zeta_0)} \right\} \geq 0$ and $\left| \frac{zp'(z)}{q'(\zeta)} \right| \leq m$,
3. $z_0 p'(z_0) = m \zeta_0 q'(\zeta_0)$
4. $\operatorname{Re} \left\{ \frac{z_0 p''(z_0)}{p'(z_0)} + 1 \right\} \geq m \operatorname{Re} \left\{ \frac{\zeta_0 q''(\zeta_0)}{q'(\zeta_0)} + 1 \right\}$
5. $\operatorname{Re} \left\{ \frac{z_0^2 p'''(z_0)}{p'(z_0)} \right\} \geq m^2 \operatorname{Re} \left\{ \frac{\zeta_0^2 q'''(\zeta_0)}{q'(\zeta_0)} \right\}.$

Theorem 1 Let $0 \leq \alpha < 1$, $0 < \lambda \leq 1$, $\beta, \gamma, \delta, \mu \in \mathbb{R}$. For an analytic function p defined in \mathbb{D} with $p(0) = 1$, if

$$\operatorname{Re} \left\{ (p(z) - \alpha)^\lambda \left(\mu + \beta(p(z) - \alpha) + \frac{\gamma}{p(z) - \alpha} + \frac{\delta zp'(z)}{p(z) - \alpha} \right) \right\} > g(\epsilon(\alpha, \lambda), \alpha, \lambda), \quad (1)$$

where

$$g(u, \alpha, \lambda) = -u^{\lambda+1} \left(\beta + \frac{\delta}{2(1-\alpha)} \right) \sin \left(\frac{\lambda\pi}{2} \right) + \mu \cos \left(\frac{\lambda\pi}{2} \right) u^\lambda \\ + \left(\gamma - \frac{\delta(1-\alpha)}{2} \right) \sin \left(\frac{\lambda\pi}{2} \right) u^{\lambda-1}$$

and

$$\epsilon(\alpha, \lambda) = \frac{\mu \lambda \cos \left(\frac{\lambda\pi}{2} \right) + \sqrt{\mu^2 \lambda^2 \cos^2 \left(\frac{\lambda\pi}{2} \right) + 4(\lambda^2 - 1) \left(\beta + \frac{\delta}{2(1-\alpha)} \right) \left(\gamma - \frac{\delta(1-\alpha)}{2} \right) \sin^2 \left(\frac{\lambda\pi}{2} \right)}}{2(\lambda + 1) \left(\beta + \frac{\delta}{2(1-\alpha)} \right) \sin \left(\frac{\lambda\pi}{2} \right)}$$

then $p \in \mathcal{P}(\alpha)$.

Proof. Define the analytic function $p : \mathbb{D} \rightarrow \mathbb{C}$ as

$$p(z) = \frac{1 + (1 - 2\alpha)w(z)}{1 - w(z)}, \quad (2)$$

where w is an analytic function in \mathbb{D} with $w(0) = 0$. Suppose that there exists a point $z_0 \in \mathbb{D}$, such that

$$\operatorname{Re}\{p(z)\} > \alpha \quad \text{for} \quad |z| < |z_0| \quad \text{and} \quad \operatorname{Re}\{p(z_0)\} = \alpha, \quad (3)$$

then we have

$$|w(z)| < 1 \quad \text{for} \quad |z| < |z_0| \quad \text{and} \quad |w(z_0)| = 1. \quad (4)$$

By Lemma 1, we have $z_0 w'(z_0) = k w(z_0)$, where k is a real number with $k \geq 1$. Now,

$$\frac{z_0 p'(z_0)}{2k(1-\alpha)} = \frac{w(z_0)}{(1-w(z_0))^2} = \frac{2\{\operatorname{Re} w(z_0) - 1\}}{|1-w(z_0)|^4}.$$

Putting $p(z_0) = \alpha + iy$, we have $w(z_0) = 1 - \frac{2(1-\alpha)^2}{(1-\alpha)^2 + y^2} + i \frac{2(1-\alpha)y}{(1-\alpha)^2 + y^2}$ and

$$z_0 p'(z_0) = -k \frac{(1-\alpha)^2 + y^2}{2(1-\alpha)}, \quad (5)$$

which is a non positive real number. Also we observe that for the case $0 < \lambda < 1$

$$\begin{aligned} & \operatorname{Re} \left\{ (p(z_0) - \alpha)^\lambda \left(\mu + \beta(p(z_0) - \alpha) + \frac{\gamma}{p(z_0) - \alpha} + \delta \frac{z_0 p'(z_0)}{p(z_0) - \alpha} \right) \right\} \\ &= \operatorname{Re} \{ \beta(p(z_0) - \alpha)^{\lambda+1} + \mu(p(z_0) - \alpha)^\lambda + \gamma(p(z_0) - \alpha)^{\lambda-1} \\ & \quad + \delta z_0 p'(z_0)(p(z_0) - \alpha)^{\lambda-1} \} \\ &= \operatorname{Re} \left\{ \beta(iy)^{\lambda+1} + \mu(iy)^\lambda + \left(\gamma - k\delta \frac{(1-\alpha)^2 + y^2}{2(1-\alpha)} \right) (iy)^{\lambda-1} \right\} \\ &= \operatorname{Re} \left\{ \left(-\beta \sin \frac{\lambda\pi}{2} + i\beta \cos \frac{\lambda\pi}{2} - k \frac{\delta}{2(1-\alpha)} \left(\sin \frac{\lambda\pi}{2} - i \cos \frac{\lambda\pi}{2} \right) \right) |y|^{\lambda+1} \right. \\ & \quad \left. + \mu \left(\cos \frac{\lambda\pi}{2} + i \sin \frac{\lambda\pi}{2} \right) |y|^\lambda + \left(\gamma - k\delta \frac{1-\alpha}{2} \right) \left(\sin \frac{\lambda\pi}{2} - i \cos \frac{\lambda\pi}{2} \right) |y|^{\lambda-1} \right\} \\ &= \left(-\beta \sin \frac{\lambda\pi}{2} - k \frac{\delta}{2(1-\alpha)} \sin \frac{\lambda\pi}{2} \right) |y|^{\lambda+1} + \mu \left(\cos \frac{\lambda\pi}{2} \right) |y|^\lambda \\ & \quad + \left(\gamma - k \frac{\delta(1-\alpha)}{2} \right) \left(\sin \frac{\lambda\pi}{2} \right) |y|^{\lambda-1} \\ &\leq -\left(\beta + \frac{\delta}{2(1-\alpha)} \right) \sin \frac{\lambda\pi}{2} |y|^{\lambda+1} + \mu \cos \frac{\lambda\pi}{2} |y|^\lambda + \left(\gamma - \frac{\delta(1-\alpha)}{2} \right) \sin \frac{\lambda\pi}{2} |y|^{\lambda-1} \\ &= g(|y|, \alpha, \lambda) \leq \max_{u \in (0, \infty)} g(u, \alpha, \lambda) = g(\epsilon(\alpha, \lambda), \alpha, \lambda), \end{aligned}$$

which is a contradiction to (1). For the case when $\lambda = 1$,

$$\operatorname{Re} \left\{ (p(z_0) - \alpha) \left(\mu + \beta(p(z_0) - \alpha) + \frac{\gamma}{p(z_0) - \alpha} + \delta \frac{z_0 p'(z_0)}{p(z_0) - \alpha} \right) \right\}$$

$$\begin{aligned}
&= \operatorname{Re} \left\{ (p(z_0) - \alpha) + \left(\mu + \beta(p(z_0) - \alpha) + \frac{\gamma}{p(z_0) - \alpha} - \delta k \left(\frac{(1 - \alpha)^2 + y^2}{2(1 - \alpha)p(z_0) - \alpha} \right) \right) \right\} \\
&\leq \operatorname{Re} \left\{ (iy)\mu + \beta(iy)^2 + \gamma - \delta \left(\frac{(1 - \alpha)^2 + y^2}{2(1 - \alpha)} \right) \right\} \\
&= -\left(\beta + \frac{\delta}{2(1 - \alpha)} \right) y^2 + \gamma - \delta \frac{(1 - \alpha)}{2} \\
&\leq \gamma - \delta \frac{(1 - \alpha)}{2} = g(\epsilon(\alpha, 1), \alpha, 1).
\end{aligned}$$

This contradicts (1). Hence the proof. \square

Remark 1 By taking $\mu = \delta = 1$ and $\beta = \gamma = 0$ in Theorem 1, we get the result obtained in [6, Theorem 2.29].

By taking $p(z) = zf'(z)/f(z)$ and $\alpha = 0$ in Theorem 1, we have the following result:

Corollary 1 For a function $f \in \mathcal{A}$ and $0 < \lambda \leq 1$, if

$$\operatorname{Re} \left\{ (\beta - \delta) \left(\frac{zf'(z)}{f(z)} \right)^{\lambda+1} + \left(\mu + \delta + \delta \frac{zf''(z)}{f'(z)} \right) \left(\frac{zf'(z)}{f(z)} \right)^{\lambda} + \gamma \left(\frac{zf'(z)}{f(z)} \right)^{\lambda-1} \right\} > g(\epsilon(0, \lambda), 0, \lambda),$$

then $f \in \mathcal{S}^*$.

Theorem 2 For an analytic function p in \mathbb{D} with $p(0) = 1$ and $0 \leq \alpha < 1$, if

$$\frac{zp'(z)}{(p(z) - \alpha)^{1/\beta}} \neq i\delta \tag{6}$$

for all $\delta \in \mathbb{R}$, with $|\delta| \geq 1$ and $\beta = \frac{1}{2n-1}$, $n \in \mathbb{N}$, then $p \in \mathcal{P}(\alpha)$.

Proof. Let

$$h(z) = \left(\frac{p(z) - \alpha}{1 - \alpha} \right)^{1/\beta}.$$

We note that h is analytic in \mathbb{D} , with $h(0) = 1$. Here $p(z) \neq \alpha$ for $z \in \mathbb{D}$, suppose that there exist a point $z_1 \in \mathbb{D}$ such that $p(z_1) = \alpha$, then z_1 is a zero of multiplicity $m \geq 1$ such that

$$h(z) = (z - z_1)^m g(z) \quad (m \in \mathbb{N}),$$

where $g(z)$ is analytic in \mathbb{D} and $g(z_1) \neq 0$. Therefore we have

$$\frac{zp'(z)}{(p(z) - \alpha)} = \beta \left(\frac{mz}{z - z_1} + \frac{zg'(z)}{g(z)} \right). \quad (7)$$

But the imaginary part of right-hand side of (7) can take any value when z approaches z_1 . This contradicts (6). Therefore $p(z) \neq \alpha$, that is $h(z) \neq 0$. Suppose that there exist a point $z_0 \in \mathbb{D}$ such that

$$\operatorname{Re}\{(h(z))^\beta\} > 0 \quad \text{for } |z| < |z_0| \quad \text{and} \quad \operatorname{Re}\{(h(z_0))^\beta\} = 0 \quad (h(z_0) \neq 0).$$

Setting

$$\phi(z) = \frac{1 - (h(z))^\beta}{1 + (h(z))^\beta},$$

we observe that

$$|\phi(z)| < 1 \quad \text{for } |z| < |z_0|, \quad |\phi(z_0)| = 1 \quad \text{and} \quad \phi(0) = 0.$$

Hence the conditions of Lemma 1 are satisfied. By taking

$$(h(z_0))^\beta = iy,$$

where y is a non zero positive real number, and by using Lemma 1, we obtain

$$\frac{z_0\phi'(z_0)}{\phi(z_0)} = \frac{-2\beta(h(z_0))^{\beta-1}z_0h'(z_0)}{1 - (h(z_0))^{2\beta}} = k,$$

and

$$-z_0h'(z_0) = \frac{k(1 + y^2)}{2\beta(h(z_0))^{-1}(iy)}.$$

Now,

$$\begin{aligned} \frac{z_0p'(z_0)}{(p(z_0) - \alpha)^{1/\beta}} &= \frac{-k(1 - \alpha)^{1-(1/\beta)}(1 + y^2)}{2((iy)^{1/\beta})} \\ &= \frac{-k(1 - \alpha)^{1-(1/\beta)}(1 + y^2)}{2y^{1/\beta}} \left(\cos \frac{\pi}{2\beta} - i \sin \frac{\pi}{2\beta} \right). \end{aligned}$$

For $\beta = \frac{1}{2n-1}$, $n \in \mathbb{N}$

$$\frac{z_0p'(z_0)}{(p(z_0) - \alpha)^{1/\beta}} = \frac{-k(1 - \alpha)^{1-(1/\beta)}(1 + y^2)}{2y^{1/\beta}} (-1)^{n+1}i = i\delta, \quad \delta \in \mathbb{R},$$

which is a contradiction to (6), where

$$\begin{aligned} |\delta| &= \left| \frac{k(1-\alpha)^{1-(1/\beta)}(1+y^2)(-1)^{n+2}}{2y^{1/\beta}} \right| \\ &\geq \frac{(1-\alpha)^{1-(1/\beta)}(1+y^2)}{2y^{1/\beta}} \geq 1. \end{aligned}$$

Hence the proof. \square

For the choice of $p(z) = zf'(z)/f(z)$, $\beta = 1$ and $p(z) = f'(z)$, $\beta = 1$ and $\alpha = 0$, in Theorem 2, we have the following Corollary 2 and Corollary 3 respectively.

Corollary 2 For $0 \leq \alpha < 1$, if the function $f \in \mathcal{A}$ satisfies

$$\frac{zf'(z)\left(1 - \frac{zf'(z)}{f(z)}\right) + z^2f''(z)}{zf'(z) - \alpha f(z)} \neq i\delta \quad (\delta \in \mathbb{R}, |\delta| \geq 1),$$

then $f \in \mathcal{S}^*(\alpha)$.

Corollary 3 If $f \in \mathcal{A}$ satisfies

$$\frac{zf''(z)}{f'(z)} \neq i\delta \quad (\delta \in \mathbb{R}, |\delta| \geq 1),$$

then f is univalent.

Theorem 3 Let $\alpha, \beta, \gamma, \delta \in \mathbb{R}$ with $0 \leq \alpha < 1$, $\gamma < \frac{\beta + \delta}{2}$ and let $G(z)$ be a complex valued function defined in \mathbb{D} . If p is analytic in \mathbb{D} with $p(0) = 1$ and

$$\begin{aligned} &\operatorname{Re}\{\gamma z^3 p'''(z) + (3\gamma + \beta)z^2 p''(z) + (\gamma + 2\beta + \delta)z p'(z) + G(z)p(z)\} \\ &> \mu(\alpha, \beta, \gamma, \delta, G(z)), \end{aligned} \quad (8)$$

where

$$\mu(\alpha, \beta, \gamma, \delta, G(z)) = \frac{(1-\alpha)[\operatorname{Im}(G(z))]^2 - [\delta + \beta - 2\gamma]^2}{2(\delta + \beta - 2\gamma)} + \alpha \operatorname{Re}\{G(z)\},$$

then $p \in \mathcal{P}(\alpha)$.

Proof. Let the function p be defined as in (2), suppose that there exists a point z_0 in \mathbb{D} satisfying (3). By defining $h : \mathbb{D} \rightarrow \mathbb{C}$ as $h(z) = (1 + (1 - 2\alpha)z)/(1 - z)$, we have $p \not\prec h$. By Lemma 2, there exist a $\zeta_0 \in \partial\mathbb{D}$ and $m \geq 1$ such that

$$\operatorname{Re} \left\{ 1 + \frac{z_0 p''(z_0)}{p'(z_0)} \right\} \geq m \operatorname{Re} \left\{ 1 + \frac{\zeta_0 h''(\zeta_0)}{h'(\zeta_0)} \right\} = 0$$

and (9)

$$\operatorname{Re} \left\{ \frac{z_0^2 p'''(z_0)}{p'(z_0)} \right\} \geq m^2 \operatorname{Re} \left\{ \frac{\zeta_0^2 h'''(\zeta_0)}{h'(\zeta_0)} \right\} > 0.$$

Using (5) and (9), we obtain

$$\operatorname{Re}\{z_0^2 p''(z_0)\} \leq -z_0 p'(z_0) \quad \text{and} \quad \operatorname{Re}\{z_0^3 p'''(z_0)\} \leq 0. \quad (10)$$

From (5), (10) and by taking $p(z) = \alpha + iy$ ($y \in \mathbb{R}$), we have the following inequality

$$\begin{aligned} & \operatorname{Re}\{\gamma z_0^3 p'''(z_0) + (3\gamma + \beta)z_0^2 p''(z_0) + (\gamma + 2\beta + \delta)z_0 p'(z_0) + G(z)p(z_0)\} \\ & \leq (\delta + \beta - 2\gamma)z_0 p'(z_0) + \alpha \operatorname{Re}\{G(z_0)\} - \operatorname{Im}\{G(z_0)\}y \\ & \leq -(\delta + \beta - 2\gamma) \left(\frac{(1 - \alpha)^2 + y^2}{2(1 - \alpha)} \right) + \alpha \operatorname{Re}\{G(z_0)\} - \operatorname{Im}\{G(z_0)\}y \\ & \leq \frac{(1 - \alpha)(\operatorname{Im}\{G(z_0)\})^2 - (\delta + \beta - 2\gamma)^2}{2(\delta + \beta - 2\gamma)} + \alpha \operatorname{Re}\{G(z_0)\} \\ & = \mu(\alpha, \beta, \gamma, \delta, G(z_0)), \end{aligned}$$

which contradicts (8) and completes the proof. □

On taking $\alpha = \beta = \gamma = 0$, $\delta = 1$ and $G(z) \equiv 1$ in Theorem 3, we get the following Corollary that improves the result of Miller [7, p.80].

Corollary 4 *For an analytic function p in \mathbb{D} with $p(0) = 1$, if*

$$\operatorname{Re}\{p(z) + zp'(z)\} > -\frac{1}{2},$$

then $p \in \mathcal{P}$.

By taking $\gamma = \beta = 0$, $G(z) \equiv 1$ and $\gamma = \beta = 0$, $\delta = 1$, $G(z) \equiv 1$ in Theorem 3, we obtain the Corollary 5 and Corollary 6 respectively, which are due to Kim et al. [6, Theorem 2.6].

Corollary 5 For an analytic function p in \mathbb{D} with $p(0) = 1$, if

$$\operatorname{Re}\{\delta zp'(z) + p(z)\} > \alpha - \frac{(1-\alpha)\delta}{2}, \quad (0 \leq \alpha < 1)$$

then $p \in \mathcal{P}(\alpha)$.

Corollary 6 For an analytic function p in \mathbb{D} with $p(0) = 1$, if

$$\operatorname{Re}\{zp'(z) + p(z)\} > \frac{(3\alpha - 1)}{2}, \quad (0 \leq \alpha < 1)$$

then $p \in \mathcal{P}(\alpha)$.

Theorem 4 Let p be an analytic function in \mathbb{D} , with $p(0) = 1$ for $\beta > 0$, if

$$\left| \operatorname{Im} \left\{ (p(z))^{1/\beta} + \frac{zp'(z)}{p(z)} \right\} \right| < \frac{1}{2|(p(z))^{1/\beta}|} \left((2 - \beta)|(p(z))^{2/\beta} - \beta \right), \quad (11)$$

then $(p(z))^{1/\beta} \in \mathcal{P}$.

Proof. Define the function $p : \mathbb{D} \rightarrow \mathbb{C}$ as

$$p(z) = \left(\frac{1 + w(z)}{1 - w(z)} \right)^\beta$$

or equivalently

$$w(z) = \frac{p(z)^{1/\beta} - 1}{p(z)^{1/\beta} + 1},$$

then w is analytic in \mathbb{D} with $w(0) = 0$. Suppose that there exist a point z_0 in \mathbb{D} such that

$$\operatorname{Re}\{(p(z))^{1/\beta}\} > 0 \quad \text{for } |z| < |z_0| \quad \text{and} \quad \operatorname{Re}\{(p(z_0))^{1/\beta}\} = 0,$$

we obtain

$$|w(z)| < 1 \quad \text{for } |z| < |z_0| \quad \text{and} \quad |w(z_0)| = 1.$$

Therefore by using Jack's Lemma, a simple calculation yields

$$\frac{1}{\beta} \frac{z_0 p'(z_0)}{p(z_0)} = \frac{2z_0 w'(z_0)}{1 - (w(z_0))^2} = \frac{2kw(z_0)}{1 - (w(z_0))^2}.$$

Hence

$$\frac{1}{2k\beta} \frac{z_0 p'(z_0)}{p(z_0)} = \frac{w(z_0)}{1 - (w(z_0))^2}.$$

On taking $(p(z_0))^{1/\beta} = iy$, where y is a nonzero real number, we obtain

$$w(z_0) = \frac{y^2 - 1}{1 + y^2} + i \frac{2y}{1 + y^2}$$

and

$$\frac{z_0 p'(z_0)}{p(z_0)} = \frac{2k\beta w(z_0)}{1 - (w(z_0))^2},$$

where k is a real number, with $k \geq 1$. Therefore

$$\begin{aligned} \left| \operatorname{Im} \left\{ (p(z_0))^{1/\beta} + \frac{z_0 p'(z_0)}{p(z_0)} \right\} \right| &= \left| \operatorname{Im} \{ (p(z_0))^{1/\beta} \} + \operatorname{Im} \left\{ \frac{z_0 p'(z_0)}{p(z_0)} \right\} \right| \\ &= \left| y + 2k\beta \frac{(1 + y^2)}{4y} \right| \\ &\geq \left| y + \beta \frac{(1 + y^2)}{2y} \right| \\ &\geq \left| y - \beta \frac{(1 + y^2)}{2y} \right| \geq |y| - \beta \frac{(1 + |y|^2)}{2|y|} \\ &= |(p(z_0))^{1/\beta}| - \frac{\beta(1 + |(p(z_0))^{2/\beta}|)}{2|(p(z_0))^{1/\beta}|}, \end{aligned}$$

which contradicts (11). Hence the proof. \square

Theorem 5 For an analytic function p in \mathbb{D} with $p(0) = 1$, if p satisfies

$$\operatorname{Re} \left\{ (p(z))^2 + \frac{zp'(z)}{p(z)} \right\} < 1 - c - \frac{c}{2(1 - c)} \quad (0 < c < 1), \quad (12)$$

then $p \in \mathcal{P}$. Also $p(z) \prec \sqrt{1 + cz}$.

Proof. Define a function $p : \mathbb{D} \rightarrow \mathbb{C}$ by

$$\begin{aligned} p(z) &= \sqrt{1 + cw(z)}, \quad (z \in \Delta) \\ &= 1 + p_1 z + p_2 z^2 + \dots, \end{aligned}$$

or equivalently

$$w(z) = \frac{p^2(z) - 1}{c} = w_1 z + w_2 z^2 + \dots,$$

we observe that w is analytic in \mathbb{D} and $w(0) = 0$.

Suppose that there exist a point z_0 in \mathbb{D} , such that

$$\operatorname{Re}\{p(z)\} > 0 \quad \text{for} \quad |z| < |z_0| \quad \text{and} \quad \operatorname{Re}\{p(z_0)\} = 0$$

and

$$\max |w(z)| = |w(z_0)| = 1 \quad |z| \leq |z_0|.$$

By Lemma 1, there exist a number $k \geq 1$ such that $z_0 w'(z_0) = kw(z_0)$. Without loss of generality we may assume that $w(z_0) = e^{i\theta}$, where $\theta \in [-\pi, \pi]$, for this z_0 , we have

$$\begin{aligned} \operatorname{Re} \left\{ (p(z_0))^2 + \frac{z_0 p'(z_0)}{p(z_0)} \right\} &= \operatorname{Re}\{1 + cw(z_0)\} + \operatorname{Re} \left\{ \frac{ckw(z_0)}{2(1 + cw(z_0))} \right\} \\ &= \operatorname{Re}\{1 + ce^{i\theta}\} + \frac{ck}{2} \operatorname{Re} \left\{ \frac{e^{i\theta}}{1 + ce^{i\theta}} \right\} \\ &= \operatorname{Re}\{1 + c \cos \theta + i \sin \theta\} + \frac{ck}{2} \operatorname{Re} \left\{ \frac{\cos \theta + i \sin \theta}{1 + ce^{i\theta}} \right\} \\ &\geq 1 + c \cos \theta + \frac{c}{2} \left(\frac{\cos \theta + c}{1 + c^2 + 2c \cos \theta} \right) = H(\cos \theta). \end{aligned}$$

Let $t = \cos \theta$ then

$$H(t) = 1 + ct + \frac{c}{2} \left(\frac{t + c}{1 + c^2 + 2ct} \right).$$

Since $H(t)$ is an increasing function,

$$\begin{aligned} H(t) &\geq H(-1) = 1 - c + \frac{c}{2} \left(\frac{c - 1}{1 + c^2 - 2c} \right) \\ &= 1 - c - \frac{c}{2} \left(\frac{1 - c}{(1 - c)^2} \right) \\ &= 1 - c - \frac{c}{2(1 - c)}, \end{aligned}$$

which is a contradiction to (12) and implies that, $\operatorname{Re}\{p(z)\} > 0$ and $|w(z_0)| < 1$. That is $p(z) \prec \sqrt{1 + cz}$ and $p \in \mathcal{P}$, $z \in \mathbb{D}$. \square

Following results are obtained as the consequence of Theorem 5.

For the choice of $p(z) = \frac{zf'(z)}{f(z)}$ in Theorem 5, we have the following:

Corollary 7 If $f \in \mathcal{A}$ satisfies

$$\operatorname{Re} \left\{ \left(\frac{zf'(z)}{f(z)} \right)^2 + \frac{zf''(z)}{f'(z)} - \frac{zf'(z)}{f(z)} + 1 \right\} < (1 - c) - \frac{c}{2(1 - c)} \quad (0 < c < 1),$$

then $f \in \mathcal{S}^*(q_c)$.

By taking $p(z) = \frac{z\sqrt{f'(z)}}{f(z)}$ in Theorem 5, we have the following:

Corollary 8 If $f \in \mathcal{A}$ satisfies

$$\operatorname{Re} \left\{ \frac{z^2 f'(z)}{(f(z))^2} + \frac{zf''(z)}{2f'(z)} - \frac{zf'(z)}{f(z)} + 1 \right\} < (1 - \lambda) - \frac{\lambda}{2(1 - \lambda)} \quad (0 < \lambda < 1),$$

then $f \in \mathcal{U}(\lambda)$ and hence it is univalent.

Acknowledgement

The authors are thankful to the referee for the insightful suggestions.

References

- [1] J. A. Antonino, S. S. Miller, Third-order differential inequalities and subordinations in the complex plane, *Complex Var. Elliptic Equ.*, **56** (5) (2011), 439–454.
- [2] P. L. Duren, *Univalent functions*, Grundlehren der Mathematischen Wissenschaften, 259, Springer-Verlag, New York, 1983.
- [3] Huo Tang, H. M. Srivastava, Shu-Hai Li, Li-Na Ma, Third-order differential subordination and superordination results for meromorphically multivalent functions associated with the Liu-Srivastava operator, *Abstr. Appl. Anal.*, **2014**, Art. ID 792175, 11 pp.
- [4] I. S. Jack, Functions starlike and convex of order α , *J. London Math. Soc.*, **3** (2) (1971), 469–474.
- [5] I. H. Kim, N. E. Cho, Sufficient conditions for Carathéodory functions, *Comput. Math. Appl.*, **59** (6) (2010), 2067–2073.
- [6] I. H. Kim, Y. J. Sim, N. E. Cho, New criteria for Carathéodory functions, *J. Inequal. Appl.*, **13** (2019), 16 pp.

- [7] S. Miller, Differential inequalities and Carathéodory functions, *Bull. Amer. Math. Soc.*, **81** (1975), 79–81.
- [8] S. S. Miller, P. T. Mocanu, Differential subordinations and univalent functions, *Michigan Math. J.*, **28** (2) (1981), 157–172.
- [9] S. S. Miller, P. T. Mocanu, *Differential subordinations*, Monographs and Textbooks in Pure and Applied Mathematics, 225, Marcel Dekker, Inc., New York, 2000.
- [10] S. S. Miller, P. T. Mocanu, M. O. Reade, All α -convex functions are starlike, *Rev. Roumaine Math. Pures Appl.*, **17** (1972), 1395–1397.
- [11] N. Xu, Sufficient conditions and applications for Carathéodory functions, *J. Math.*, **2013**, Art. ID 930290, 4 pp.
- [12] M. Nunokawa, Differential inequalities and Carathéodory functions, *Proc. Japan Acad. Ser. A Math. Sci.*, **65** (10) (1989), 326–328.
- [13] M. Nunokawa, A. Ikeda, N. Koike, Y. Ota, H. Saitoh, Differential inequalities and Carathéodory functions, *J. Math. Anal. Appl.*, **212** (1) (1997), 324–332.
- [14] M. Nunokawa, S. Owa, N. Takahashi, H. Saitoh, Sufficient conditions for Carathéodory functions, *Indian J. Pure Appl. Math.*, **33** (9) (2002), 1385–1390.
- [15] M. Nunokawa, M. Obradović, S. Owa, One criterion for univalence, *Proc. Amer. Math. Soc.*, **106** (4) (1989), 1035–1037.
- [16] S. Ponnusamy, P. Vasundhra, Starlikeness of nonlinear integral transforms, *J. Anal.*, **15** (2007), 195–210.
- [17] J. Sokół, On some subclass of strongly starlike functions, *Demonstratio Math.*, **31** (1) (1998), 81–86.
- [18] J. Sokół, J. Stankiewicz, Radius of convexity of some subclasses of strongly starlike functions, *Zeszyty Nauk. Politech. Rzeszowskiej*, **19** (1996), 101–105.

Received: July 4, 2020



On γ –countably paracompact sets

Amani Rawshdeh

Department of Mathematics,
Al-Balqa Applied University,
Asalt 19117, Jordan

email: amanirawshdeh@bau.edu.jo
(Corresponding author)

Heyam H. Al-Jarrah

Department of Mathematics,
Faculty of science Yarmouk University,
Irbid 21163, Jordan

email: hiamaljarah@yahoo.com;
heyam@yu.edu.jo

Khalid Y. Al- Zoubi

Department of Mathematics,
Faculty of science Yarmouk
University,

Irbid 21163, Jordan
email: khalidz@yu.edu.jo

Wasfi A. Shatanawi

Department of Mathematics and General
Courses, Prince Sultan University, Riyadh
11586, Saudi Arabia.

Department of Medical Research, China
Medical University Hospital, China Medical
University, Taichung 40402, Taiwan
email: wshatanawi@yahoo.com

Abstract. In this paper we introduce and study a new class of sets, namely γ –countably paracompact sets. We characterize γ –countably paracompact sets and we study some of its basic properties. We obtain that this class of sets is weaker than α –countably paracompact sets and stronger than β –countably paracompact sets.

1 Introduction

In [3] C. E. Aull, presented and studied the concept of α –countably paracompact and β –countably paracompact sets. In connection with the definition of α –countably paracompact sets and β –countably paracompact sets we obtain the definition of γ –countably paracompact sets. In section 2 of this work, we

2010 Mathematics Subject Classification: 54A05, 54C08, 54D10

Key words and phrases: countably paracompact, α –countably paracompact, β –countably paracompact, γ –countably paracompact

present γ -countably paracompact sets and then we investigate several characterizations to this types of sets and study some of its basic properties. In section 3 of this work, some of relationships between γ -countably paracompact sets and other well-known sets are investigated. In particular, we show that this class of sets lies between the classes of α -countably paracompact sets and β -countably paracompact sets. Finally, in section 4, we introduce a class of spaces namely locally γ -countably paracompact spaces characterized by γ -countably paracompact sets and study some of their fundamental properties.

Throughout this work a space will always mean a topological space on which no separation axiom is assumed unless explicitly stated. Let (X, τ) be a space and A be a subset of X . The closure of A , interior of A and the relative topology on A in (X, τ) will be denoted by $\text{cl}(A)$, $\text{int}(A)$ and τ_A , respectively. A space (X, τ) is called countably paracompact [4] if every countable open cover of X has an open locally finite refinement. Now we begin with some known notions and definitions which will be used in this work.

Definition 1 [5] *A subset A of a space (X, τ) is called generalized closed if $\text{cl}(A) \subseteq U$ whenever $A \subseteq U$ and U is open in (X, τ) .*

Theorem 1 [4] *If A is dense in X , then for every open $U \subseteq X$ we have $\text{cl}(U) = \text{cl}(U \cap A)$.*

Definition 2 *Let A, B, C and Y be subsets of a space (X, τ) . Then:*

- i. *A cover \mathcal{U} of Y is called an A -open cover of Y [1] if $Y \subseteq \bigcup_{U \in \mathcal{U}} U$ and U is open in (A, τ_A) for every $U \in \mathcal{U}$.*
- ii. *A collection $\mathcal{U} = \{U_\alpha : \alpha \in \Delta\}$ is called A -locally finite [1] if \mathcal{U} is locally finite in (A, τ_A) .*
- iii. *If \mathcal{U} and \mathcal{V} are covers of Y , then \mathcal{V} is called A -refinement of \mathcal{U} [1] if for every $V \in \mathcal{V}$, $V \subseteq A$ and there exists $U \in \mathcal{U}$ such that $V \subseteq U$. If for every $V \subseteq \mathcal{V}$, V is open in (A, τ_A) then \mathcal{V} is called an A -open refinement of \mathcal{U} .*
- iv. *Y is called α -countably paracompact of (X, τ) [3] if every countable open cover of Y by members of τ has a locally finite open refinement by members of τ .*
- v. *Y is called β -countably paracompact of (X, τ) [3] if (Y, τ_Y) is countably paracompact as a subspace.*

The proof of the following proposition is obvious.

Proposition 1 *Let Y be a subset of a topological space (X, τ) . If a collection $\mathcal{U} = \{U_\alpha : \alpha \in I\}$ is X -locally finite, then \mathcal{U} is Y -locally finite.*

The following example shows that the converse of the above proposition is not true in general.

Example 1 *Let $X = \mathbb{R}$ with the topology $\tau = \{U : 0 \notin U\} \cup \{\mathbb{R}\}$. Put $Y = \mathbb{Q}^* = \mathbb{Q} - \{0\}$. Then the collection $\{\{y\} : y \in Y\}$ is Y -locally finite but it is not X -locally finite.*

In the following proposition, we shall show when the converse of the above proposition is true.

Proposition 2 *Let Y be a closed subset of a topological space (X, τ) . If $\mathcal{U} = \{U_\alpha : \alpha \in I, U_\alpha \subseteq Y\}$ is a Y -locally finite collection of subsets of Y , then \mathcal{U} is X -locally finite.*

Proof. Let $\mathcal{U} = \{U_\alpha : \alpha \in I\}$ be Y -locally finite such that $U_\alpha \subseteq Y$ for each $\alpha \in I$. If $x \in X$, then either $x \in Y$ or $x \notin Y$. If $x \in Y$, then there exists an open set W in (Y, τ_Y) such that $x \in W$ and W intersects at most finitely many members of \mathcal{U} . Now $W = M \cap Y$ for some $M \in \tau$. As \mathcal{U} is a collection of subsets of Y , so M intersects at most finitely many members of \mathcal{U} . Now if $x \notin Y$, then $X - Y$ is open in (X, τ) containing x which intersects no member of \mathcal{U} . \square

Corollary 1 *Let Y be a closed subset of a topological space (X, τ) . The collection $\{U_\alpha : \alpha \in I, U_\alpha \subseteq Y\}$ is Y -locally finite iff \mathcal{U} is X -locally finite.*

2 γ -countably paracompact sets

In this section we shall present the concept of γ -countably paracompact sets.

Definition 3 *Let A, B, C and Y be subsets of a space (X, τ) . Then Y is called ABC -countably paracompact set of (X, τ) , if every countable A -open cover of Y has a B -locally finite C -open refinement.*

Note that a subset Y of a space (X, τ) is α -countably paracompact iff it is XXX -countably paracompact and it is β -countably paracompact iff it is YYY -countably paracompact.

Definition 4 Let Y be a subset of a topological space (X, τ) . Then Y is called γ -countably paracompact if it is XXY -countably paracompact.

Example 2 Let $X = \mathbb{R}$ with the topology $\tau = \{U : \mathbb{R} - \mathbb{Q} \subseteq U\} \cup \{\emptyset\}$. Then $Y = \mathbb{Q}$ is γ -countably paracompact.

Proposition 3 Let Y be a subset of a topological space (X, τ) . Then Y is γ -countably paracompact iff for every countable X -open cover $\mathcal{U} = \{U_n : n \in \mathbb{N}\}$ of Y there exists an X -locally finite Y -open cover $\mathcal{V} = \{V_n : n \in \mathbb{N}\}$ of Y such that $V_n \subseteq U_n$ for $n = 1, 2, \dots$

Proof. Let Y be a γ -countably paracompact set. If $\mathcal{U} = \{U_n : n \in \mathbb{N}\}$ is a countable X -open cover of Y , then there exists an X -locally finite Y -open refinement of \mathcal{U} , say \mathcal{W} . So for every $W \in \mathcal{W}$ choose a natural number $n(W)$ such that $W \subseteq U_{n(W)}$. Then define $V_n = \bigcup_{n \in \mathbb{N}} \{W : n(W) = n\}$. Hence $\mathcal{V} = \{V_n : n \in \mathbb{N}\}$ is open in Y and it is X -locally finite such that $V_n \subseteq U_n$ for $n = 1, 2, \dots$ \square

Proposition 4 Let Y be a subset of a topological space (X, τ) . If Y is γ -countably paracompact, then for every increasing countable X -open cover $\mathcal{U} = \{U_n : n \in \mathbb{N}\}$ of Y there exists a Y -open cover $\mathcal{V} = \{V_n : n \in \mathbb{N}\}$ of Y such that $cl_Y(V_n) \subseteq U_n$ for $n = 1, 2, \dots$

Proof. Let $\mathcal{U} = \{U_n : n \in \mathbb{N}\}$ be an increasing countable X -open cover of Y . Then, by Proposition 3, there exists an X -locally finite Y -open cover $\mathcal{V} = \{V_n : n \in \mathbb{N}\}$ of Y such that $V_n \subseteq U_n$. To show that $cl_Y(V_n) \subseteq U_n$, set $F_n = Y - \bigcup_{m > n} V_m$. Then F_n is closed in Y such that $V_n \subseteq F_n \subseteq \bigcup_{m \leq n} V_m \subseteq U_n$ and so $cl_Y(V_n) \subseteq U_n$. \square

Proposition 5 Let Y be a subset of a topological space (X, τ) and suppose that for every countable X -open cover $\mathcal{U} = \{U_n : n \in \mathbb{N}\}$ of Y there exists an X -locally finite Y -open cover $\mathcal{V} = \{V_n : n \in \mathbb{N}\}$ of Y such that $V_n \subseteq U_n$ for $n = 1, 2, \dots$. If $W_1 \subseteq W_2 \subseteq \dots$ is an increasing sequence of open sets in X such that $\bigcup_{n \in \mathbb{N}} W_n = Y$, then there exists a sequence $F_1 \subseteq F_2 \subseteq \dots$ of closed subsets of Y such that $F_n \subseteq W_n$ for $n = 1, 2, \dots$ and $\bigcup_{n \in \mathbb{N}} \text{int}_Y(F_n) = Y$.

Proof. Let $W_1 \subseteq W_2 \subseteq \dots$ be an increasing sequence of X -open sets such that $\bigcup_{n \in \mathbb{N}} W_n = Y$. Then, there exists an X -locally finite Y -open cover $\mathcal{V} = \{V_n : n \in \mathbb{N}\}$

$n \in \mathbb{N}$ of Y such that $V_n \subseteq W_n$ for all n . Now, define $F_n = Y - \bigcup_{j>n} V_j$ which is closed in Y and for $n \in \mathbb{N}$ we have $F_n \subseteq \bigcup_{j \leq n} V_j \subseteq \bigcup_{j \leq n} W_j = W_n$. To show that $\bigcup_{n \in \mathbb{N}} \text{int}_Y(F_n) = Y$, it is enough to show that $Y \subseteq \bigcup_{n \in \mathbb{N}} \text{int}_Y(F_n)$. Let $y \in Y$. Then there exists an open O in (X, τ) such that $y \in O$ and $(O \cap Y) \cap \bigcup_{m>n} V_m = \emptyset$ for some $n \in \mathbb{N}$, so we have $y \in O \cap Y \subseteq Y - \bigcup_{m>n} V_m = F_n$. Hence $y \in \bigcup_{n \in \mathbb{N}} \text{int}_Y(F_n)$. \square

Proposition 6 *Let Y be a closed subset of a topological space (X, τ) . Then every countable X -open cover $\mathcal{U} = \{U_n : n \in \mathbb{N}\}$ of Y has an X -locally finite Y -open cover $\mathcal{V} = \{V_n : n \in \mathbb{N}\}$ such that $V_n \subseteq U_n$ for all n iff for every increasing sequence $W_1 \subseteq W_2 \subseteq \dots$ of open sets in X such that $Y = \bigcup_{n \in \mathbb{N}} W_n$ there exists a sequence $F_1 \subseteq F_2 \subseteq \dots$ of closed subsets of Y such that $F_n \subseteq W_n$ for $n = 1, 2, \dots$, moreover $\bigcup_{n \in \mathbb{N}} \text{int}_Y(F_n) = Y$.*

Proof. We show the sufficiency part. Let $\mathcal{U} = \{U_n : n \in \mathbb{N}\}$ be a countable X -open cover of Y . Set $W_n = \bigcup_{j \leq n} U_j$. Then $W_1 \subseteq W_2 \subseteq \dots$, such that $\bigcup_{n \in \mathbb{N}} W_n = Y$. So there exists $F_1 \subseteq F_2 \subseteq \dots$ of closed subsets of Y such that $F_n \subseteq W_n$ for $n = 1, 2, \dots$ and $\bigcup_{n \in \mathbb{N}} \text{int}_Y(F_n) = Y$. Define $V_n = (U_n \cap Y) - \bigcup_{j < n} F_j$. Then V_n is open in Y and $V_n \subseteq U_n$ for $n = 1, 2, \dots$. To show that $\bigcup_{n \in \mathbb{N}} V_n = Y$, let $y \in Y$ and j be the first index such that $y \in (U_j \cap Y)$. Therefore, $y \in V_j$. To complete the proof we show that $\mathcal{V} = \{V_n : n \in \mathbb{N}\}$ is Y -locally finite. Let $y \in Y$. Then there exists j such that $x \in \text{int}_Y(F_j)$ and $\text{int}_Y(F_j) \cap V_n = \emptyset$ for $n > j$. Therefore, \mathcal{V} is Y -locally finite and so by Proposition 2, $\mathcal{V} = \{V_n : n \in \mathbb{N}\}$ is X -locally finite. \square

From above discussion we can get the following Theorem.

Theorem 2 *Let Y be a closed subset of a topological space (X, τ) . Then the following are equivalent:*

- i. Y is γ -countably paracompact.
- ii. For every countable X -open cover $\mathcal{U} = \{U_n : n \in \mathbb{N}\}$ of Y , there exists an X -locally finite Y -open cover $\mathcal{V} = \{V_n : n \in \mathbb{N}\}$ of Y such $V_n \subseteq U_n$ for all n .

- iii. For every increasing sequence $W_1 \subseteq W_2 \subseteq \dots$ of open sets in X such that $\bigcup_{n \in \mathbb{N}} W_n = Y$, there exists $F_1 \subseteq F_2, \dots$ of closed subsets of Y such that $F_n \subseteq W_n$ for $n = 1, 2, \dots$, moreover $\bigcup_{n \in \mathbb{N}} \text{int}_Y(F_n) = Y$.
- iv. For every increasing countable X -open cover $\mathcal{U} = \{U_n : n \in \mathbb{N}\}$ of Y , there exists a Y -open cover $\mathcal{V} = \{V_n : n \in \mathbb{N}\}$ of Y such that $\text{cl}_Y(V_n) \subseteq U_n$ for $n = 1, 2, \dots$.
- v. For every decreasing X -closed collection $\mathcal{F} = \{F_n : n \in \mathbb{N}\}$ such that $(\bigcap_{n \in \mathbb{N}} F_n) \cap Y = \emptyset$, there exists a Y -open cover $\mathcal{O} = \{O_n : n \in \mathbb{N}\}$ of Y such that $\text{cl}_Y(O_n) \cap F_n = \emptyset$ for $n = 1, 2, \dots$.

Proof. Only we prove (iv \rightarrow i). Let $\mathcal{U} = \{U_n : n \in \mathbb{N}\}$ be a countable X -open cover of Y . Define $W_n = \bigcup_{j \leq n} U_j$. Then the collection $\{W_n : n \in \mathbb{N}\}$ is an increasing countable X -open cover of Y , by (iv), there exists a Y -open cover $\{V_n : n \in \mathbb{N}\}$ of Y such that $\text{cl}_Y(V_n) \subseteq W_n$. Define $O_n = (U_n \cap Y) - \bigcup_{j < n} \text{cl}_Y(V_j)$. Then $\{O_n : n \in \mathbb{N}\}$ is an X -locally finite Y -open refinement of \mathcal{U} . \square

To identify more characterization of γ -countably paracompact we need the following theorem.

Theorem 3 Let Y be a γ -countably paracompact set in a space (X, τ) . If F is a generalized closed subset of (X, τ) such that $F \subseteq Y$, then F is γ -countably paracompact set in (X, τ) .

Proof. Let $\mathcal{U} = \{U_n : n \in \mathbb{N}\}$ be a countable X -open cover of F . Then $F \subseteq \bigcup_{n \in \mathbb{N}} U_n = U$. Since F is a generalized closed subset in (X, τ) and U is open in X , then $\text{cl}(F) \subseteq U$. Therefore, the collection $(X - \text{cl}(F)) \cup \{U_n : n \in \mathbb{N}\}$ is an X -open cover of the γ -countably paracompact set Y and so it has an X -locally finite open refinement, say \mathcal{V}^* . Put $\mathcal{V} = \{V \in \mathcal{V}^* : \exists U_V \in \mathcal{U} \text{ such that } V \subseteq U_V\}$. Finally, define $\mathcal{W} = \{V \cap F : V \in \mathcal{V}\}$. Then, it is clear that \mathcal{W} is X -locally finite and it is F -open refinement of \mathcal{U} since for each $V \in \mathcal{V}$ there exists an open O_V in (X, τ) such that $V = O_V \cap Y$ and so $V \cap F = O_V \cap Y \cap F = O_V \cap F$, which is open in F . \square

Corollary 2 If $F \subseteq Y \subseteq X$ such that Y is a γ -countably paracompact set and F is a closed set in (X, τ) . Then F is γ -countably paracompact set in (X, τ) .

Corollary 3 A closed subset of a countably paracompact space is γ -countably paracompact set.

Let $\{(X_\alpha, \tau_\alpha) : \alpha \in I\}$ be a collection of topological spaces such that $X_\alpha \cap X_\beta = \emptyset$ for each $\alpha \neq \beta$. Let $X = \bigcup_{\alpha \in I} X_\alpha$ be topologized by $\tau_s = \{G \subseteq X : G \cap X_\alpha \in \tau_\alpha \text{ for each } \alpha \in I\}$. Then (X, τ_s) is called the sum of the spaces $\{(X_\alpha, \tau_\alpha) : \alpha \in I\}$ and we write $X = \bigoplus_{\alpha \in I} X_\alpha$.

Theorem 4 *Let $A_\alpha \subseteq X$ for all $\alpha \in I$ and $A = \bigcup_{\alpha \in I} A_\alpha$. Then A is γ -countable paracompact set in X iff A_α is γ -countable paracompact set in X_α for all $\alpha \in I$.*

Proof. Let $\alpha \in I$ and \mathcal{U} be a countable X_α -open cover of A_α . Then the collection $\{\mathcal{U} : \mathcal{U} \in \mathcal{U}\} \cup (\bigcup_{\beta \neq \alpha} X_\beta)$ is a countable X -open cover of the γ -countable paracompact set A and so it has an X -locally finite A -open refinement, say \mathcal{V} . Put $\mathcal{V}_\mathcal{U} = \{V \cap A_\alpha : V \in \mathcal{V} \text{ and } V \subseteq \mathcal{U} \text{ for some } \mathcal{U} \in \mathcal{U}\}$. It is clear that $\mathcal{V}_\mathcal{U}$ is X_α -locally finite A_α -open collection such that $\mathcal{V}_\mathcal{U} < \mathcal{U}$. To show that $\mathcal{V}_\mathcal{U}$ is a cover for A_α . Let $x_\alpha \in A_\alpha$, then there exists $V \in \mathcal{V}$ such that $x_\alpha \in V$. Since $x_\alpha \notin X_\beta$ for all $\beta \neq \alpha$, then $V \subseteq \mathcal{U}$ for some $\mathcal{U} \in \mathcal{U}$ and $x_\alpha \in V \cap A_\alpha$. Conversely, Let \mathcal{U} be a countable X -open cover of A . For all $\alpha \in I$, the collection $\mathcal{U}_\alpha = \{\mathcal{U} \cap X_\alpha : \mathcal{U} \in \mathcal{U}\}$ is a countable X_α -open cover of the γ -countable paracompact set A_α in X_α , so it has an X_α -locally finite A_α -open refinement, say \mathcal{W}_α . For all $W \in \mathcal{W}_\alpha$, there exists an open set $H_{\alpha(W)}$ in X_α such that $W = A_\alpha \cap H_{\alpha(W)} = A \cap H_{\alpha(W)}$. Put $\mathcal{H} = \{W : W \in \mathcal{W}_\alpha, \alpha \in I\}$. Then, it is clear that \mathcal{H} is an A -open refinement of \mathcal{U} . To show that \mathcal{H} is X -locally finite, let $x \in X$. Then there exists $\alpha_0 \in I$ such that $x \in X_{\alpha_0}$ and $x \notin X_\beta$ for all $\beta \neq \alpha_0$. Since \mathcal{W}_{α_0} is X_{α_0} -locally finite, then there exists an open set K in X_{α_0} (and so in X) such that K is intersect at most finitely many numbers of \mathcal{W}_{α_0} and $K \cap W = \emptyset$ for all $W \in \mathcal{W}_\alpha, \alpha \neq \alpha_0$. Therefore, \mathcal{H} is X -locally finite and so A is γ -countable paracompact set in X . \square

Theorem 5 *Let $f : X \rightarrow Y$ be a perfect onto function and let B be a γ -countably paracompact set in the space (Y, σ) . Then $f^{-1}(B)$ is γ -countably paracompact set in (X, τ) .*

Proof. Let $\mathcal{U} = \{U_n : n \in \mathbb{N}\}$ be a countable X -open cover of $f^{-1}(B)$. For each $y \in B$, \mathcal{U} is an X -open cover of the compact set $f^{-1}(y)$, so there exists a finite subset $\{U_1, U_2, \dots, U_n\}$ of \mathcal{U} such that $f^{-1}(y) \subseteq \bigcup_{i=1}^n U_i = U_y$ and U_y is open in (X, τ) . Put $V_y = Y - f(X - U_y)$. Since f is closed then the collection $\mathcal{V} = \{V_y : y \in B\}$ is a countable Y -open cover of B , and so it has a Y -locally finite B -open refinement, say $\mathcal{W} = \{W_j : j = 1, 2, \dots\}$. Since f is continuous,

the family $f^{-1}(\mathcal{W}) = \{f^{-1}(W_j) : j = 1, 2, \dots\}$ is an X -locally finite $f^{-1}(B)$ -open cover of $f^{-1}(B)$ such that for each $j = 1, 2, \dots$ $f^{-1}(W_j) \subseteq U_{y_j}$ for some $y_j \in B$. Finally, the collection $\{f^{-1}(W_j) \cap U_i : j = 1, 2, \dots, i \in i_{y_j}\}$ is an X -locally finite $f^{-1}(B)$ -open refinement of \mathcal{U} . Therefore, $f^{-1}(B)$ is γ -countably paracompact. \square

An E_1 space [2] is a topological space such that every point is the intersection of a countable number of closed neighborhoods. Note that in [2] show that every E_1 space is T_2 .

Theorem 6 *Every γ -countable paracompact subset of E_1 space is closed.*

Proof. Let Y be a γ -countably paracompact subset of an E_1 space (X, τ) and let $x \notin Y$. Let $\{C_n : n \in \mathbb{N}\}$ be a countable family of closed neighborhoods of x such that $\{x\} = \bigcap C_n$. Now, $\{X - C_n : n \in \mathbb{N}\}$ is a countable X -open cover of Y and $x \notin \text{cl}(X - C_n)$ for any n . Hence there is an X -locally finite Y -open refinement of $\{X - C_n : n \in \mathbb{N}\}$, say \mathcal{W} . Put $H = \bigcup \{W : W \in \mathcal{W}\}$, then $\text{cl}(H) = \bigcup \{\text{cl}(W) : W \in \mathcal{W}\}$. Finally, put $H^* = X - \text{cl}(H)$. So H^* is open in (X, τ) such that $x \in H^*$ and $H^* \cap Y = \emptyset$. Therefore, $x \notin \text{cl}(Y)$ and Y is closed. \square

3 The relationship between α -countably paracompact, β -countably paracompact and γ -countably paracompact sets

In this section we study the relationship between α -countably paracompact, β -countably paracompact and γ -countably paracompact sets.

It follows from the definition that every α -countably paracompact set is γ -countably paracompact and every γ -countably paracompact set is β -countably paracompact. The following two examples show that the converse are not true in general.

Example 3 *Let $X = \mathbb{R}$ with the topology $\tau = \{U : \mathbb{R} - \mathbb{Q} \subseteq U\} \cup \{\emptyset\}$. Put $Y = \mathbb{Q}$. Then Y is γ -countably paracompact, note that if \mathcal{U} is a countable X -open cover of Y , then the collection $\{\{y\} : y \in Y\}$ is an X -locally finite Y -open refinement of \mathcal{U} . Now, to show Y is not α -countably paracompact, let $\mathcal{U} = \{(\mathbb{R} - \mathbb{Q}) \cup \{y\} : y \in Y\}$. Then \mathcal{U} is a countable X -open cover of Y . If \mathcal{V} is an X -locally finite X -open refinement of \mathcal{U} , then for every $y \in Y$ there exists $y \in V \in \mathcal{V}$ such that $y \in V \subseteq (\mathbb{R} - \mathbb{Q}) \cup \{y\}$. Thus, $V = (\mathbb{R} - \mathbb{Q}) \cup \{y\}$ which means \mathcal{V} is not X -locally finite.*

Example 4 Let $X = \mathbb{R}$ with the topology $\tau = \{U : 0 \notin U\} \cup \{\mathbb{R}\}$. Then $Y = \mathbb{Q}^* = \mathbb{Q} - \{0\}$ is β -countably paracompact, since $\tau_Y = \tau_{\text{dis}}$. On the other hand, Y is not γ -countably paracompact, since $\mathcal{U} = \{\{y\} : y \in Y\}$ is a countable X -open cover of Y by members of τ and it is not X -locally finite.

So what are the additional conditions that make the reversal of previous relationships true? This is what will be shown in the following Theorem.

Theorem 7 [3] Let Y be a closed β -countably paracompact set in a normal space. Then Y is α -countably paracompact

Theorem 8 Let Y be a γ -countably paracompact set in a space (X, τ) . Then Y is α -countably paracompact if one of the following holds:

- i. Y is closed in the normal space (X, τ) .
- ii. Y is open set in the space (X, τ) .

Proof. The proof of (ii) is clear. The proof of (i) follows by Theorem 7 and from the fact that every γ -countably paracompact set is β -countably paracompact. \square

Theorem 9 Let Y be a closed β -countably paracompact set in a space (X, τ) . Then Y is γ -countably paracompact.

Proof. Let Y be a closed β -countably paracompact subset of (X, τ) and let \mathcal{U} be a countable X -open cover of Y . Then the collection $\mathcal{W} = \{U \cap Y : U \in \mathcal{U}\}$ is a countable Y -open cover of Y and so it has a Y -locally finite Y -open refinement say \mathcal{V} . Since Y is closed set, by Proposition 2, \mathcal{V} is X -locally finite. Also as for every $V \in \mathcal{V}$, there exists $U \in \mathcal{U}$ such that $V \subseteq U \cap Y \subseteq U \in \mathcal{U}$, so \mathcal{V} is X -locally finite Y -open refinement of \mathcal{U} . Hence Y is γ -countably paracompact. \square

Corollary 4 Let Y be closed in a normal space (X, τ) . The following are equivalent:

- i. Y is γ -countably paracompact.
- ii. Y is α -countably paracompact.
- iii. Y is β -countably paracompact.

4 Locally γ -countably paracompact spaces

In this section we introduce locally γ -countably paracompact spaces and we study their properties.

Definition 5 A space (X, τ) is called locally γ -countably paracompact if each point $x \in X$ has an open neighborhood U in (X, τ) such that $\text{cl}(U)$ is γ -countably paracompact in (X, τ) .

The following result follow immediately from Theorem 9.

Proposition 7 Let (X, τ) be a space. Then (X, τ) is locally γ -countably paracompact iff for all $x \in X$ there exists an open neighborhood U in (X, τ) such that $\text{cl}(U)$ is β -countably paracompact.

Theorem 10 Every closed subspace of a locally γ -countably paracompact space is locally γ -countably paracompact.

Proof. Let F be a closed subspace of a locally γ -countably paracompact space (X, τ) . For every $x \in F$, there exists an open neighborhood U of the point x in the space (X, τ) such that $\text{cl}(U)$ is γ -countably paracompact space. The intersection $F \cap U$ is an open neighborhood of the point x in the subspace F and, by Corollary 3, $\text{cl}_F(F \cap U) = \text{cl}(F \cap U) \cap F = \text{cl}(F \cap U)$ is γ -countably paracompact, being a closed subset of the γ -countably paracompact set $\text{cl}(U)$, by Theorem 3. \square

Theorem 11 Every locally γ -countably paracompact E_1 space is T_3 .

Proof. Let F be a closed subset of a locally γ -countably paracompact space (X, τ) and $x \notin F$. Let $\text{cl}(P_x)$ be the γ -countably paracompact such that P_x is neighborhood of x and let $\{C_n : n \in \mathbb{N}\}$ be a countable family of closed neighborhood of x such that $\{x\} = \bigcap C_n$. Put $H = \text{cl}(P_x) \cap F$. Then, by Theorem 3, H is γ -countably paracompact set such that $x \notin H$. Thus the collection $\{X - C_n : n \in \mathbb{N}\}$ is a countable X -open cover of H and so it has an X -locally finite H -open refinement, say $\mathcal{U} = \{U_\alpha : \alpha \in \Delta\}$. Since H is closed in X , then $V = (\bigcup U_\alpha) \cup (X - \text{cl}(P_x))$ is an open set containing F such that $x \notin \text{cl}(V)$. Hence (X, τ) is regular. \square

Example 5 Let the Hausdorff neighborhoods of a point p in the Euclidean plane consist of open circles with p at the center excluding the points on the

vertical diameters except \mathfrak{p} itself. Since the resulting topology is a strengthening of the usual topology of the Euclidean plane it is an E_1 topology ([2], Example 2). Since this is a T_2 space which is not T_3 , it can not be locally γ -countably paracompact.

Lemma 1 *Let Y be an α -countably paracompact Lindelöf subset of a regular locally γ -countably paracompact space X . If W is an open set in (X, τ) such that $Y \subseteq W$, then there is an X -locally finite collection $\{F_n : n \in \mathbb{N}\}$ of closed γ -countably paracompact sets such that $Y \subseteq \bigcup_{n \in \mathbb{N}} \text{int}(F_n) \subseteq \bigcup_{n \in \mathbb{N}} F_n \subseteq W$.*

Proof. By the regularity of the space X , then for every $x \in Y$, there exists an open set U_x in X such that $x \in U_x \subseteq \text{cl}(U_x) \subseteq W$. On the other hand, X is locally γ -countably paracompact space and so there exists an open set H_x in X such that $\text{cl}(H_x)$ is γ -countably paracompact set. Put $V_x = \text{cl}(H_x) \cap \text{cl}(U_x)$. Then, by Theorem 3, V_x is a closed γ -countably paracompact set such that $x \in \text{int}(V_x) \subseteq W$. Therefore, the collection $\mathcal{V} = \{\text{int}(V_x) : x \in Y\}$ is an X -open cover of the Lindelöf set Y , so it has a countable subcover, say \mathcal{V}^* . Since Y is γ -countably paracompact set, then \mathcal{V}^* has an X -locally finite X -open refinement \mathcal{H} which cover Y . Now, for every $H \in \mathcal{H}$, $\text{cl}(H)$ is a closed set in X such that $\text{cl}(H) \subseteq V_x$ for some $x \in Y$ and so $\text{cl}(H)$, by Theorem 3, is γ -countably paracompact set. Thus, the collection $\{\text{cl}(H) : H \in \mathcal{H}\}$ is the required collection. \square

Theorem 12 *Let $f : (X, \tau) \rightarrow (Y, \sigma)$ be a perfect function from a space (X, τ) onto a locally γ -countably paracompact space (Y, σ) . Then (X, τ) is locally γ -countably paracompact.*

Proof. Let $x \in X$. Then there exists an open set V in (Y, σ) such that $f(x) \in V$ and $\text{cl}(V)$ is γ -countably paracompact in (Y, σ) . Now, by Theorem 5, $f^{-1}(\text{cl}(V))$ is γ -countably paracompact subset of X . Since $\text{cl}(f^{-1}(V)) \subseteq f^{-1}(\text{cl}(V))$, then by Theorem 3, $\text{cl}(f^{-1}(V))$ is γ -countably paracompact subset of X . \square

Corollary 5 *The product of a compact space (X, τ) and a locally γ -paracompact space (Y, σ) is locally γ -countably paracompact.*

Acknowledgment

The publication of this paper was supported by Yarmouk University Research council.

References

- [1] K. Y. Al-Zoubi, On γ –Paracompact Sets, *Questions and Answers in General Topology*, **34** (2016), 93–100.
- [2] C. E. Aull, A certain class of topological spaces, *Prace Matematyczne*, **11** (1967), 49–53.
- [3] C. E. Aull, *Paracompact subsets*, Proc. of the Second Prague Topological Symposium, Prague (1966), 45–51.
- [4] R. Engelking, *General topology*, Polish scientific publishers, Warszawa and New York, 1989.
- [5] N. Levine, Generalized closed sets in topology, *Rend. Circ. Mat. Palermo*, **19** (2) (1970), 89–96.

Received: February 19, 2020



On some Hall polynomials over a quiver of type \tilde{D}_4

Csaba Szántó

Babeş-Bolyai University, Cluj-Napoca
Faculty of Mathematics and
Computer Science
Str. Mihail Kogalniceanu nr. 1
RO-400084 Cluj-Napoca
Romania
email: szanto.cs@gmail.com

István Szöllősi

Babeş-Bolyai University, Cluj-Napoca
Faculty of Mathematics and
Computer Science
Str. Mihail Kogalniceanu nr. 1
RO-400084 Cluj-Napoca
Romania
email: szollosi@gmail.com

Abstract. Let k be an arbitrary field and Q a tame quiver of type \tilde{D}_4 . Consider the path algebra kQ and the category of finite dimensional right modules $\text{mod-}kQ$. We determine the Hall polynomials F_{xy}^z associated to indecomposable modules of defect $\partial z = -2$, $\partial x = \partial y = -1$ or dually $\partial z = 2$, $\partial x = \partial y = 1$.

1 Introduction

Classical Hall algebras associated with discrete valuation rings were introduced by Steinitz and Hall to provide an algebraic approach to the classical combinatorics of partitions. The multiplication is given by Hall polynomials which play an important role in the representation theory of the symmetric groups and the general linear groups. In 1990 Ringel defined Hall algebras for a large class of rings, namely finitary rings, including in particular path algebras of quivers over finite fields. Far reaching analogues of the classical ones, these Ringel-Hall algebras provided a new approach to the study of quantum groups using the representation theory of finite dimensional algebras. They

2010 Mathematics Subject Classification: 16G20

Key words and phrases: tame quiver, Hall polynomial

can also be used successfully in the theory of cluster algebras or to investigate the structure of the module category.

In case of Ringel-Hall algebras corresponding to Dynkin quivers and tame quivers we know due to Ringel and Hubery, that the structure constants of the multiplication are again polynomials in the number of elements of the base field. These are the generalized Hall polynomials. If we are looking at Hall polynomials associated to indecomposable modules, the classical ones are just 0 or 1, the generalized ones in the Dynkin case are also known and have degree up to 5, however we do not have too much information about the generalized ones in the tame case. The first lists of particular tame Hall polynomials were given by the authors in [6] and in [7]. In [6] we presented all the tame Hall polynomials associated to indecomposable modules of defect $-1, 0, 1$. In [7] we listed the tame Hall polynomials corresponding to exact sequences of the form $0 \rightarrow P \rightarrow R \rightarrow I \rightarrow 0$, where P is a preprojective, I a preinjective indecomposable and R is a homogeneous module of dimension δ (the minimal radical vector of the tame quiver).

In this paper we restrict ourselves to the tame quiver of type \tilde{D}_4 and determine all the tame Hall polynomials F_{xy}^z associated to indecomposable modules of defect $\partial z = -2$, $\partial x = \partial y = -1$ or dually $\partial z = 2$, $\partial x = \partial y = 1$.

2 Preliminaries

We begin with some facts related to representations of tame quivers. For a detailed description we refer to [1, 2, 3].

Let $Q = (Q_0, Q_1)$ be a tame quiver without oriented cycles. Suppose that the vertex set Q_0 has n elements and for an arrow $\alpha \in Q_1$ we denote by $t(\alpha), h(\alpha) \in Q_0$ the tail and head of α . The Euler form of Q is a bilinear form on $\mathbb{Z}Q_0 \cong \mathbb{Z}^n$ given by $\langle x, y \rangle = \sum_{i \in Q_0} x_i y_i - \sum_{\alpha \in Q_1} x_{t(\alpha)} y_{h(\alpha)}$. Its quadratic form q_Q (called Tits form) is independent from the orientation of Q and in the tame case it is positive semidefinite with radical $\mathbb{Z}\delta$, where δ is a minimal positive imaginary root of the corresponding Kac-Moody root system (which is also the minimal radical vector of the Tits form). The defect of $x \in \mathbb{Z}Q_0$ is then $\partial x = \langle \delta, x \rangle$.

Let k be a field. The category $\text{mod-}kQ$ will be identified with the category $\text{rep-}kQ$ of the finite dimensional k -representations of the quiver. We will denote by $[M]$ the isomorphism class of the module M , by α_M the number of its automorphisms, by $\dim M \in \mathbb{Q}_0 \cong \mathbb{Z}^n$ its dimension vector and by $\partial M =$

$\partial(\dim M)$ its defect. Using the Euler form one has for $X, Y \in \text{mod-}kQ$

$$\langle \underline{\dim} X, \underline{\dim} Y \rangle = \dim_k \text{Hom}(X, Y) - \dim_k \text{Ext}^1(X, Y).$$

For $\dim_k \text{Hom}(X, Y)$ we will use the notation (X, Y) .

The indecomposable modules in $\text{mod-}kQ$ are of three types: preprojectives (having negative defect), preinjectives (having positive defect) and regulars (having zero defect).

For P preprojective (i.e. with all its indecomposable components preprojective), I preinjective and R regular module we have $\text{Hom}(R, P) = \text{Hom}(I, P) = \text{Hom}(I, R) = \text{Ext}^1(P, R) = \text{Ext}^1(P, I) = \text{Ext}^1(R, I) = 0$. It follows that the submodules of a preprojective module are always preprojective, preinjectives can project only on preinjectives, a submodule of a regular module cannot have preinjective components and a regular cannot project on preprojectives. Preprojective and preinjective indecomposables are exceptional (i.e. their endomorphism space is one dimensional and they have no self extensions) and are uniquely determined up to isomorphism by their dimension vector, which is a positive real root of the root system of Q . Note also that the possible defects of a preprojective indecomposable are -1 in the \tilde{A}_n case, $-1, -2$ in the \tilde{D}_n case, $-1, -2, -3$ in the \tilde{E}_6 case, $-1, -2, -3, -4$ in the \tilde{E}_7 case and $-1, -2, -3, -4, -5, -6$ in the \tilde{E}_8 case.

The category of regular modules is an abelian, exact subcategory which decomposes into a direct sum of serial categories with Auslander-Reiten quiver of the form $\mathbb{Z}\tilde{A}_\infty/\mathfrak{m}$, called tubes of rank \mathfrak{m} . These tubes are indexed by the points of the projective line \mathbb{P}_k^1 , the degree of a point $\mathfrak{a} \in \mathbb{P}_k^1$ being denoted by $\deg \mathfrak{a}$. A tube of rank 1 is called homogeneous, otherwise it is called non-homogeneous. We have at most 3 non-homogeneous tubes indexed by points \mathfrak{a} of degree $\deg \mathfrak{a} = 1$. All the other tubes are homogeneous. We assume that the non-homogeneous tubes are labelled by some subset of $\{0, 1, \infty\}$, whereas the homogeneous tubes are labelled by the closed points of the scheme $\mathbb{H}_k = \mathbb{H}_{\mathbb{Z}} \otimes k$ for some open integral subscheme $\mathbb{H}_{\mathbb{Z}} \subset \mathbb{P}_{\mathbb{Z}}^1$. Let $\mathbb{X}_k \subseteq \mathbb{H}_k$ be the set of points of degree 1. The indecomposables on a homogeneous tube labelled by $\mathfrak{a} \in \mathbb{H}_k$ are denoted by $R^k(1, \mathfrak{a}) \subset R^k(2, \mathfrak{a}) \subset \dots$. For a partition $\lambda = (\lambda_1, \dots, \lambda_n)$ let $R^k(\lambda, \mathfrak{a}) = R^k(\lambda_1, \mathfrak{a}) \oplus \dots \oplus R^k(\lambda_n, \mathfrak{a})$. Note that the homogeneous modules of dimension δ are up to isomorphism $R^k(1, \mathfrak{a})$, with $\mathfrak{a} \in \mathbb{X}_k$. For simplicity we will denote them by $R^k(\mathfrak{a})$. Note that $\dim_k \text{End}(R^k(\mathfrak{a})) = 1$.

A module without homogeneous regular components can be described combinatorially, field independently, using a system of positive real roots together with the dimension of quasi-socles for the non-homogeneous regular components of dimension $t\delta$. We denote this system by μ and let $M(\mu, k)$ be the cor-

responding (up to isomorphism) unique module in $\text{mod-}kQ$. A Segre symbol is a multiset $\sigma = \{(\lambda^1, d_1), \dots, (\lambda^r, d_r)\}$, where λ^i are partitions and $d_i \in \mathbb{N}^*$. It will describe the homogeneous regular components of the module. Using the definitions above, a decomposition symbol is pair $\alpha = (\mu, \sigma)$. Given a decomposition symbol $\alpha = (\mu, \sigma)$ and a field k , we define the decomposition class $S(\alpha, k)$ to be the set of isomorphism classes of modules of the form $M(\mu, k) \oplus R$, where $R = R^k(\lambda^1, a_1) \oplus \dots \oplus R^k(\lambda^r, a_r)$ for some distinct points $a_1, \dots, a_r \in \mathbb{H}_k$ such that $\deg a_i = d_i$. We also mention that for a decomposition symbol α the polynomial $n_\alpha(q) = |S(\alpha, k)|$ is strictly increasing in $q > 1$.

Note that for k finite with q elements $|\mathbb{X}_k| = q + 1$, q or $q - 1$ in the \tilde{A}_n case and $q - 2$ for other tame quivers. So if k has 2 elements and the quiver is not of \tilde{A}_n type there are no homogeneous modules of dimension δ .

For simplicity denote by x the decomposition symbol corresponding to a preprojective (preinjective) indecomposable given by the root x . Also denote by δ the symbol corresponding to homogeneous modules of dimension δ .

We mention next some needed facts about Ringel-Hall algebras. Suppose that k is finite. We consider the rational Ringel-Hall algebra $\mathcal{H}(kQ)$ of the algebra kQ . Its \mathbb{Q} -basis is formed by the isomorphism classes $[M]$ from $\text{mod-}kQ$ and the multiplication is defined by $[N_1][N_2] = \sum_{[M]} F_{N_1 N_2}^M [M]$. The structure constants $F_{N_1 N_2}^M = |\{\mathbf{U} \subseteq M \mid \mathbf{U} \cong N_2, M/\mathbf{U} \cong N_1\}|$ are called Ringel-Hall numbers. The associativity of the Ringel-Hall algebra follows from the equality $\sum_{[N]} F_{N_1 N}^M F_{N N_3}^N = \sum_{[N]} F_{N_1 N_2}^N F_{N N_3}^M$.

Hubery proved the existence of generalized Hall polynomials in tame cases with respect to the decomposition classes.

Theorem 1 ([4]) *Given decomposition symbols α, β and γ , there exists a rational polynomial $F_{\alpha\beta}^\gamma$ such that for any finite field k with $|k| = q$,*

$$F_{\alpha\beta}^\gamma(q) = \sum_{\substack{A \in S(\alpha, k) \\ B \in S(\beta, k)}} F_{AB}^C \quad \text{for all } C \in S(\gamma, k)$$

and moreover

$$n_\gamma(q) F_{\alpha\beta}^\gamma(q) = n_\alpha(q) \sum_{\substack{B \in S(\beta, k) \\ C \in S(\gamma, k)}} F_{AB}^C \quad \text{for all } A \in S(\alpha, k),$$

$$n_\gamma(q) F_{\alpha\beta}^\gamma(q) = n_\beta(q) \sum_{\substack{A \in S(\alpha, k) \\ C \in S(\gamma, k)}} F_{AB}^C \quad \text{for all } B \in S(\beta, k).$$

Remark 1 The polynomials F_{rx}^z or F_{yr}^z where r is the symbol of a homogeneous regular will denote in our article Hubery's polynomial divided by $n_r(q)$, which is again a polynomial.

We list now the known tame Hall polynomials associated to indecomposables (see the introduction).

Proposition 1 ([6, 7]) *We have the following:*

a) Suppose we limit ourselves to defects in $\{-1, 0, 1\}$. For two roots x, y with $\partial x = \partial y = -1$ and $\langle x, y \rangle > 0$ we have that $F_{rx}^y = 1$ for any symbol r corresponding to regular indecomposables of dimension $y - x$. This dualizes for roots with defect 1. For roots x, y with $\partial x = -1$, $\partial y = 1$ and $\langle x, y \rangle \neq 0$ we have that $F_{yx}^r = \frac{1}{q-1} \alpha_r$ for any symbol r corresponding to regular indecomposables of dimension $y - x$ (where α_r is the number of automorphisms). For three symbols corresponding to regular indecomposables the Hall polynomial is classical so it is 0 or 1. In all the other cases the Hall polynomial is 0.

b) Let x be a positive real root with $\partial x < 0$. Then $F_{\delta-xx}^\delta = h_{-\partial x}$, where

$$\begin{aligned} h_1 &= 1, \\ h_2 &= q - 3, \\ h_3 &= q^2 - 5q + 7, \\ h_4 &= q^3 - 6q^2 + 15q - 14, \\ h_5 &= q^4 - 7q^3 + 22q^2 - 37q + 26, \\ h_6 &= q^5 - 7q^4 + 22q^3 - 45q^2 + 62q - 39. \end{aligned}$$

We end this section with a well known lemma:

Lemma 1 Let P and P' be preprojective indecomposables with $\partial P = -1$. Then every nonzero morphism $f: P \rightarrow P'$ is a monomorphism.


3 Reductions

From now on we suppose that Q is of \tilde{D}_4 type.

Our aim is to determine the tame Hall polynomials F_{xy}^z associated to indecomposable modules of defect $\partial z = -2$, $\partial x = \partial y = -1$ or dually $\partial z = 2$, $\partial x = \partial y = 1$.

Using reflection functors (and the fact that Hall numbers are preserved via these functors) one can see that we only need to consider a particularly oriented quiver of \tilde{D}_4 type (see for example [6] for all the details).

By the arguments above we will consider the quiver Q' of \tilde{D}_4 type with all arrows pointing to a non-central vertex (say vertex 1, the central vertex being 5). Thus the unique sink in Q' is 1 (one of the marginal vertexes):



We end this section with the main tool, which will provide us the recursions permitting to compute the Ringel-Hall numbers above.

Proposition 2 [5] *Let $X, Y, Z \in \text{mod-}kQ$ where Q is an arbitrary quiver and k is finite. Denote by s_X^Y the number of submodules of Y isomorphic to X , by f_X^Y the number of submodules of Y with factor isomorphic to X , by e_X^Y the number of epimorphisms from Y to X , by α_X the number of automorphisms of X and by h_{XY} the number of morphisms from X to Y . Then we have the following formula:*

$$e_X^Y = h_{YX} - \sum_{\substack{Z \in \text{mod-}kQ \\ \dim Z < \dim X}} f_Z^Y \alpha_Z s_Z^X.$$

Moreover $e_X^Y = \alpha_X f_X^Y$.


4 Recursions and Hall polynomials

Consider the quiver Q' and the indecomposable preprojectives P_0, P', P with $\dim P = \dim P_0 + \dim P'$ and $\partial P_0 = \partial P' = -1, \partial P = -2$. Let S_1 be the projective simple corresponding to the unique sink 1. The rest of the indecomposable preprojectives are:

- $P_2(n)$ the indecomposable preprojective (of defect -2) with dimension vector $(1, 0, 0, 0, 1) + n\delta$;

- $P_1^i(n)$ (for $i = \overline{1,4}$) the indecomposable preprojectives (of defect -1) of dimensions $(0, 0, 0, 0, 1) + n\delta$, $(1, 1, 0, 0, 1) + n\delta$, $(1, 0, 1, 0, 1) + n\delta$, $(1, 0, 0, 1, 1) + n\delta$;
- $P_1^{2i}(n)$ (for $i = \overline{1,4}$) the indecomposable preprojectives (of defect -1) of dimensions $(2, 1, 1, 1, 2) + n\delta$, $(1, 0, 1, 1, 2) + n\delta$, $(1, 1, 0, 1, 2) + n\delta$, $(1, 1, 1, 0, 2) + n\delta$.

The segment of the preprojective component of the Auslander-Reiten quiver which we will use is the following:



Proposition 3 $F_{P/P_0}^P = g_{n-1}(q)$, where $n = \langle \underline{\dim} P_0, \underline{\dim} P \rangle = \langle \underline{\dim} P, \underline{\dim} P' \rangle$ and

$$g_n = X^n - 3X^{n-1} + \cdots + (-1)^{n-1}(2n-1)X + (-1)^n(n+1)$$

(with $g_0 = 1$ and $g_{-1} = 0$).

Proof. First of all note that $n = \langle \underline{\dim} P_0, \underline{\dim} P \rangle = \langle \underline{\dim} P_0, \underline{\dim} P_0 + \underline{\dim} P' \rangle = 1 + \langle \underline{\dim} P_0, \underline{\dim} P' \rangle = \langle \underline{\dim} P_0 + \underline{\dim} P', \underline{\dim} P' \rangle = \langle \underline{\dim} P, \underline{\dim} P' \rangle$. Also if $n = \langle \underline{\dim} P_0, \underline{\dim} P \rangle = (P_0, P) = 0$, then $F_{P/P_0}^P = 0 = g_{-1}$.

We will use induction on $n \geq 1$. For $n = 1$, the assertion is trivial since $n = 1 = \langle \underline{\dim} P_0, \underline{\dim} P \rangle = (P_0, P)$. Using successive Auslander-Reiten translations, the fact that the modules are indecomposable preprojectives and $\underline{\dim} P = \underline{\dim} P_0 + \underline{\dim} P'$, one can see that $F_{P/P_0}^P = F_{P_1 S_1}^{P_2}$, where $n = 1 = \langle \underline{\dim} P_0, \underline{\dim} P \rangle = \langle \underline{\dim} S_1, \underline{\dim} P_2 \rangle = (\underline{\dim} P_2)_1$. This means (looking at the dimensions) that $P_2 = P_2(0)$, and $P_1 = P_1^1(0)$, so $F_{P/P_0}^P = F_{P_1 S_1}^{P_2} = 1$.

Suppose the assertion is true for values under n and prove it for n .

Using again successive Auslander-Reiten translations, one can see (as above) that $F_{P/P_0}^P = F_{P_1 S_1}^{P_2}$, where $n = \langle \underline{\dim} P_0, \underline{\dim} P \rangle = \langle \underline{\dim} S_1, \underline{\dim} P_2 \rangle = (\underline{\dim} P_2)_1$.

By Proposition 2 we have that

$$e_{P_1}^{P_2} = h_{P_2 P_1} - \sum_{\substack{Z \in \text{mod-}kQ' \\ \dim Z < \dim P_1}} f_Z^{P_2} \alpha_Z s_Z^{P_1}.$$

Note that $F_{P_1 S_1}^{P_2} = f_{P_1}^{P_2} = \frac{e_{P_1}^{P_2}}{\alpha_{P_1}} = \frac{e_{P_1}^{P_2}}{q-1}$ and $h_{P_2 P_1} = q^{(P_2, P_1)}$. Also if there is a monomorphism $Z \rightarrow P_1$ and an epimorphism $P_2 \rightarrow Z$ it follows that $Z = 0$ or Z is a indecomposable preprojective of defect -1 such that $(Z, P_1) \neq 0$ and $(P_2, Z) \neq 0$ (here we use the fact that submodules of preprojectives are preprojective and a preprojective of defect -2 can't project on a different preprojective of defect -2). Using the fact that the indecomposable preprojectives are directing, one can see that in the Auslander-Reiten quiver Z follows after P_2 and precedes P_1 .

Suppose $n = 2m$. Denote by $g'_{2m} = f_{P_1^{11}(2m)}^{P_2(2m)}$.

Using the previous formula and observations and the Auslander-Reiten segment presented above, performing all the calculations we obtain:

$$\begin{aligned} g'_{2m} &= f_{P_1^{11}(2m)}^{P_2(2m)} \\ &= \frac{q^{2m+1} - 1}{q - 1} - \sum_{\substack{i=\overline{1,4} \\ j=0, m-1}} f_{P_1^{1i}(m+j)}^{P_2(2m)} s_{P_1^{1i}(m+j)}^{P_1^{11}(2m)} - \sum_{\substack{i=\overline{1,4} \\ j=0, m-1}} f_{P_1^{2i}(m+j)}^{P_2(2m)} s_{P_1^{2i}(m+j)}^{P_1^{11}(2m)} \quad (1) \end{aligned}$$

By Lemma 1 we have that

$$s_{P_1^{1i}(m+j)}^{P_1^{11}(2m)} = \frac{q^{(P_1^{1i}(m+j), P_1^{11}(2m))} - 1}{q - 1}$$

where $(P_1^{11}(m+j), P_1^{11}(2m)) = m - j + 1$ and $(P_1^{1i}(m+j), P_1^{11}(2m)) = m - j$ for $i = \overline{2, 4}$. Also

$$s_{P_1^{2i}(m+j)}^{P_1^{11}(2m)} = \frac{q^{(P_1^{2i}(m+j), P_1^{11}(2m))} - 1}{q - 1}$$

where $(P_1^{21}(m+j), P_1^{11}(2m)) = m - j - 1$ and $(P_1^{2i}(m+j), P_1^{11}(2m)) = m - j$ for $i = \overline{2, 4}$.

The kernel of an epimorphism $P_2(2m) \rightarrow P_1^{1i}(m+j)$ is preprojective and of defect -1 , so it is indecomposable and unique. Denote it by X . This implies that $f_{P_1^{1i}(m+j)}^{P_2(2m)} = F_{P_1^{1i}(m+j)X}^{P_2(2m)}$. Using the induction hypothesis one can deduce that $F_{P_1^{1i}(m+j)X}^{P_2(2m)} = g_{2j}(q)$ and $F_{P_1^{2i}(m+j)X}^{P_2(2m)} = g_{2j+1}(q)$, since $\langle P_2(2m), P_1^{1i}(m+j) \rangle = 2j + 1$ and $\langle P_2(2m), P_1^{2i}(m+j) \rangle = 2j + 2$.

Substituting everything in (1) we obtain:

$$\begin{aligned} g'_{2m} &= \frac{q^{2m+1} - 1}{q - 1} - \frac{q^{m+1} - 1}{q - 1} g_0(q) \\ &\quad - \sum_{j=1, m-1} \frac{q^{m-j+1} - 1}{q - 1} (g_{2j}(q) + 3g_{2j-1}(q) + 3g_{2j-2}(q) + g_{2j-3}(q)) \\ &\quad - 3g_{2m-1}(q) - 3g_{2m-2}(q) - g_{2m-3}(q). \end{aligned}$$

In case $n = 2m + 1$ a similar recursion can be obtained for g'_{2m+1} . More precisely we get:

$$\begin{aligned} g'_{2m+1} &= \frac{q^{2m+2} - 1}{q - 1} - \frac{q^{m+1} - 1}{q - 1} (g_1(q) + 3g_0(q)) \\ &\quad - \sum_{j=1, m-1} \frac{q^{m-j+1} - 1}{q - 1} (g_{2j+1}(q) + 3g_{2j}(q) + 3g_{2j-1}(q) + g_{2j-2}(q)) \\ &\quad - 3g_{2m}(q) - 3g_{2m-1}(q) - g_{2m-2}(q). \end{aligned}$$

By direct calculation we get that $g'_{2m} = g_{2m}(q)$ and $g'_{2m+1} = g_{2m+1}(q)$ that is, $g'_n = g_n(q)$ for all n . \square

Remark 2 Based on calculations done with a computer we conjecture that the polynomials above are irreducible (as integer polynomials).

Acknowledgment

Supported by a grant of the Babeş-Bolyai University (GTC 2019-2020).

References

- [1] I. Assem, D. Simson, A. Skowronski, *Elements of Representation Theory of Associative Algebras, Volume 1: Techniques of Representation Theory*, LMS Student Texts (No. 65), Cambridge Univ. Press 2006.
- [2] M. Auslander, I. Reiten, S. Smalø, *Representation Theory of Artin Algebras*, Cambridge Stud. in Adv. Math. 36, Cambridge Univ. Press 1995.
- [3] V. Dlab, C. M. Ringel, *Indecomposable representations of graphs and algebras*, AMS Memoirs 173, 1976.

- [4] A. Hubery, *Hall polynomials for affine quivers*, Represent. Theory, **14** (2010), 355–378.
- [5] C. M. Ringel, *Hall algebras*, in: Topics in algebra, Banach Center Publ., **26** (1990), 433–447.
- [6] Cs. Szántó, On some Ringel-Hall products in tame cases, *Journal of Pure and Applied Algebra*, **216** (2012), 2069–2078.
- [7] Cs. Szántó, I. Szöllősi, Hall polynomials and the Gabriel-Roiter submodules of simple homogeneous modules, *Bulletin of the London Mathematical Society*, **47** (2), (2015), 206–216.

Received: October 27, 2020

Contents

Volume 12, 2020

V. K. Bhat, P. Singh, S. Sharma

On weak (σ, δ) -rigid rings over Noetherian rings.....5

L. Bognár, A. Joós, B. Nagy

Assessing the effect size of users' consciousness for computer networks vulnerability 14

S. S. Dragomir

Trapezoid type inequalities for generalized Riemann-Liouville fractional integrals of functions with bounded variation 30

W. Fechner, Á. Szász

Composition iterates, Cauchy equations, translation equations, and Sincov inclusions 54

Z. Finta

Direct and converse theorems for King operators 85

V. Gupta, A. H. Ansari, N. Mani

Fixed point theorem for new type of auxiliary functions97

A. Gutierrez

Collatz conjecture revisited: an elementary generalization 112

A. Keddi, F. Madani, A. A. Bouchentouf

Nonparametric estimation of trend function for stochastic differential equations driven by a bifractional Brownian motion128

M. Lee

**Orbital shadowing property on chain transitive sets for
generic diffeomorphisms 146**

A. J. Mathias, V. Sangeetha, M. Acharya

Restrained domination in signed graphs 155

V. A. Minuță

Graded Morita theory over a G -graded G -acted algebra 164

V. Popa, A.-M. Patrîciu

**Fixed points for a pair of weakly compatible mappings satisfying
a new type of ϕ - implicit relation in S - metric spaces 179**

G. Román

On Euler products with smaller than one exponents 193

B. Roy

More on decomposition of generalized continuity 212

A. H. Ansari, M. S. Khan, V. Rakočević

Maia type fixed point results via C -class function 227

P. Baliarsingh, L. Nayak, S. Samantaray

**On the convergence difference sequences and the related operator
norms 245**

D. Bród

On some properties of split Horadam quaternions 260

J. Bukor, J. T. Tóth

On topological properties of the set of maldistributed sequences 272

C. M. da Fonseca

On the connection between tridiagonal matrices, Chebyshev polynomials, and Fibonacci numbers 280

P. V. Danchev

A note on nil-clean rings 287

H. Ö. Güney, S. Owa

Generalized operator for Alexander integral operator 294

S. F. Namiq, E. Rosas

On $\lambda^D - R_0$ and $\lambda^D - R_1$ spaces 307

Ş. Özlü, A. Sezgin

Soft covered ideals in semigroups 317

M. Shahraki, S. Sedghi, S. M. A. Aleomraninejad, Z. D. Mitrović

Some fixed point results on S-metric spaces 347

V. Soleymanivarniab, R. Nikandish, A. Tehranian

On the metric dimension of strongly annihilating-ideal graphs of commutative rings 358

V. Agnes S. J. Lavanya, M. P. Jeyaraman, H. Aaisha Farzana

On application of differential subordination for Carathéodory functions 370

A. Rawshdeh, H. H. Al-Jarrah, K. Y. Al-Zoubi, W. A. Shatanawi

On γ -countably paracompact sets 383

Cs. Szántó, I. Szöllősi

On some Hall polynomials over a quiver of type \tilde{D}_4 395

Acta Universitatis Sapientiae

The scientific journal of Sapientia Hungarian University of Transylvania publishes original papers and surveys in several areas of sciences written in English.

Information about each series can be found at

<http://www.acta.sapientia.ro>.

Main Editorial Board

László DÁVID Editor-in-Chief

Adalbert BALOG Executive Editor

Angella SORBÁN Managing Editor

Csaba FARKAS Member

Zoltán KÁSA Member

Laura NISTOR Member

Ágnes PETHŐ Member

Acta Universitatis Sapientiae, Informatica

Editorial Board

Executive Editors

Csaba FARKAS (Sapientia Hungarian University of Transylvania, Romania)

Emil HOROBET (Sapientia Hungarian University of Transylvania, Romania)

Pál KUPÁN (Sapientia Hungarian University of Transylvania, Romania)

Róbert SZÁSZ (Sapientia Hungarian University of Transylvania, Romania)

Members

Sébastien FERENCZI (Institut de Mathématiques de Luminy, France)

Kálmán GYÖRY (University of Debrecen, Hungary)

Zoltán MAKÓ (Sapientia Hungarian University of Transylvania, Romania)

Ladislav MIŠÍK (University of Ostrava, Czech Republic)

János TÓTH (Selye University, Slovakia)

Adrian PETRUŞEL (Babeş-Bolyai University, Romania)

Alexandru HORVÁTH (Sapientia Hungarian University of Transylvania, Romania)

Árpád BARICZ (Babeş-Bolyai University, Romania)

Csaba SZÁNTÓ (Babeş-Bolyai University, Romania)

Szilárd ANDRÁS (Babeş-Bolyai University, Romania)

Each volume contains two issues.



Sapientia University



Sciencido by De Gruyter

sciencido



Scientia Publishing House

ISSN 1844-6094

<http://www.acta.sapientia.ro>

Information for authors

Acta Universitatis Sapientiae, Mathematica publishes original papers and surveys in various fields of Mathematics. All papers are peer-reviewed.

Papers published in current and previous volumes can be found in Portable Document Format (pdf) form at the address: <http://www.acta.sapientia.ro>.

The submitted papers should not be considered for publication by other journals. The corresponding author is responsible for obtaining the permission of coauthors and of the authorities of institutes, if needed, for publication, the Editorial Board is disclaiming any responsibility.

Submission must be made by email (acta-math@acta.sapientia.ro) only, using the L^AT_EX style and sample file at the address <http://www.acta.sapientia.ro>. Beside the L^AT_EX source a pdf format of the paper is necessary too.

Prepare your paper carefully, including keywords, 2010 Mathematics Subject Classification (MSC 2010) codes (<http://www.ams.org/msc/msc2010.html>), and the reference citation should be written in brackets in the text as [3]. References should be listed alphabetically based on the Instructions for Authors given at the address <http://www.acta.sapientia.ro>.

Contact address and subscription:

Acta Universitatis Sapientiae, Informatica
RO 400112 Cluj-Napoca, Str. Matei Corvin nr. 4.
Email: acta-math@acta.sapientia.ro

Printed by F&F INTERNATIONAL
Director: Enikő Ambrus

ISSN 1844-6094
<http://www.acta.sapientia.ro>

Supported by:

