

# GEOLOGICA HUNGARICA

FASCICULI INSTITUTI GEOLOGICI HUNGARIAE  
AD ILLUSTRANDAM NOTIONEM GEOLOGICAM  
ET PALAEONTOLOGICAM

## SERIES PALAEONTOLOGICA

FASCICULUS 42

DR. MÜLLER PÁL: A BÁDENI EMELET TÍZLÁBÚ RÁKJAI  
DR. PÁL MÜLLER: DECAPOD CRUSTACEA OF THE BADENIAN

INSTITUTUM GEOLOGICUM HUNGARICUM  
BUDAPESTINI, NOVEMBER 1984

Lektor:  
DR. DUDICH ENDRE

Szakszerkesztő:  
PIROS CHRISTA

Az angol szöveg a szerző munkája

Nyelvi lektor:  
DR. MIKLÓS MÜLLER

Technikai szerkesztő:  
CSABAY LÁSZLÓNÉ  
GARAMSZEGI KÁROLYNÉ

HU ISSN 03 67 4150

A kézirat lezárva 1984. január 15-én

Megjelent a Műszaki Könyvkiadó gondozásában  
Budapest, 1984

Felelős kiadó: DR. HÁMOR GÉZA

Műszaki vezető: Kórizs Károly  
Műszaki szerkesztő: Metzker Sándor

A könyv formátuma: A4

Terjedelem: 40 A5 ív

Példányszám: 880

Ábrák száma: 12 + fényképtáblák

Betűsalád és -méret: Extended, gm/gm

Papír minősége: 80 g ofszet és 120 g műnyomó

Azonossági szám: 01486



84.2339 Egyetemi Nyomda, Budapest, 1984  
Felelős vezető: Sümeghi Zoltán igazgató

## TARTALOM — CONTENTS

Bevezetés .....	5
A bádeni emelet és a Középső Paratethys .....	6
A bádeni emelet rákjaival foglalkozó irodalom áttekintése .....	7
A bádeni emelet rákjainak gyűjteményi lelőhelyei .....	8
Munkamódszerek, preparálás .....	9
A fontosabb lelőhelyek leírása .....	10
Taxonómiai alapelvek .....	14
A bádeni tízlábú rák fauna jellegzetességei .....	16
A bádeni rákok rétegtani értéke .....	17
Ősföldrajzi következtetések .....	19
Ökológiai kérdések .....	20
* * *	
Summary .....	27
Introduction .....	28
Acknowledgements .....	28
The Badenian stage and the Central Paratethys .....	30
Previous studies on Badenian decapods .....	32
Depositories of Badenian decapods .....	34
Methods used in field-work, preparation of the material .....	35
Description of the localities .....	36
Taxonomic principles .....	48
Systematical part .....	49
General characteristics of the Badenian decapod fauna .....	100
Stratigraphic significance of Badenian decapods .....	102
Palaeogeographic implications .....	105
Ecology .....	107
Addendum .....	112
References .....	113
List of species .....	118
Plates .....	123

A tízlabú rákok az ósmaradványanyagban árnylag ritkák, pedig a mai tengeri élővilág fontos, gyakori tagjai. Kevés olyan lelőhely van, ahol húsznál több fajt találtak, ezek nagy része is meozós vagy eocén. A neogén nagy kövületgazdagsága ellenére többnyire csak szórványleleteket adott. A Középső Paratethys bádeni emelete kivételes. Ebben eddig 120 különböző alakra bukkantak, ezek közül 102-t lehetett fajra vagy alfajra meghatározni. A gazdagság egyik oka a viszonylag keskeny tengerágakban, tehát jórészt nyugodt körülmények között lerakódott mészkövek nagy mennyisége. Talán ennél is fontosabb, hogy sok jó lelőhely olyan világvárosok közelében (Budapest és Bécs) van, ahol az őslénytani kutatásnak nagy a hagyománya. A kövült rákok vizsgálatának is több mint egy évszázados múltja van itt. A régi közlemények revíziójának időszerű volta mellett a szerző által összegyűjtött anyag is indokolta e munka elkészítését.

A szétszórt lelőhelyekről származó szórványleletek néha gyönyörű példányokat adtak, ezekkel foglalkozik a régebbi közlemények jó része. Ennél fontosabbnak ítélem azt, hogy egyes rétegek alaposan begyűjtött anyagából, számos faj alapján, az egykor együttesekre is tudunk következtetni. Ennek az elvnek alapján néhány igéretes lelőhelyet sokszor, egyeseket (Rákos, Örs vezér tere, Törökmező, Tétényi-fennsík) száznál is többször meglátogattam, vagy viszonylag rövid idő alatt nagy mennyiségű kőzetanyagot gyűjtöttem, amit aztán később vizsgáltam át (Gross-Höflein). Más, kevésbé igéretes vagy távolabbi helyeket csak egy-egy, vagy néhány alkalommal vizsgáltam (pl. Mátraverebély—Szentkút, Zebegény, Szob, Deutsch-Altenburg, Grobie). Ezért az összegyűlt anyag nemileg egyoldalú. A budapesti felső-bádeni lelőhelyekről sokszorta több anyag gyűlt össze, mint a távolabbi, alsó-bádeni rétegek jó részéből. Ennek ellenére érdemes volt az egész ismert bádeni anyagot egységen feldolgozni.

Több ezer példány (izolált ollók, carapaxok, más testrészek, ritkán teljes vázak) képezi a monográfia alapját. A 102 meghatározható alak között 21 faj és egy alfaj újnak bizonyult, s négy fajt nem lehetett ismert nemzettségebe sorolni, ezért új genusként írtam le ezeket.

Célom az volt, hogy leírjam vagy revideáljam az összes eddig publikált, gyűjteményekben elérhető, vagy újonnan gyűjtött tízlabú rákot, az elkülönült ujjak kivételével. Az ujjak vizsgálata mindaddig nem ígér sok eredményt, amíg a carapax vagy olló alapján leírt rákokkal nem sikerül azokat azonosítani. Ezért az ujjakat csak akkor vizsgáltam, ha az adott réteg más, felismerhető maradványt is tartalmazott, s az ujjaknak azokkal való faji azonosítása valamilyen módszerrel lehetséges volt.

Cluster-elemzés segítségével értékeltetem a rák-együttesekre jellemző ökológiai viszonyokat. Az összetételükben egymáshoz hasonló együttesek csoportjaira jellemző (állandó és hűséges) alakok mai rokonságának igényeiből próbáltam következtetni a bezáró kőzetek lerakódásakor uralkodó viszonyokra, figyelembe véve néhány, üledékkőzettani módszerrel nyert eredményt is. Ősföldrajzi és rétegtani következetetésre is jutottam.

Mind a gyűjtés, minden a kiértékelés munkáját egyedül, szabadidőben végeztem, csaknem harminc éven keresztül. Munkámat azonban családom tagjai, sok barátom, hazai és külföldi kollégám önzetlenül támogatta. Nevüket az angol szövegrészben sorolom fel, s ott köszönöm meg segítségüket.

# A BÁDENI EMELET ÉS A KÖZÉPSŐ PARATETHYS

A bádeni emelet regionális rétegtani egység, melynek érvényessége a Középső Paratethysre terjed ki. Radiometrikus adatok alapján nagyjából a 16,5–13,3 millió év közötti időtartamnak felel meg. A Középső Paratethys a Kárpátok ívközi medencéjében és külső előmályedéseiben levő beltenge, majd tó volt, de elérte a Balkán-hegység, Alpok és Dinári Alpok pereméig is. A Földközi-tengerrel (a kutatók egy része ezt még Tethys-nek nevezi) eleinte az ős-Alpok északi pereme mentén húzódó molassz-vályú kapcsolta össze. Ez a tengérág azonban a bádeni korszak kezdetére már végleg elzáródott. Az ősi Adriával viszont valószínűleg összeköttetésben maradt a Középső Paratethys a bádeni korszak végéig. Kelet felé a Keleti Paratethyshez kapcsolódott ez a tenger, mely a mai Balhas-tón is túl ért. A Paratethys két részének élővilága erősen eltérő volt, kimutatható faunavándorlások (KÓKAY J. szóbeli közlés) ellenére, ami a két tengerrész önállóságát, viszonylagos elszigeteltségét bizonyítja.

A bádeni korszak legnagyobb részében a tenger sótartalma az óceánokhoz közel állott, amit változatos korall-, tengeri sün- és nagyon gazdag puhatestű fauna bizonyít. A középső- és felső-bádeni alemeletek faunája azonban időleges, néhány ezrelékes sótartalom-ingadozásra utal. Az óceánokhoz hasonló sótartalom bizonyítja, hogy a világengerekkel egy vagy több tengerszoros kötötte össze a Középső Paratethyst, amelyeken át az óceáni eredetű víz beáramlott, s egyidejűleg, folyamatosan ömlött ki a felhígult vagy besűrűsödött tengervíz, a mai Gibraltári-szoroshoz hasonlóan. Ezt a kapcsolatot háromféleképpen képzelhetjük el:

1. az Atlanti-óceán felé, a Földközi-tengeren és az ős-Adrián át,
2. az Indiai-óceán felé, a Keleti Paratethysen és a Zagrosz-hegységgel párhuzamos vályún át,
3. az Indiai-óceán felé, feltételezett molassz-vályúkon át, melyek a Balkán-hegység, a Hellenidák, a Tauridák és a Zagrosz-hegység mentén húzódhattak.

A bádeni időszak nagy részében a Keleti Paratethys vize hígabb volt, mint a Középső Paratethysé. Ezért a sótartalmat stabilizáló, óceánvizet szállító kapcsolat semmiképpen sem lehetett erre. minden valószínűség szerint a Földközi-tengerrel az összeköttetés a bádeni időszak teljes tartama alatt megmaradt, s ez biztosította a víz állandó sótartalmát. A puhatestű faunában olyan nagy a rokonság a két tenger között, hogy elszigeteltségük aligha képzelhető el. Nem zárható ki viszont, hogy ezen kívül egy részben csökkentsősvizű, részben túlsós kapcsolat is megnyílt az indo–nyugat–pacifikus területek felé (RÖGL—STEININGER—C. MÜLLER 1978).

A. E. REUSS (1859) említ először egy rákot bécsei-medencei rétegekből, majd Wieliczkából (A. E. REUSS 1867). R. REUSS (1871) szintén egy fajt ír le Baden mellől. BITTNER (1877) volt az első, aki több fajt említ egyetlen lelőhelyről, Gamlitzból. Magyarországról először BROCCHI (1883) ismertet bádeni rákokat: a budapest-rákosi vasúti bevágásból hat új fajt ír le és egy hetedikhez tartozó ollót is ábrázol. BITTNER (1884, 1893) további fajokat ír le különböző lelőhelyekről.

LÖRENTHEY IMRE egyike volt a harmadidőszaki rákok legsikeresebb kutatóinak. (Német nyelvű közleményeiben keresztneve német változatát használta: Emerich.) Figyelmét elsősorban a kisvábhegyi (ma: Martinovics-hegy) páratlanul gazdag és jó megtartású felső-eocén faunára fordította, a bádeni (akkori megnevezéssel: felsőmediterrán) lelőhelyekkel kevesebbet foglalkozott. Mégis számos új fajt írt le ebből is, s újra leírta BROCCHI (1883) fajait is (LÖRENTHEY 1897, 1898a, 1898b, 1898c, 1901a, 1901b). Jó leírásokat adott a lelőhelyekről és rétegekről, modern fejlődéstanit elveket vallott s kitűnő formaérzékkel a fajon belüli változékonyságokat mindig jól felismerte. Mindezért kora legmodernebb paleontológusai között kell őt számon tartanunk. Hirtelen és korai halál szakította meg munkáját. Befejezetlen, hatalmas kézirata a háborús idők miatt sokáig kallódott, szerzője halála után sok évvel K. BEURLEN rendezte sajtó alá (LÖRENTHEY – BEURLEN 1929). Ezt a munkát sok tekintetben kitűnően végezte, de sajnos nem vizsgálta meg a kéziratban talált ábrákat. LÖRENTHEY életében megjelent cikkeinek rajzait mindig gondosan ellenőrizte, ezek a képek mind körvonalukban, mind részleteikben nagyon hűek. A posztumusz művében leírt új fajok ábráit azonban valószínűleg más rajzolta, mint a régebbieket, s azok, ellenőrzés híján sajnos megbízhatatlanok, sokszor használhatatlanok. Feltehető, hogy éppen azért késlekedett a közléssel, mert újra akarta rajzolatni ezeket. Például a *Calappa heberti* faj jobb ollójának képe (LÖRENTHEY – BEURLEN 1929, 6. tábla 11) – mely régebbi közleményből származik – kitűnő, míg a 12. képen bemutatott bal olló alig hasonlít az eredetire (vesd össze a XXXVIII\* tábla 4. és 6. képeivel). A műben (LÖRENTHEY – BEURLEN 1929) leírt új fajok szerzőiként az irodalom csaknem kivétel nélkül LÖRENTHEYT és BEURLENT idézi. A szövegből azonban egyértelműen megállapítható, hogy a leírásokat LÖRENTHEY fogalmazta, BEURLEN pedig nem változtatta meg azokat (LÖRENTHEY – BEURLEN 1929, p. 8), mivel a megváltoztatott szövegreszeket a könyvben szögletes zárójelbe tette BEURLEN, ilyen pedig a bádeni rákok leírásának szövegében nincs. Ezért a Zoológiai Nőmeneklatúra Nemzetközi Kódexének (STOLL et al. 1961) 50. cikke értelmében egyedül a magyar kutatót illeti meg a szerzőség a következő fajoknál: *Dorippe margaretha*, *Portumnus tricarinatus*, *Portunus rakesensis* (= *Liocarcinus r.*), *Zozymus mediterraneus* (= *Chlorodiella m.*), *Microcorystes latifrons* (= *Corystes l.*), *Maia biaensis* (= *Maja b.*), *Maia hungarica* (= *Micippa h.*), *Calappa praelata* és *Calianassa pseudorakesensis* (= *Callianassa ps.*).

Ezalatt Ausztriában TOULA (1904), majd GLAESSNER (1924, 1928, 1929) foglalkozott bádeni rákokkal, az utóbbit szerző a teljesség igényével. Munkájukat a második világháború után BACHMAYER folytatta (PAPP et al. 1947, BACHMAYER 1953a, 1953b, 1962, BACHMAYER – TOLLMANN 1953, BACHMAYER – KÜPPER 1952). Anyagának jelentős részét egy lelkes amatőr, K. OROSZY gyűjtötte. Az 1950-es évekig tehát a kutatások nagyrészt Magyarországra és Ausztriára korlátozódtak. A hatvanas évektől azonban a vizsgálat a Középső Paratethys egészére kiterjedt (FÖRSTER 1979a, 1979b, JANAKEVICS 1969, 1977, STANCU – ANDREESCU 1968, MÜLLER 1979b), de e sorok írója továbbra is főleg magyarországi anyagon dolgozott (MÜLLER 1974a, 1974b, 1976a, 1976b, 1979a).

A bádeni rákok kutatásának hagyománya tehát jóval több mint száz esztendős. Ennek ellenére kevés szerző foglalkozott a témaival, s a közlemények száma sem nagy.

\* Római számmal saját fényképtábláimat jelölöm.

A régebbi művek revíziója érdekében a típuspéldányokat meg kellett vizsgálnom. Ezért hazai és külföldi gyűjteményeket látogattam meg, vagy ott dolgozókat kértem másolatok küldésére. A nagy valószínűséggel elveszett anyagokon kívül csaknem minden példányt sikerült megvizsgálni. A leíró részben ezt a körülményt „material seen” (látott anyag) szavakkal jelööm.

LÖRENTHEY (1897, 1901a, LÖRENTHEY — BEURLEN 1929) anyaga rendezett állapotban a Földtani Intézet gyűjteményében van (az intézmények levélcímét az angol szövegváltozatban adom meg). Saját gyűjtését eredetileg a Pázmány Péter Tudományegyetem (ma Eötvös Loránd Tudományegyetem) Őslénytani Intézetében tartotta, ezt később, részletekben, áadták a Földtani Intézetnek. Annak ellenére, hogy az egyetemi gyűjtemények általában nem bizonyultak biztonságosnak típusanyagok tárolására, a veszteség mérsékelt, különösen, ha külföldi esetekhez hasonlítjuk. A következők vesztek el: *Microcorystes latifrons* holotypus (= *Corystites l.*, még a szerző életében elveszett, l. LÖRENTHEY — BEURLEN 1929, p. 138.), *Portumnus tricarinatus* holotypus, *Maia biaensis* holotypus (de ennek gipszmásolata megvan), és *Calianassa rakosiensis* holotypus. Az elsőként említett két fajból sajnos újabb példány sem került elő azóta. Ezzel szemben BROCCHI (1883) példányainak sorsa ismeretlen. Anyagát HEBERT és MUNIER-CHALMAS gyűjtötték, s SZABÓ (1879) még láttá a példányokat a Sorbonne gyűjteményében. A két geológus magyarországi gyűjteménynek nagy részét megtaláltam a Université Pierre et Marie Curie-n, az Institut de Macropaléontologie múzeumában (a Sorbonne egyik jogutódja), de egyetlen rák sem volt ott. A példányok valószínűleg elvesztek, más párizsi gyűjteményben sem tudnak róla.

GLAESSNER (1924, 1928) és BACHMAYER (1950, 1953a, 1953b, 1962, BACHMAYER — TOLLMANN 1953, BACHMAYER — KÜPPER 1952) anyagai valószínűleg hiánytalanul megvannak a bécsi Naturhistorisches Museum gyűjteményében, az anyag jelentős részét láttam (lásd a leíró részben). A. E. REUSS (1859, 1867) és R. REUSS (1871) típusanyagai valószínűleg elvesztek. Ezeknek legalább egy része magángyűjteményben volt. BITTNER (1877, 1884, 1893) típusainak sem sikerült nyomára bukkanni, kivéve a *Ranidina rosaliae* BITTNER 1884-ét, melynek syntypusai a Geologisches Bundesanstalt gyűjteményében vannak (STOJASPALE 1975), valamint a *Cancer illyricus* BITTNER 1884 holotypusát, melyet az Universität Graz gyűjteményében őriznek. BITTNER többi anyaga, legalábbis nagyrészt, szintén a Geologisches Bundesanstalt gyűjteményében volt, a szerző eredeti közlései szerint\*. FÖRSTER (1979a, 1979b) lengyelországi anyagát az Uniwersytet Warszawski gyűjteményében őrzik, míg JANAKEVICS (YANAKEVICH 1969, 1977) anyaga Tyiraszpolban, a Goszpedinsztitútban van. A KOJUMDZSIEVA gyűjtötte bulgáriai anyagot (MÜLLER 1979b) a Szófiai Egyetem gyűjteményében őrzik. A szerző saját anyagát a Természettudományi Múzeum Föld- és Őslénytárának felügyelete alatt, részben a múzeumban, részben lakásán tartja, mint a Művelődésügyi Minisztérium által védetté nyilvánított gyűjteményt.

Minden elérhető típuspéldányról, de legalább egy syntypusról (a leíró rendszertani részben ezt a körülményt mint „material seen” tüntettem fel) szilikongumi öntőmintát készítettem, kivéve a *Maia oroszyi* (= *Pisa o.*) és a *Necronectes schafferi* törékeny példányait. Ezért, tudományos célra, kérésre, öntvényeket küldök.

\* Lásd az „Addendumot”.

## MUNKAMÓDSZEREK, PREPARÁLÁS

A legtöbb példányt felszíni feltáráskban gyűjtöttem, melyek közül többet sokszor felkerestem. Lehetőleg nagyobb tömböket feszítettem le kibúvások falából, melyeket aztán 4—5 cm-es darabokra aprítva, binokuláris fejnagyító segítségével ( $1,5 - 3 \times$ -os nagyítással) átvizsgáltam. A lenyomatokat és törédekeket is eltettem. A szabad szemmel látható kőzettani jellegek alapján elkülöníthető minden egyes réteget külön gyűjtöttem, esetenként azonban, a rétegenkénti gyűjtéstől szigorúan elkülönítve, rézsűlábi törmelékből is vettettem anyagot, s később is tekintettel voltam annak kevert voltára. Az egységesnek tekinthető rétegek vastagsága 10 cm és 5 m között változott a gyakorlatban. A későbbi kiértékelés igazolta (4. táblázat), hogy a faunaegyüttesek típusai és a kőzettani jellegek közt szoros kapcsolat van, tehát az üledék jellege függ a leülepedés körülményeitől, melyek viszont az élővilágot szabályozó körülményekkel vannak összhangban. Ezért a praktikus szempont (a réteg helyszíni azonosításának és elkiülönítésének lehetősége) az ökológiai feldolgozás igényeit is kielégítette.

Ha a rákpáncél puha, rossz megtartású volt, sokszor nem lehetett a kemény kőzetet róla eltávolítni. Ilyen esetben a lenyomatról tisztítottam le a héj maradványait, s szilikongumival öntvényt készítettem róla. Így ugyanannak a példánynak külső és belső oldalát egyaránt lehetett tanulmányozni. A gumi odatapadásának elkerülésére hígított színtelen nitrozománcal itattam át a lenyomatokat. Ha a héj jó megtartású volt, akkor a szokásos eszközökkel, főleg tükkel, vibro-tool-lal, csípőfogóval, vésővel preparáltam. A héjakról szintén szilikongumival készíthető negatív öntőminta, amit aztán gipszzel öntöttem ki, másolat készítése céljából. Buborékok képződése ellen minden folyékony gumin, minden gipszet finom, de erős sortejű ecsettel kentem fel. A gipszöntvényt száradás után hígított zománcal szilárdítani kell, hogy műanyag öntvényekkel egyenlő szilárdságú, de optikailag kedvezőbb, átlátszatlan fehér példányokat kapjunk. Öntvény készítésére még epoxigyanta is alkalmas (a Műanyagipari Kutató Intézet által gyártott FM-4 és T-30 jelű komponensek titándioxid-liszttel keverve kis viszkozitású anyagot adtak, melyet centrifugával lehetett a finom részekbe bejuttatni), szilárdulás után a kőzetanyagot sósavval kell eltávolítani. Ez az eljárás főleg korall-lenyomatok kiöntésére volt alkalmas, de rákok esetében is jól használható. A szilikongumi-öntvényt toluolba vagy toluol-tartalmú nitrohígítóba áztatva mintegy 1,4-szeres torzulásmentes lineáris duzzadást lehet elérni. Az eljárást megismételve, 5—8-szoros méretű öntvények készíthetők.

Ha a kőzetanyag laza (mészhomok, homok) volt, a vázak pedig elég szilárdak, szitálással is gyűjtöttem. Ilyen esetben a kis példányok megtalálásának valószínűsége nagyobb, mint kalapáccsal történő gyűjtésnél, ezért a számszerű kiértékelés érdekében a példányszámokat korrigálni kellett (20. old.).

## A FONTOSABB LELŐHELYEK LEÍRÁSA

A lelőhelyeket, illetve a több rétegből álló helyeknél az egyes rétegeket kódossal jelöltem, mely egyúttal a leltári szám első tagja is (1. táblázat). Ugyanez a kód szerepel az együttesek értékelésénél is. Az egyes rétegekből gyűjtött rákfajok felsorolását az angol szövegrészben adom, néhány, kisebb jelentőségű külföldi lelőhely leírása is csak ott szerepel.

A Budapest környéki kibúvások adták a feldolgozott anyag zömét (1. ábra).

A Tétényi-fennsík (2. ábra) egri és neogén kőzetekből álló magaslatán a bádeni rétegek, kis vastagságuk miatt, csak kis foltokban és sávokban bukkannak elő. A Kamaraerdei út bevágása (MK, 2. ábra) ma is hozzáférhető. Korábban Katona út (Militärstrasse) néven szerepelt (LÖRENTHEY 1911). Ma csak két réteg látható (3. ábra), az MK jelű 1 m vastag rosszul osztályozott kavicsból álló konglomerátum, melybe korall-lenyomatok (*Tarrellastraea* sp., *Porites* sp., *Siderastraea* sp. és egy *Tarrellastraea*-habitusú, de valószínűleg más nemzettséghöz tartozó zátonyépítő korall) települnek lencsésen, kimosott helyzetben. A feltáras északi végén azonban *Porites* sp. élő helyzetben betemetett oszlopai láthatók.

A rákok leírása kapcsán KÓKAY (in MÜLLER 1976a) felsorolta és kiértékelte a puhatestű faunát. Szerinte a víz 30 méternél lényegesen sekélyebb, jól mozgatott, oxigéndús volt, óceánihoz közelálló sótartalommal. A korallos réteget 1,5 m vastag mészhomokkő fedи (MKF), gyengén diszkordánsan. Kőzettanilag mikroonkoidos váztörmelékes „grainstone”. A Balatoni út és a Szabadkai út csatlakozásánál levő bevágások (MF, 2., 3. ábra) ma is hozzáférhetők, alapszelvényé kiépítve. A rétegsor egy részét, az akkor meglevő, sekélyebb feltárasok alapján SCHAFARZIK—VENDL (1929) is ismerteti. A szürke, nagy *Pectinidae* fajokat tartalmazó eggenburgi korú homokot diszkordánsan fedik itt a bádeni rétegek: rosszul osztályozott durva kavics, 20 cm-t elérő nagyságú görgetegekkel, melyek egy részét kagylók fúrták meg. Erre 0,8 m vastag homokos—kavicsos mészkő (MFK) települ, sok *Glycimeris*-lenyomattal. Puhatestűit KÓKAY vizsgálta (in MÜLLER 1976a). A következő 1,6 m vastag szint puhább és keményebb rétegekből álló, az előbbinél kevésbé kavicsos és homokos mészkő. Ezekből külön nem tudtam gyűjteni rákokat, de a részű lábánál talált tömbököt vettet mintákat. Ez a kevert anyag kapta az MFA jelet. A bevágás legfelső rétegét más feltárasban vizsgáltam (MM, MB). A feltárástól nyugatra egy kisebb gödörben kváromokkó fedte az előbb említett rétegeket (MA). KÓKAY idézett leírása, mely az MFK rétegre vonatkozott, az MA és MFA együttesekre is érvényesnek látszik: szerinte az óceánihoz közelálló sótartalmú, erősen mozgatott vizű környezetben rakódott le az üledék, 30 méternél sekélyebb tengerben. A rétegen a kavicsszemek nagysága dél felé csökken, ami valószínűleg azt jelenti, hogy észak felé kavicsos anyagból álló part lehetett (valószínűleg kárpáti korú rétegekből eredő kaviccsal), s a víz enyhén mélyült dél felé. A nem túlságosan nagy hullámenergia a kavicsokat csak néhányszor tíz méter távolságra tudta elhordani, így a délebbi részeken már csak homokkövet találunk.

A kvárcszemcséket tartalmazó rétegek fedője tiszta mészhomokkő, melyben sztromatolitok és onkoidok vannak (MB és MM). A kis gödröket, melyekben ezek fel voltak tárva, már nagyrészt betöltötték, de a felszínen elszórt anyagból még lehet gyűjteni, a sztromatolitok pedig nagy területen a felszínen láthatók. A képződmények kriptalga-Foraminifera eredetűek (LELKES—MÜLLER in press). A beágyazó kőzetanyag mikroonkoidos „grainstone”, sok puhatestű-lenyomattal, főleg *Cerithium*-mal. A réteg vastagsága NyDNy felé csökken, s ugyanarra a fennőtt sztromatolitok helyét fokozatosan gömbszerű, görgetett onkoidok váltják fel, melyek száma, ha továbbhaladunk, egyre csökken. Mindebből partközeli, sekélyvízű tengerrézsre gondolhatunk, ahol a fenék dél felé mélyült, hasonlóan ahhoz a helyzethez, amire a fekü homokos—kavicsos kőzetek alapján következtettünk. KÓKAY (in MÜLLER 1974a, p. 119) a puhatestűek alapján sekély szublitorális tengerre következtetett, 26–28%-os sótartalommal. A *Cerithium*-ok gyakorisága sűrű alga vagy tengerifű borítást jelez. A tizlábú rákok fajainak száma is növekszik dél felé. A szelvény déli végén a fent említett réteget részben fedи, részben pedig helyettesíti egy finomszemű, rosszul osztályozott szemesékből képződött mészhomokkő, mely sosem tartalmaz makroonkoidokat (MDG). A rákokon kívül ott talált faunaelemeket KÓKAY hatá-

rozta meg (in MÜLLER 1974b, p. 275). Szerinte a víz 30–40 méternél sekélyebb volt, a sótartalom közel állott az óceánihoz, de időnként változott.

**Diósdon** (2. ábra), a Tétényi-fennsík déli részén kis területen bukkannak ki a bádeni kőzetek az Országos Érc- és Ásványbányák elhagyott kőfejtőjében („krétabánya”) és a Tégla- és Cserépipari Tröszt működő homokbányának felső peremén. A pannóniai homok itt jelentős ugrómagasságú vető mentén érintkezik a bádeni, és az arra erős diszkordanciával települő szarmata mészkövekkel. Az MDZ jelű lelőhely a vető mentén van, a homokbánya peremén. A kemény, tömör mészkő vastagsága két méternél több. *Tarbellastraea* sp., *Porites* sp. s két másik zátonyépítő korallfaj lenyomatának tömegét tartalmazza, mind kimosott helyzetben. A beágyazó anyag zöme rosszul osztályozott, s egyes lencsékben tartalmaz rákmaradványokat, köztük igen nagy példányokat is. A kőfejtő egy másik részén néhány tömb másfajta faunát tartalmazott (MND), mely valószínűleg fiatalabb a korallas rétegnél, de kapcsolatuk fedett.

**Bián** (1. ábra), a Tétényi-fennsík nyugati nyúlványainál bádeni és szarmata rétegek tűnnek ki a „Nyakaskő” nevű szikla körül. Ezt STRAUSZ (1924) és SCHAFARZIK—VENDL (1929) írták le a legrészletesebben, a korábbi irodalom megemlítéseihez. A legalsó feltárt réteg a szikla alatti árokban bukkan ki kis felületen: szürke aleurit, aleuritos homok (MNA). E fölött négy méter vastag homokos mészkő következik. Ez gazdag puhatestű faunát tartalmaz lenyomatok formájában (MN).

Budapest keleti részén, a klasszikus rákosi vasúti bevágás környékén van az a lelőhely csoport, mely a Paratethys leggazdagabb és legszebb megtartott rákanyagát adta (1., 4., 5. ábra). A rétegsorokat számos cikk tárgyalja, legújabban, összefoglaló igénnnyel KÓKAY J.—MIHÁLY S.—MÜLLER P. (in press) írták le. Kóbánya-felső és Rákos vasútállomások között van a delta alakú bevágás, melyet a múlt században mélyítettek, s azóta is többször bővíttettek. A ma is hozzáérhető alapszelvény bádeni és szarmata rétegeket tár fel. SZABÓ (1879), FRANZENAU (1881), VADÁSZ (1906), majd a legrészletesebben SCHAFARZIK—VENDL (1929) írták le e szelvénnyt. BROCCHI (1883) faunáját is itt gyűjtötte HEBERT és MUNIER-CHALMAS. Az MRC jelű réteg alatti szintek (5. ábra) tufaeredetű ásványokat is bőven tartalmaznak, melyek részben a fekü kárpáti (?alsó-bádeni) „Tari Dáicitufa Formáció” anyagának áthalmozásából, részben új kitörésekkel származhatnak. A felsőbb, MRF jelű réteg biotit-jának kora ugyanis lényegesen fiatalabb ( $13,4 \pm 0,6$  millió év a szomszédos, Keresztúri úti feltárást azonos, MEG jelű rétegeből vett minta alapján), mint az MRZ jelű réteggel egyidős *Cepaea*-s tufitét ( $15,6 \pm 0,8$  millió év, BALOGH—ÁRVÁNÉ-SÓS—PÉCSKAY 1980). Valószínű, hogy a mélyebb szint biotitja csaknem kizárolag a régebbi tufák áthalmozásából ered, a felső rétegé pedig zömmel friss kitörés terméke. A bevágás keleti részén a fekü riolit- vagy dácittufát diszkordánsan *Porites*-es korallzátony fedeti, 1,5–2 méter vastag lencse alakjában (MRZ). *Porites*-lenyomatokon kívül mészalgák (?*Neogoniolithon*), kevés *Bryozoa* és mindenkorral két töredék *Tarbellastraea* került elő. Az 5–10 cm vastag, 0,5–1 méter magas korallágak eredeti helyzetükben ágyazódtak be a tufitos mészkőbe. A *Porites*-t eredetileg *Millepora*-nak határozták meg (LÖRENTHEY—BEURLEN 1929, p. 35), de *Hydrozoa*-maradvány itt egyáltalán nem került elő. A korallzátonyt részben helyettesíti, részben fedi az MR45 és MR8 jelű homokkő, mely középszemű, erősen bioturbált, rosszul rétegzett. Ebben a rákok kitűnő megtartásúak. A homokkövet homokos mészkő fedi, melynek alsó részében a *Linga columbella* (LAMARCK) a leggyakoribb puhatestű, míg felső része osztrigapadokat tartalmaz (MRL, ill. MRO). A kvarc- és egyéb terrigén szemcsék mennyisége felfelé csökken. A vasúti bevágás legvastagabb rétege a mészkőre települő finomszemű, rosszul osztályozott tufitos homokkő (MRF). Ugyanez a réteg a Keresztúri úton is látható (MEG). Jellemző a tömeges *Callianassa*-ásásnyom (*Ophiomorpha*). Gyakoriak a nagy *Pectinidae* fajok és a szabálytalan sünök.

A homokkövet tiszta mészkő fedi (MRC). Gyakoriak benne a *Cardium* és *Cerithium* fajok. Közöttani lag mikroonkoidos váztörmelékes „grainstone”. Erre kereszt- (ferde) rétegzett mészhomokkő települ, melyben egy 5–15 cm vastag réteg tömegesen tartalmaz makroonkoidokat, melyek elérik a 20 cm-es méretet is (MRS), hasonlóan a Tétényi-fennsík megfelelő rétegéhez (MB). A beágyazó kőzet mikroonkoidos váztörmelékes „grainstone”, kevés ooidal (LELKES—MÜLLER in press). Az e fölött települő rétegekből még nem került elő rák-maradvány.

Mintegy egy kilométernyire Rákostól az Örs vezér tere környékén szintén gazdag lelőhelycsoport volt, de az építkezések befejeztével ezeket mind betemették. Az utóbbi húsz évben sok árkot és alapgödröt ástak itt, de már a húszas években is került elő anyag. A rétegsorokat és faunákat STRAUSZ (1927), BARTKÓ—KÓKAY (1966), MIHÁLY (1969), MÜLLER (1976b, 1979a) és KÓKAY—MIHÁLY—MÜLLER (in press) ismertették. Az alsó homokos rétegek nagyjából egyformák voltak valamennyi feltárástban, míg a felső tiszta mészkövek fáciése helyről helyre változott. A legalsó rák-tartalmú réteg vékony korallas tufás mészkő lehetett, de ezt csak a Kerepesi út mentén mélyített csatorna-árokban kidobott anyagban sikerült felismerni. Szegényes rákanyaga az MRZ lelőhelyéhez hasonló. E fölött mintegy 2 m vastag homokos mészkő (MKC) települt, ez a Kerepesi út mentén volt feltárvá. Ez a rákosi MRL és MRO rétegeknek felelhet meg, de azoknál mozgatottabb, sekélyebb vízből ülepedhetett le. Jellemző puhatestű a *Glycymeris obtusatus* (BARTKÓ—KÓKAY 1966). A mészkő fedője 4–5 m vastag laza, középszemű homok, melyet a Gyakorló úton (MGY) és a Sugár üzletház pince-

tömbjében (MOH) tártak fel (4., 5. ábra). Megjelenésében ez a kváromoknak tűnik, de kvarctartalma alig 5%, a többi víztiszta szemcse kőzetüvegnek bizonyult (RAVASZNÉ BARANYAI L. szóbeli közlése). CaCO<sub>3</sub>-tartalma 40% körüli, ezenkívül földpátok, csillámok és agyagásványok mutatkoztak a röntgen-diffraktogramon, tehát sok benne a vulkáni eredetű anyag, de erőteljesen átmosva. A finomabb szemcsék az egykorú MRF réteg anyagában keresendők. Az erőteljes átmosás és az abból következő viszonylag jó osztályozottság nem a hullámenergiának, hanem a bioturbációt a következménye. Nagy *Pectinidae* fajokat, tengeri sünöket és Foraminiferákat tartalmaz. A rákok aránylag jó megtartásúak voltak, ezért itt szitálással lehetett gyűjteni.

A homokot mészkarbaszorozat fedi, két különböző helyen más-más kifejlődésben. A Gyakorló útnál (5. ábra) a homokra 0,4 m vastag, rákokat nem tartalmazó *Ervilia*-s mészkarbász települ, ezen pedig a felterábsan 0,5–0,8 m vastagságban látható, *Cerithium*-lenyomatokat tömegesen tartalmazó mészkarbász (MGF) fekszik, melyben makroonkoidok vannak, mészhomokkő—kavics vagy „black pebble” maggal (LELKES—MÜLLER in press). A Sugár üzletházánál a homokot vékony tufitréteg fedi. Ezen 2,2 m vastag, részben laza, széteső, részben cementált mészhomokkő (MOE) fekszik (5. ábra), mely a Gyakorló úti MGF mészkarbász gyakorlatilag egykorú. Ez váztormelékes „grainstone”, kevés mikroonkoiddal. Jellemzőek a kis osztrigák, *Chlamys* fajok, *Cardium*, szabályos sünök, *Lingula* sp. A rákok megtartása tökéletes, s ahol a kőzetanyag laza volt, ott szitálással lehetett gyűjteni. A Paratethys leggazdagabb ismert rák anyaga innét származik.

A Mező Imre úton mélyült a Metró H. 34 sz. fúrása. Negyedíoszaki homok és kavics alatt finomszemű aleuritos homokkő és mészhomokkő változók (KÓKAY, szóbeli közlés), mely alatt tarka, középső-bádeni agyag és aleurit települ. A felső-bádeni rétegekben 25,8 és 33,0 m között (MH34) találtunk rákokat. A Telepy utcában (MTE, 1. ábra) a század elején csatornaárokban egyetlen rák faj került elő. A feltárást SCHAFARZIK (1903) írta le.

Visegrádon (6. ábra), a Fekete-hegy oldalában rosszul feltárt bádeni rétegsor található (SCHAFARZIK—VENDL 1929, SCHOLZ 1970, BÁLDI—KÓKAY 1970). Valószínűleg ez volt REUSS (1872) „Nagy-maros” megjelölésű koralljainak típuslelőhelye is. Andezittömbös—kavicsos tufitokat alsó-bádeni márgás, tufitos mészkarbászok takarnak. Ezek alsóbb része (MV) korallos, gyakoribb alakok: *Tarbellastraea* sp., *Favia magnifica* REUSS, *Stylophora subreticulata* REUSS, *Porites* sp. A mészkarbászok felsőbb része (MVA) vörös-mészalgás. A feltárasok hiánya miatt a tényleges rétegsort nem lehetett felvenni. A rákok megtartása mindenkor rétegben gyenge, de az MV mészkarbász faunája fajokban gazdag. Törökmezőn (6. ábra) mindenkor nem ismertetett új lelőhelycsoport van a turistaház közelében. Az új út bevágása (MTH) *Heterostegina giganteoformis* PAPP óriás Foraminiferát tartalmazó tufás mészkarbászban van. Innen 500 méterre északra már már nagyrészt betöltött kis kőfejtő (MTZ) korall-lenyomatokban gazdag mészkarbász tárt fel. Mintegy 20 korallfajt lehetett elkülöníteni, gyakori a *Stylophora subreticulata* REUSS, *Tarbellastraea* sp., *Montastraea* sp. és egy vékony, lemezszerű alak, mely Nagyhöflányban (MNH) is előfordul. A rákok megtartása aránylag gyenge. Tovább északra egy anyagnyerő-gödör (a helybeliek megnevezése szerint „murvabánha”) (MTM), mintegy 3 m vastagságban tufitos Mollusca-s márgát tár fel. Ennek nannoflórája alsó-bádeni jellegű (NAGYMAROSI, szóbeli közlés). Zebegényben, a bakókuti (bőszobi) kőfejtő (MZZ, 6. ábra) lágy, vörös-mészalgás agyagos mészkarbász tár fel, melybe 4 m vastag, kemény, tiszta mészkarbász települ. Ez a mikrites mészhomokkő tele van egy kicsiny, elágazó ahermatipikus korall lenyomataival, mely egyes mai *Oculina* fajokhoz hasonlít. Kevés *Porites* sp. társul hozzá. A rákok ebben a korallos mészkarbászban vannak, kisebb lencsékben. A szobi (MSZ, 6. ábra) lelőhely a puhatestű maradványokról ismert. Valószínűleg középső-bádeni (KÓKAY, szóbeli közlés). A Damásdi-patak bevágásában alul szürke, feljebb sárga laza homokkő települ, kb. 6 m vastagságban. Erre 0,1 m vastag lumasella, majd 0,5 m vastag sötétbarna faunás tufit települ. Mindegyik rétegben van rák, de a puhatestűkhez képest nagyon kevés. Eddig csak ollókat találtam.

A Mátra északnyugati vége és a Cserhát csatlakozó része (7. ábra) szintén szép rák anyagot adott. A területet NOSZKY (1940, pp. 102–113.) és HÁMOR (in press) írta le. Mátraverebély—Szentkúton a Szent László-forrásnál és a Meszes-tető oldalában a Noszky-féle „bázisrétegek”, homokkövek, homokos és márgás mészkarbászok tartalmaznak rákokat. Sámonházán a Természettudományi Múzeum gyűjteményében levő anyag cékulái szerint két lelőhely van, ezek közül az egyik fekvése ismeretlen: Halastó-hegy, bázishomok. A másik lelőhely: Csúcs-hegy alja, kis kőfejtő a Kis-Zagyva völgyében, s talán azonos HOJNOS (1923, 1933) feltárással.

Nyírádon (MNN) a falu temetőjétől 500 méterre nyugatra (8. ábra) a Sümeg felé vezető országúton kis bevágás van. Egyetlen réteget, durvaszemű mészhomokkötőt tár fel mintegy 2,5 m vastagságban. Ebben nagy puhatestűek és néhány koralltelep lenyomata található. Ezek még nincsenek meghatározva. Balatonakaliban, a BAK-40. sz. fúrás (BAK) szarmata és bádeni rétegeket tár fel (KÓKAY, szóbeli közlés). Ebben a 35–38 m közötti szakaszban voltak rákok. A Mecsek hegységben, Kishajmás-Szatina és Husztót állomások között az új pályának mély bevágást ástak a 70-es években, ?alsó-bádeni homokköben. Itt SOLT P. gyűjtött rákokat.

Fertőrákoson az FRK-21. sz. fúrás (9. ábra) vastag ?középső-bádeni mészkarbász tár fel, mely-

ben alig volt gerinctelen makrofauna, zömmel vörös-mészalgás mészkőből állott a rétegsor, melyben 268 m-nél mintegy 20 cm vastag zátonymaradvány települt (olisztitolit?), rákokkal.

A hazai lelőhelyeken kívül azokat a kúlföldi feltárásokat említem meg, ahol gyűjtöttem. Nagyhöflányban (9. ábra) (Gross-Höflein, Burgenland) a „Fenk” cég nagy kőfejtőjét (jelenlegi tulajdonos BUCHINGER) a bádeni lajtamészkövek faciosztratotípusának jelölték ki (PAPP et al. 1978, p. 194). A rétegsort DULLO (1983) tanulmányozta részletesen, üledékközöttani, faciológiai szempontból. Ő négy korallos szintet talált benne, ezek közül alulról a második („oberer Korallenkalk”, PAPP et al. 1978, p. 197) gazdag és jó megtartású rák faunát adott. A réteg koralljai: *Tarbellastraea* sp., *Porites* sp. (ez a kettő kőzetalkotó mennyiségben), *Stylophora* cf. *subreticulata* REUSS, *Acanthastraea* sp. és egy lemezes forma, ami Törökmezőn (MTZ) is gyakori. BACHMAYER és TOLLMANN (1953) rák anyaga valószínűleg szintén kizárolag ebből a rétegből származik, de az ő gyűjteményükből csak egy példányt tudtam megvizsgálni. Ennek ellenére valószínű, hogy az általuk említett rákok többségét megtaláltam. A fejtőt töltik, de a korallos réteget valószínűleg sikerül megvédeni a betakarástól (STEININGER, szóbeli közlés). Az Oszlopon (Oslip, Ausztria, Burgenland) levő Bauer-féle homokbánya részletes analízisét is DULLO (1983) adja (MOS, 9. ábra). Kövületmentes homok felett mintegy méter vastag *Ophiomorpha*-nyomos (*Callianassa* ásásnyom) homok sok rakollót tartalmaz. Ezt vörös-mészalgás mészkő fedi. A fejtőt ma is művelik. Alsó-Ausztriában, Deutsch-Altenburg mellett, a Hundsheimer-hegyen nagy elhagyott, de ma is hozzáférhető kőfejtő van (DAL, 9. ábra). Mezozoós mészkövön, bázis breccsa – konglomerátum fölött mintegy 30 m vastag mészhomokkő van, melybe vékony mohaállatos márga települ (BACHMAYER 1953a). A Grobie-i [déli Szentkereszthegység, Lengyelország (MGR)] lelőhelyet nyomtatott anyagban még nem közölték, csupán FÖRSTER (1979a) említi, a R. WRONA által gyűjtött rákok kapcsán. Nincs feltárás, a szántóföldön heverő kövek zöme vörös-mészalgás mészkő, melyek között néhány korallos darab akadt, ebben voltak a rákok.

A tízlabú rákok — különösen a rövidfarkúak — törzsfejlődése, speciációja gyors folyamat. Az összes ismert, jól leírt pleisztocén faj azonosnak bizonyult ugyan a maiakkal, de a pliocén anyagban már sok kihalt fajt találtak (GLAESSNER 1929). Az 5–6 millió éves messinai anyag zöme is ma élő alakokból áll (a szerző publikálatlan anyaga), de ezek egy része már alaktani különbségeket mutat a maiakkal összehasonlítva. A bádeni és a velük rokon mai fajok között ezek a különbségek már határozottabbak, például a *Pachygrapsus hungaricus*—*P. marmoratus*; a *Brachynotus februarius*—*B. sexdentatus* és *B. foresti*; vagy az *Ebalia corsi*—*E. edwardsii* esetében. A mai nemzetiségek kialakulása is gyors volt. A legtöbb valószínűleg a miocénben és a pliocénben keletkezett (MÜLLER 1979b). Így a 12–15 millió év előtti, bádeni fajok meghatározásánál óvatosan kell eljárni, ha azokat mai alakokkal hasonlítjuk össze, feltéve, hogy nincs a közben eltelt időből is elegendő adatunk a faj változásának vagy állandóságának bizonyítására. Ilyen folyamatos adatsor azonban csak a *Dardanus arrosor* (HERBST) faj esetében ismert. Ezért minden más bádeni fajnak új nevet adtam, még ha létezett is alaktanilag hasonló mai forma. Ez az új név bizonyos esetekben csak időleges megoldásnak tekintetű.

Hasonló eljárást nem követhettem a nevek esetén. Ezért a mai neveket eredeti formájukban használtam, ha egy bádeni faj a nemzetég mai fajai által megszabott keretbe illett morfológiai szempontból.

Gyűjtőnemzetiségek megjelölésére a nevet két szimpla idézőjel ('...') közé tettem. Olyankor használtam ezt a megjelölést, ha a fosszilisan megmaradó testrészek nem teszik lehetővé a biztos generikus meghatározást, de a közeli rokonság biztosra vehető. Közeli hasonlóság hiján új genus-t kellett bevezetni. Idézőjelet ("...") olyan fajoknál használtam, melyeket régi szerzők nyilvánvalóan tévesen osztottak be jól ismert, mai nemekbe (pl. „*Cancer*” *carniolicus* BITTNER), de a rossz megtartás vagy a típuspéldányok elveszett volta miatt nem lehetett jobb megoldást ajánlani.

Az ollók ujjai erősen meszesek, sokszor tömörek, ezért ezek a rákok leggyakrabban fosszilizálódó részei. Mivel ezek alakját elsősorban a funkció alakítja ki, mely azonos lehet különböző rendszertani egységekben, gyakori a konvergencia. Rendszertani vizsgálatuk mérsékelt sikерrel járt (BACHMAYER 1962, JANSEN 1972, FÖRSTER 1979a, 1979b). A sikertelenség okait FÖRSTER (1979b) részletesen elemzte. Hozzá kell tenni, hogy a zoológusok és paleontológusok ritkán adnak leírást vagy használható ábrát az ujjakról, ezért a mai fajokkal való összehasonlítás általában csak múzeumi példányok birtokában lehetséges. Egyéb testrészeket is tartalmazó rétegekből származó ujjakat csak néhány bádeni fajnál ismertettek (GLAESSNER 1928, LÖRENTHEY—BEURLEN 1929, BACHMAYER 1953a).

E munkám egyik célja, hogy az elkülönült testrészek faji azonosságát nyomozzam, beleértve az ujjakat is. E célból mai rokon formákkal hasonlítottam össze azokat (hazai gyűjteményekben, a párizsi Muséum National d'Histoire Naturelle-ben, de legtöbbször külföldről kölcsönbe kapott példányok segítségével). Egy adott rétegen a testrészek relatív gyakorisága támpontot adhat az összetartozásra, feltéve, hogy a betemetődés előtti szelekció nem volt erős. Egyes esetekben az ujjak még az olló többi részével együtt találhatók. Az ollótő (kéztő) díszítése sok csoportban hasonló a carapaxéhoz, s így azonos fajba tartozásuk eldönthető. Ha az olló-ujjak meghatározásához sikeresen valamivel közelebb jutni, ennek az lenne a jelentősége, hogy ezek sokszor találhatók olyan rétegekben, ahol rajtuk kívül csak puhatestűek és Foraminiferák gyakoriak, s így új lehetőséget biztosíthatnak rétegtani, ősföldrajzi és paleoökológiai vizsgálatokhoz. A rákok pedig ilyen kutatások céljára nagyon alkalmasak, mert fejlődésük gyors, terjedési lehetőségeik minden felnőtt állapotban, minden pelágikus lárvaként jelentős, s így óceánokon (GARTH 1974) vagy a rendestől eltérő sótartalmú víziutakon is (POR 1978) keresztül juthatnak. A paleoökológiai vizsgálatok szempontjából pedig előnyös, hogy a mai alakokkal sok, ökológiai adatot is tartalmazó monográfia foglalkozik.

A rendszertani részben (49. old.) a fajokat a GLAESSNER (1969) által használt beosztásban mutatom be, mert a legtöbb őslénytani munka ezt használta az utóbbi időben. Ennek ellenére meg vagyok

győződve arról, hogy GUINOT (1977) és DE SAINT LAURENT (1980a, 1980b) utat mutattak a tízlábúak rendszerének jobb megértéséhez.

A fejezetben használt rövidítések:

MNÖ: Természettudományi Múzeum Őslénytára  
FI: Földtani Intézet  
CA: a szerző anyaga  
NHMW: Naturhistorisches Museum, Wien

A rákok testrészeire vonatkozó rövidítések:

ca: carapax  
ch: olló  
da: dactylus  
ff: rögzített ujj

A nemzetiségek típusfajait csak akkor adom meg, ha nem szerepelnek GLAESSNER (1969) munkájában. A szinonima listákban csak azokat a munkákat idézem, melyek a taxonra vonatkozó új adatot vagy emendációt közölnek. Ezért, alapvető jelentőségük ellenére, a Fossilium Catalogus (GLAESSNER 1929) és a Treatise on Invertebrate Paleontology (GLAESSNER 1969) idevágó köteteket is csak egyes esetekben említem.

# A BÁDENI TÍZLÁBÚ RÁK FAUNA JELLEGZETESSÉGEI

A bádeni rák fauna modern jellegű. Bár a 102 meghatározható alakból csak egyet tekintettünk mai fajhoz tartozónak, a 72 nemzetsegből csak 12 bizonyult nagy valószínűsséggel kihaltnak, tehát a nemek 85%-ának ma is van élő képviselője. Ezek a számok és arányok óhatatlanul tartalmaznak szubjektív elemeket is, de közelítően érvényesek. Bizonyos archaikus jellegeit is tükröz az a tény, hogy régi eredetű családoknak elég jelentős a szerepe (*Calappidae*, *Dromiidae*) mind a fajok számát, mind a példányszámot tekintve.

A *Callianassidae* család meglepően sok bádeni fajt számlál, ez részben az ollók erősen elmeszesedett vázának, a beásó életmódnak és ebből következőleg a fosszilizálódásra való alkalmasságnak tulajdonítható. A bádeni '*Callianassa*' fajok zöménél (valószínűleg a '*C. brocchii*' kivételével) a jobb és bal olló egyforma volt (homochelia), ellentétben a mai fajokkal. Ez a kérdés további vizsgálatot igényel. A *Galatheidae* és *Porcellanidae* fajok nagyon közel állnak mai rokonaihoz, ami e régi eredetű családoknál érthető. A *Paguridae* maradványoknál, ahol a közvetlen összehasonlítás lehetséges volt (*Pagurus*, *Petrochirus*, *Diogenes*, *Dardanus arrosor*), a fosszilis ollók nagyon hasonlóak a maiakhoz.

Feltűnően nagy a bádeni *Dromioidea* fajok száma (hat), pedig általános nézet volt, hogy neogén fajaik ritkák. Ha a korallzátonyon élőket nem számítjuk, két faj marad, míg a Földközi-tenger sekély vizében ma csak egy él. Az alakok egy része hasonlít mai rokonához, de a *Kerepesia* és a *Dromiidarum?* n. gen. kihalt nemekhez tartozhatnak. Esetleg egy további faj (?*Dorippe carpatica*) tartozik még a *Dromiacea* csoportba. Bár a *Calappidae* fajok száma nem nagy, feltűnően gyakoriak a legtöbb rétegen, ami részben szilárd vázunknak köszönhető, de nagyrészt egykor tényleges gyakoriságukat tükrözi. Az alakok nem különböznek lényegesen mai rokonaitól, de a *Mursia lienharti* és a *Matuta brocchii* fajok, nemzetsegük keretén belül primitívnek tekinthetők rövid oldaltüskéjükkel, bár ilyen fajok ma is élnek mindenktől genus-on belül. Érdekes (másodlagos?) jelenség a cirpelőszerv hiánya *Mursia lienharti* fajnál. Nyolc *Leucosidae* fajt találtunk a bádeni faunában, ez a viszonylag magas szám nagyon szilárd carapaxukkal függhet össze, mely a megmaradás valószínűségét növeli. Nagy részük az *Ebalia* nembe tartozik, s ebben a fauna a mediterrán faunára hasonlít, ahol szintén nagy az e nembe tartozó fajok száma. Két vagy három faj kihalt nemzetseghez tartozik.

A bádeni fauna egyik legfeltűnőbb sajátossága a *Majidae* fajok kis száma. Bármelyik mai provinciában az uralkodó családok közé tartozik a tengeri pókoké. Az öt fajból egy zátonylakó. Különösen feltűnő a tengerifű- és algalakó kis alakok hiánya, melyek más kis formákkal együtt, a mai Földközi-tengeri és indo-nyugat-pacifikus faunák fontos elemei. A két *Parthenopidae* faj nagyon közel áll a két leggyakoribb földközi-tengeri fajhoz, valószínűleg leszármazási kapcsolatban áll azokkal. A bádeni *Cancroidea* csoport néhány érdekes jellegzetességet mutat. A „*Cancer*” gyűjtőgenus-ba sorolt alakok valószínűleg kihalt csoportot alkotnak. A csoport fajainak száma nagy, különösen, ha a mai Földközi-tengerhez hasonlítjuk, még akkor is, ha ez a szám néhány téves meghatározást is tükröz. A *Portunidae* fajok száma is nagy: legalább 17. A fajok egy része ma is élő nemzetseghez tartozik, de a nemek jelentős része biztosan kihalt. A *Portunus*, *Liocarcinus*, *Thalamita* és *Charybdis* fajok hasonlóak ma élő rokonaihoz, a Földközi-tengerben vagy az Indiai-óceánban. A mai tengerekhez hasonlóan a bádeni *Xanthidae* fajok száma is nagy, különösen a zátonylakóké.

A bádeni fauna csak kis része a világ neogén tízlabú rákjainak. Ezért a csoport törzsfejlődésére csak kevés adatot várhatunk tőle.

Megállapítható, hogy a *Majidae* csoport változása gyors volt a területen a miocéntől máig. Ez a változás (a fajok számának gyors növekedése) egyaránt jelenthet helybeli gyors fejlődést, vagy tömeges bevándorlást. Ennek eldöntéséhez jobban kellene ismerni a késő neogén és pleisztocén európai faunákat. A *Dromiacea* csoport erős csökkenést mutat a bádeni korszaktól máig, de ez a folyamat nagyrészt a korallzátonyok itteni eltűnésekének következménye. A *Cancroidea* csoport visszahúzódása, úgy tűnik, tényleges fejlődési folyamat, bár Amerika körüli vizekben szerepük ma is jelentős. A *Xanthidae* csoport regressziója, a *Dromiacea* csoportba hasonlóan, a korallzátonyok eltűnéssel kapcsolatos. Ha a mai indo-nyugat-pacifikus területhez hasonlítjuk, inkább növekedést tapasztalunk. A csoport gyors fejlődését látszik bizonyítani az is, hogy a bádeni emeletből hiányzik néhány, ma nagyon elterjedt, gyakori nemzetseg. A többi család olyan kevés fajjal szerepel, hogy fejlődésükre vonatkozólag semmi adatot nem ad a bádeni fauna.

A bádeni emeleten belül három alemeletet különböztetnek meg: az alsó (morva), középső (wieliczkai) és felső (koszvói) egységet. Ezek megkülönböztetése nem egyszerű, ha egyes feltárások, rétegsorok korát vizsgáljuk. Sztratotípusokkal és határsztratotípusokkal határozták meg ezeket (PAPP et al. 1978), főleg Foraminiferára és a nannoflórára alapozva a megkülönböztetést. Ezek a csoportok a rákok zömét adó mészkövekből hiányoznak, vagy jellemző képviselőik nem találhatók meg. Az ezzel kapcsolatos nehézségek ellenére a rág-tartalmú sekélyvízi rétegek korát korrelálhatjuk a Foraminifera-dús mélyebbvízi képződményekéhez. Ehhez geológiai, ősföldrajzi módszereket használhatunk, vagy olyan puhatestűeket vehetünk figyelembe, melyek mindenkor kőzettípusban megtalálhatók. Az ősföldrajzi módszer alkalmazását megkönyíti az a körfülmény, hogy a korai és késő-bádeni kor-szakok között jelentős tektonikai események történtek, gyökeresen átrajzolva az ősföldrajzi térképet a Középső Paratethys nagy részén.

Budapest környékén két olyan terület van, amelyik csak alsó- vagy csak felső-bádeni üledékeket tartalmaz, legalábbis a rákokban gazdag sekélyvízi eredetű kőzetek esetében.

Budapest belterületén és közvetlen környékén (Rákos, Tétényi-fennsík) a mészkövek kora kizárával felső-bádeni. Ezt a puhatestűek bizonyítják (BARTKÓ—KÓKAY 1966), és az a tény, hogy rájuk konkordánsan települnek a szarmata rétegek, legalábbis a terület jó részén. A vékony rétegsor faunája annyira egységes, hogy két alemeletbe sorolása szóba sem jöhét. Az alsóbb kőzetrétegek radiometrikus kora is még a felső-bádeni alemeletre utal ( $13,4 \pm 0,6$  millió év, BALOGH—ÁRVÁNÉ-SÓS—PÉCSKAY 1980).

A Pilis északi részén és a Börzsönyben azonban alsó-bádeni üledékeket találunk, csak Szob esetén vetődött föl a középső-bádeni kor lehetősége. A kort mikro-, makro- és nannofosszília is bizonyítja (BÁLDINÉ BEKE et al. 1980, NAGYMAROSI 1980 és szóbeli közlés).

A két terület tízlábú rágfaunája sok tekintetben eltér egymástól, bár hasonló ökológiai körfülmények nyomozhatók mindenkor helyen. A leglényegesebb eltérés az, hogy egyes gyakori fajok csak az egyik vagy csak a másik részen találhatók.

Az alsó-bádeni terület legjellemzőbb ilyen fajai:

*Dardanus arrosor*  
*Kromtis koberi*  
*Dynomene emiliae*  
*Calappa praelata*  
*Carpilius antiquus*

Csak felső-bádeni lelőhelyeken kerültek elő:

*Petrochirus priscus*  
*Dardanus substriatiformis*  
*Dromia eotvoesi*  
*Calappa heberti*  
*Matuta brochii*

Az alsó-, illetve felső-bádeni faunák közötti különbség akkor a legszembetűnőbb, ha a korallzátonyok rágjait hasonlítjuk össze. Az élőhelyek nyilvánvalóan hasonlóak voltak mindenkor alemeletben: folt-zátonyok kis kiterjedésben, sekély, aránylag védett öblökben. Az uralkodó korallok egyformán *Tarbellastraea* sp. és *Porites* sp. A különbségeket a 2. táblázat mutatja be. Bár a lista nem tekinthető véglegesnek, az eltérés elég jelentős, s feltételezhető, hogy oka rétegtani, vagyis a kör-különbség. A nagyhüflányi (Gross-Höflein, MNH) fauna minden tekintetben átmeneti jellegű az említett két Budapest környéki fauna között. STEININGER és PAPP (in PAPP et al. 1978, p. 199) szerint a korallzátony fölött néhány méterrel települő márga a felső-bádeni emelet legalsó részébe tartozik.

Így az MNH fauna vagy szintén a felső-bádeni alsó részéhez vagy — ami valószínűbbnek látszik — már a középső-bádenihez tartozik. Így a korallzátonyok faunája alapján három biozónát különböztethetünk meg feltételesen (2. táblázat).

Egy negyedik, legfelső zóna létezése nyilvánvaló a Budapest környéki és balatonakali faunák alapján. Ezt az jellemzi, hogy a 3. zónában nagyon gyakori alakok jelentős része hiányzik (*Callianassa munieri*, *C. chalmasii*, *Petrochirus priscus*, *Dromia eotvoesi*, *Calappa heberti*, *Thalamita fragilis* és minden, zátonyhoz kötött faj, mert korallzátonyt ebben a szintben már nem találunk). A harmadik és negyedik zónát az is megkülönbözteti, hogy az alsóban sokszor uralkodó terrigén anyag a felsőből gyakorlatilag hiányzik, ami éghajlatváltozást jelenthet. Legvalószínűbbnek az látszik, hogy a 3. zónát félig száraz, az eróziónak kedvező éghajlat jellemzéte, mely később nedvesbe fordult, s erózió-védő erdő telepedett meg a lejtőkön (KÓKAY — MIHÁLY — MÜLLER in press).

A négy zónát tehát fajok jelenlétével vagy hiányával jellemezzük (3. táblázat) s feltételesen "assemblage zone"-nak tekinthetők. További vizsgálatot igényel, vajon általánosnak tekinthető-e ez a beosztás a Középső Paratethysben, vagy csak helyi viszonyokat tükröz.

# ŐSFÖLDRAJZI KÖVETKEZTETÉSEK

A bádeni emeletre jellemző, gyakori nemzetiségek közül elég sok ma már csak az indo—nyugat—pacifikus faunatartományban él. Ezt már GLAESSNER is hangsúlyozta (1928, p. 205), s később más szerzők is megerősítették (FÖRSTER, 1979a, p. 104, MÜLLER 1979c, p. 865). Az említett mai faunatartomány területéről alig ismerünk miocén rákokat, így a közvetlen összehasonlítás ma még lehetetlen. Az említett jelenség könnyen megmagyarázható úgy is, hogy a ma csak keleten élő nemzetiségek a bádeni emelet után kihaltak a Paratethysból és az ahhoz kapcsolódó Földközi-tengerből és keleti Atlanti-óceánból. A korallzátonyokhoz kötött nemek kipusztulása elkerülhetetlen volt a zátonyképző korallok eltűnésevel. Az említett nemzetiségek miocén jelenléte a Paratethysben alsó-miocén faunakicsérődéssel is magyarázható. Általánosan elfogadott, hogy az eggenburgi korszakban a keleti kapcsolatok még nyitottak voltak. A mai helyzet tehát annak köszönhető, hogy a harmadidőszak végi és pleisztocén lehűlés sokkal jobban érintette az Európa körül vizeket, mint az Ázsiától délre levő óceánokat, ahol tehát refugium, „menedékhely” alakult ki.

Az indo—nyugat—pacifikus tartományban ma is élő, bádeni nemzetiségek a következők: *Matuta*, *Micippa*, *Macrophthalmus*, *Crossotonotus*, *Schizophrys* és *Actumnus* (ezek egy része, a Jóreménységfokát megkerülve, Afrika nyugati partjain is él, de ez a terjedés valószínűleg negyedidőszaki). Néhány nemzetiség elsősorban indo—nyugat—pacifikus, de egy-egy fajjal másutt is képviselve van: *Mursia*, *Trapezia*, *Daira*, *Chlorodiella*, *Carpilius*, *Dorippe*, *Pilumnopaeus* és *Achaeus*. A bizonytalanul meghatározott nemeket (pl. *Pilodius*, *Glabropilumnus*) kihagytam a felsorolásból. Az elsőnek említett hat nemzetisgóból mindegyik ismert a felső-, míg négy csak az alsó-bádeni alemeletből. A második listából (8 nemzetiség) hat felső-, négy alsó-bádeni. Korábbi nézetek ellenére (MÜLLER 1979c) nem bizonyítható az indo—nyugat—pacifikus rokonságú nemek számának növekedése a bádeni emelet során, mert az összes ismert felső-bádeni nem száma (53) nagyobb, mint az alsó-bádeniből ismerteké (36). A keleti rokonságú nemek aránya tehát gyakorlatilag állandó.

Három bádeni faj, vagy nagyon közeli rokonaik előfordulnak a Fidzsi szigetek pliocén rétegeiben (RATHBUN 1945). Ezek közül kettő (*Dardanus hungaricus* és *Daira speciosa*) az 1. és 3. számú zónában található (2. táblázat), míg a *Dynomene emiliae* fajt az 1. és 2. zónában találjuk. Így ezeknek a bizonyítottan indo—nyugat—pacifikus rokonságú alakoknak a száma még valamelyest csökken is a bádeni korszak folyamán.

Mindebből arra lehet következtetni, hogy az indo—nyugat—pacifikus rokonságú alakok egy közös Tethys-fauna utódai, s később haltak ki a nyugati vizekben. Így a bádeni tenger és az Indiai-óceán összeköttetése ma kevésbé látszik valószínűnek, mint néhány évvel ezelőtt. Ennek ellenére nem lehet kizártani egy olyan tengeri kapcsolat léteét, mely néhány euriyalin alak átvándorlását lehetővé tette, de erre faunisztikai bizonyítékunk még nincs. Vitathatatlanul látszik az, hogy a bádeni fauna lényegében azonos maradt a teljes korszak során, s a bádeni tenger óceánihoz közelálló sótartalmát a Földközi-tenger felől beáramló víz biztosította. Sajnos a bádenivel egykorú (serravalli és langhi) mediterrán rák fauna alig ismert. Mégis tudjuk, hogy legalább négy bádeni rák faj előfordul a mediterrán miocénben, míg hat másiknak közeli rokonát találjuk ott.

Egy figyelemremélő tény magyarázatával még várni kell. Kizárálag a 3. számú zónától kezdve (2. táblázat) található néhány nagyon gyakori faj: *Petrochirus priscus*, *Dromia eovoesi*, *Calappa heberti*, *Matuta brocchii*, *Micippa hungarica*, *Thalamita fragilis*, *'Pisa' oroszji*, *Actumnus telegdi*. Gyakoriságuk valószínűvé teszi, hogy nagy résziük tényleg hiányzik az alsó-bádeniből, nem csupán annak kevésből ismert volta miatt maradt ki a listákból. Ez fajok nagy része eltűnik a 3. és 4. zóna határáról, mások ritkábbá válnak. Egy lehetséges magyarázat szerint e fajok érzékenyen voltak a klímára vagy a sótartalomra, ezért haltak ki a 4. zóna kezdetén, a feltételezett éghajlatváltozás hatására. Bevándorlásuk feltételezése szükségessé teszi eredetiük helyének feltételezését is. Ez a hely aligha lehetett az indo—nyugat—pacifikus területen, hiszen nagyrészt sztenohalin fajokról van szó, melyek onnét nem vándorolhattak át, mert a jelenlegi ősföldrajzi ismereteink szerint a köztes tengerrések részben csökkent-, részben túlsós vizűek voltak. Eredhettek volna a Földközi-tenger délebbi területeiről is, de onnét eddig csak a *Petrochirus priscus* fajt (GLAESSNER 1928) ismertették közülük.

# ÖKOLÓGIAI KÉRDÉSEK

A bádeni emelet különböző típusú rák-együtteseket adott. Ezeket szubjektív módszerekkel is csoportosíthatjuk. A csoportok valószínűleg elsősorban az egykor környezet körülményeit tükrözik. Objektívebb csoportosítást kaphatunk analitikus módszerekkel. RAJČÍ M. végezte a cluster-analízis számításait. Az egyes együttesek (lelőhelyek, ill. rétegek) hasonlóságát bináris (jelenlét-hiány) és kvantitatív függvények segítségével számítottuk. Kvantitatív módszereket ritkán alkalmaznak a makropaleontológiában, főleg akkor, ha nem szitált vagy iszapolt, hanem kézzel gyűjtött anyagról van szó, hiszen az ilyen minták reprezentatív voltában joggal kételekedhetünk. Az alább bemutatott megfontolások és az eredmények meggyőző volta azonban bizonyítani látszik a kézzel gyűjtött minták számszerű kiértékelésének jogosultságát is.

Még a legnagyobb gondossággal végzett kézi gyűjtés esetén is fellép egyfajta statisztikai torzítás. Al gyűjtő a kőzetet adott átlagmérétű darabokra töri. A kőzettaraboknál kisebb kövületek rejte maradhatnak, ha a darabok belsőjében vannak. Ezért a kisebb példányok megtalálásának valószínűsége csökken. Tegyük fel azonban (ez a rákok esetén megengedhető, mert a maradványoknak csak elenyésző tört részét teszik), hogy a kőzet a fossziliák helyétől független helyeken törik el, így annak valószínűsége, hogy egy kövületet észreveszünk,

$$p = 1 - \frac{(r-f)^3}{r^3} \quad (p_{\max} = 1)$$

esz, ahol  $p$  az  $f$  méretű maradvány megtalálásának valószínűsége  $r$  méretű darabokra tört kőzetben, feltéve, hogy minden a kövületek, minden a kőzettarabok nagyjából izometrikusak. Ezzel a képlettel korrigáltuk a szitált minták darabszámaikat, hogy azok a (jóval számosabb) kézzel gyűjtött minta darabszámaihoz hasonlóan torzuljanak. Így az eloszlási függvények már összehasonlíthatóvá váltak. Bár egyfajta torzítás hatását így redukáltuk, mások elkerülhetetlenül megmaradnak, pl. szelektív beágyazódás, fosszilizálódás, téves határozás következtében.

Az így korrigált adatokat használtuk a hasonlóság kiszámítására, bináris (SØRENSEN 1948, SOKAL—MICHELER 1958) és kvantitatív (HORN 1966, CZEKANOWSKI 1909, PINKHAM—PEARSON 1976, HUMMON 1974) egyenletekkel. A csoportképzéshez súlyozott átlagszámításos módszert használtunk. A számítások alapjául először a nemzetiségek, majd a fajok példányszámaikat, ill. jelenlétét—nemlétét vettük. E két számításnál teljesen azonos dendrogramokat kaptunk, ami érthető, mert a legtöbb nemzetiség csak egy fajjal szerepelt.

A cluster-analízis fő célja az volt, hogy az egyedi lelőhelyekből, illetve rétegekből csoportokat képezzünk, melyekben a hasonló faji összetétel hasonló ökológiai viszonyokra enged következtetni. A csoportokból kiválasztottuk a leghűségesebb és legállandóbb fajokat s ezek mai rokonainak ökológiai igénye alapján következtettem tovább. Nyilvánvaló, hogy egy-egy csoportra a legjellemzőbbek a húséges és állandó alakok. A csoportokat vagy halmazokat (clusterek) úgy nyerhetjük, hogy a számítás eredményeképpen kapott dendrogramokat (12. ábra) egy adott szinten vízszintes vonallal elvágjuk. E szint megválasztása bizonyos mértékig önkényes, ezért a módszer sem mentes szubjektív elemektől. Ezt a szubjektivitást erősen csökkenti, ha az egymástól eltérő módszerek végeredménye hasonló. Négy csoportot kaptunk a SØRENSEN, HUMMON, SOKAL—MICHELER és HORN-módszer alapján, s öt, többé-kevésbé homogén csoportot a PINKHAM—PEARSON és a CZEKANOWSKI-függvénytel (11. ábra, 4. táblázat). További hierarchikus szintek nem mutatkoztak, sem alacsonyabb, sem magasabb szinten. Az előzetesen legkohérencebbnek ítélt zátonyokat (MRZ csoport, lásd alább) a függvények többsége egybe sorolta, a PINKHAM—PEARSON és a SOKAL—MICHELER-módszer mégis kettéosztotta. Négy eltérő módszer — a SØRENSEN, CZEKANOWSKI, HORN és HUMMON-függvény — csaknem teljesen azonos eredményt adott. Csupán néhány lelőhely, illetve réteg helyzete változott a módszertől függően, ezek: MKC, WEG és MDN. Ezek közül az első valóban átmeneti jellegű, míg a másik kettő mintája nagyon kevés elemből állott, mivel ezek kevessé begyűjtött lelőhelyek. Az eredmények

jól egybevágnak a kőzetek üledéktani jellegeivel, a szemnagysággal, de főleg az onkoidok meglétével vagy hiányával és a zátonyszerű kifejlődésekkel, valamint a terrigén anyag mennyiségével (4. táblázat).

A legegységesebbnek mutatkozó csoport (MGY) még az MRO, MRF, MR8, MR45, MA, MOH, MEG és MFA rétegeket tartalmazza (a csoportnak az egyik, jellemzőnek ítélt elem jelét tartottam meg), s a dendrogramok egy részében az MKC is idekerült. Az alsó-bádeni Mátraverebély–Szentkút és Nyírád (MNU) lelőhelyek is hasonló együttest adtak, melyben a *Calappa heberti* a *C. praelata* helyettesíti. Ezek mintának kis elemszáma miatt nem vontuk be őket az analízisbe. Ezeket az együtteseket a '*Callianassa*', *Calappa*, *Matuta*, *Mursia*, *Thalamita*, *Dromia eotvoesi* és *Portunus* (*Portunus*) taxonok gyakorisága jellemzi, míg *Albunea* és *Portunus* (*Monomia*) csak egy-két helyen került elő. Ezek a taxonok húségesek, az első hat pedig állandó is, mert minden egyes elemben előfordul.

A '*Callianassa*' fajok sokasága (DE SAINT LAURENT–LE LEUFF 1979) a 10–30 méter közötti vízmélységek tulajdonítható, mert a ma élő *Callianassa* és *Callichirus* fajok ezt a mélységet kedvelik, bár egyesek előfordulnak az árapályövben, mások pedig 100 méternél mélyebben is. A helyek egy részénél egy nagy méretű faj (*C. munieri*) gyakorisága homogén üledéket jelez. A *Matuta* fajok zöme sekély vízben él (SAKAI 1976), de a bádeni fajhoz leginkább hasonló *M. curtispina* 35–80 m között, homokos fenéken él. A *Mursia* fajok lágy aljzatot kedvelnek, 50–150 m mélységen (SAKAI 1976), de a *M. cristimana* fajt 17 méterből is jeleztek (BARNARD 1950). A *Thalamita fragilis*-hez hasonló *Th. crenata* egészen sekély vízben él, iszapon vagy lágy homokon. Más fajok 80 méterig szállnak le (SAKAI 1976). Az egyes *Calappa* fajok eltérő mélységet kedvelnek, 10–150 méter között (SAKAI 1976). A *Calappa praelata*-hoz hasonló *C. granulata* 30–150 m között él (ZARIQUIEY ALVAREZ 1968). A *Portunus* (*Portunus*) fajok sekély vízben élnek, 20–30 méterig. Az *Albunea carabus* lágy homokos fenéken él, általában 30–50 méterig (ZARIQUIEY ALVAREZ 1968).

Az adatok aktualisztikus értelmezése szerint a csoportba tartozó rétegek zöme 20–30 méter mély vízben keletkezett. Ahol az üledék homogén volt (MRF, MR8, MR45), ott a nagy *Callianassa* fajok domináltak, másutt a kis. *C. pseudorakosiensis* élt tömegesen (MFA, MFK). A *Calappidae* fajok sokfélesége és gyakorisága normális sótartalmat, meleg–mérsékelt éghajlatot jelez. Az *Albunea* időnkénti gyakorisága (MGY, MRF) 30 méternél valamivel mélyebb vizet jelenthet. A csoport rétegei néhány rákfajt megesztnak más csoportokkal, főleg az MOE-vel (lásd alább). Ezek közül a *Maja* a legfontosabb, melynek mélységgénye 15–30 m közötti, ritkán 10–200 m között él (SAKAI 1976). Az európai *Maja squinado* 2–73 m között él (INGLE 1980). A *Dorippe* fajok 15–50 m között (SAKAI 1976) vagy 40–100 m között (ZARIQUIEY ALVAREZ 1968) élnek. A *Liocarcinus* fajok a parttól 220 méterig terjedő zónát osztják fől egymás között (INGLE 1980).

Az MOE csoport még az MDG, MDN réteget tartalmazza, valamint a klasszifikációk egy részében a DAL és WEG jelűt is. A csoport jellemző alakjai, melyek állandóak és húségesek: *Micippa hungarica*, *Ebalia* fajok, '*Lissocarcinus*', *Liocarcinus*. A jobban begyűjtött lelőhelyeken ehhez társul az *Achaeus*, *Actumnus*, *Dorippe*, *Maja*, *Parthenope*, *Cancer*, *Xantho*, *Palicus*. A *Micippa hungarica*-hoz közelálló *M. thalia* 20–100 m között él, homokon vagy héjtörmeléken (SAKAI 1976). A legtöbb *Achaeus* faj 10–50 m között él, de egyesek sekélyebb vízben is előfordulnak. Az *Ebalia* fajok nagyon változó mélységből kerültek elő: kettőtől több száz méteres mélységgig (INGLE 1980). A *Palicus* fajok általában 50 méter alatt élnek (SAKAI 1976, ZARIQUIEY ALVAREZ 1968). A *Cancer* fajok a parttól 1000 méterig találhatók (RATHBUN 1930). A *Parthenope* fajok 10 és 40 m között gyakoriak (ZARIQUIEY ALVAREZ 1968). A mai *Actumnus* fajok általában mélyebben élnek, mint a *Pilumnus* alakok, általában 20–35 m alatt, de két alak árapályövben él (SAKAI 1976). Ezek az adatok a gyakoribb és állandóbb fajok adatait nagyobb súlyjal véve, 20 méternél valamivel nagyobb vízmélységet sugallnak, hasonlóan az előző csoporthoz. Lényeges különbség a nagy és közepes méretű *Callianassa* fajok hiánya, amit az inhomogén üledéknak tulajdoníthatunk. A *Calappa*, *Dromia*, *Thalamita*, *Portunus* fajok hiánya, a *Matuta* és *Mursia* ritkasága lehűlést és valamelyest változó sótartalmat jelez, mert ezek a nemek ma elsősorban trópusi és meleg–mérsékelt zónákban, rendes sótartalmú vízben élnek. Mint említettem, az MOE csoportot terrigén anyag hiánya (18. old.), ami éghajlatváltozást jelenthet.

Az MB jelű csoport nagyon érdekes, mert olyan környezetet képvisel, ami fosszilisan nagyon ritkán marad meg. A kőzetek minden tartalmaznak onkoidokat, vagy szstromatolitokat (LELKES–MÜLLER in press). Ez nagyon sekély vizet jelez. A következő egységek tartoznak ide: MRS, MGF, MB, MBK, MBH, MJL, MM és fenntartásokkal a DAL és WEG. Jellemzője a *Pisidia kokayi*, '*Pisa*' oroszyi és *Pachygrapsus hungaricus*, melyek ritkák másutt, a legtöbb csoportban teljesen hiányoznak. Jellemző még a *Brachynotus februarius* kizárolagos jelenléte, valamint a *Pilumnus mediterraneus* és *Xantho moldavicus* gyakorisága. Az utolsónak említett két faj kivételével ezek mind húséges alakok, s meglehetősen állandóak is, tehát az együttesre nagyon jellemzőek. A mai *Pachygrapsus* fajok a szikláslás árapályövre jellemzőek (PÉRÈS–PICARD 1964), ami ezt a leletet különösen érdekessé teszi. A mai *Brachynotus* fajok leginkább 0,5–2 m mélység között találhatók (ZARIQUIEY ALVAREZ 1968). A legtöbb *Pisidia* faj szintén ilyen körülmények között él, de néha mélyebben is előfordul. Mindhárom említett alak bűvöhelyeket igényel: sziklahasadékokat, vagy kövek alatt, *Upogebia*-járatokban keres

menedéket (PESTA 1918). A *Pisa* fajok 1 és 90 méter között élnek, kedvelik a szilárd aljzatot (INGLE 1980). A *Xantho* és *Pilumnus* fajok meglehetősen igénytelenek, nagyon sekély vízben is előfordulnak. A fentiek szerint az MB típusú rétegek nagyon sekély, parti sávban rakódtak le. Az onkoidok és szstromatolitok szilárd aljzatot jelentettek s bűvöhelyet is. Időnkénti regressziók, tengerszint-ingadozások következtében sziklás-köves árapályövvé vált ez a terület.

A rákok mai rokonai zömmel euryhalin szervezetek: *Xantho*, *Pilumnus*, *Brachynotus*, *Pisidia*, *Liocarcinus*, *Pachygrapsus* fajok ma is élnek a Fekete-tenger félénk vizében (PESTA 1918), még néhány *Leucosidae* faj áthatolt a Szuezi-csatorna túlsós vizein, mások pedig a Márvány-tengerben élnek (HOLTHUIS—GOTTLIEB 1958, POR 1978). Ez összhangban van azzal a ténytel, hogy az árapályöv közelében minden változó a sótartalom, de különösen így lehetett ez a bádeni emelet vége felé, amikor a Keleti Paratethys csökkent sótartalmú vizének beáramlását kell feltételezni (KÓKAY in KÓKAY—MIHÁLY—MÜLLER in press).

Az MRZ csoport (korallzátonyok) az MDZ, MK, MV, MTZ és MSH helyekből áll. A zátonyépítő korallok jelenlétéből a kőzettípus azonnal felismerhető. Ezek részben élő helyzetben, részben kimosva temetődtek be, de mennyiségeük mindegyik helyen kőzetalkotó. A betemetődés módja néha egészen rövid távon, néhány méteren megváltozik (lásd az MK lelőhely leírásánál). Egész sor nemzetseg és faj kizárolag ezeken a helyeken fordul elő: minden *Chlorodiella* faj, *Trapezia*, *Carupa*, *Rakosia*, *Daira speciosa*, *Carpilius*, *Charybdis* (*Goniosupradens*), *Dynomene*, *Dromia neogenica*, *Actaea*, 'Polidius', *Haydnella*, *Crossotonotus*, *Schizophrys*, *Petrolisthes*. (Ritkaságként egy-egy példány *Chlorodiella mediterranea* és *Dromia neogenica* került elő az MKC rétegből, mely bizonyos mértékig átmeneti jellegű.) A felsorolt nemek mai képviselőinek többsége zátonylakó: *Charybdis* (*Goniosupradens*), *Daira perlata*, *Chlorodiella*, a legtöbb *Dynomene* faj, *Carpilius*, *Trapezia* (SAKAI 1976, TAYLOR 1968). Mások nem kizárolagos zátonylakók, de szilárd aljzatot igényelnek, ezért zátonyon is élnek (*Schizophrys*, *Carupa*, *Daira americana*, *Crossotonotus*). Kevésbé érthető a *Panopeus* és *Cancer* jelenléte. A mai *Panopeus* fajok jó része lágy aljzatot kedvel, de alkalmilag zátonyon is előfordul, ezzel szemben *Cancer* fajt zátonyról nem jeleztek, elterjedésük sem esik egybe a zátonyokkal (RATHBUN 1930). *Calappa heberi* is került elő ezekről a lelőhelyekről, mai *Calappa* fajok is élnek alkalmilag zátonyon. Jellemző, hogy a másutt gyakori alakok közül néhány következetesen hiányzik a nagyon gazdag, jól begyűjtött MRZ típusú anyagból: *Parthenope*, *Mursia*, *Leucosiidae*. Ezek mai képviselői ritkák a zátonyokon, vagy teljesen hiányzanak (TAYLOR 1968). Két helyen (MDZ és MRZ) a *Pisidia kokayi* és *Pachygrapsus hungaricus* is előfordul, jelezve, hogy ezek a zátonyok időnként az árapályövbe kerültek, amit üledéktani bélyegek (gyökérnyomok és száradási repedések) is jeleznek az MRZ lelőhelyen. A korallzátonyok jelenléte óceánihoz közelálló sótartalmat és meleg—mérsékelt vagy trópusi éghajlatot jelent. A korall nemzetsegégek számának csökkenése a bádeni emelet során (mintegy 20-ról négyre vagy ötre, lásd 2. táblázat) arra utal, hogy a sótartalom a felsőbb részek keletkezésekor már nem volt minden teljesen állandó, de az éghajlat romlására is utalhat. A zátonyok mérete viszont a felső-bádeni emeletben sem kisebb, mint az alsóban, ami inkább a sótartalom változásának hatását támasztja alá.

Különleges együttest ismertem meg legújabban az MZZ (Zebegény) rétegen (6. ábra). Ez még nem került be a cluster-analízisbe. A finomszemű mikrites kőzet egy kis nem-zátonyépítő, ágas-bogas telepes korallfaj tömegeit tartalmazza, mely *Oculina*-ra emlékeztet. Ahermatipikus korallból álló mezőket a legkülönbözőbb mélységekben és klímákon találunk, egy métertől több száz méter mélységig, a trópusoktól a hideg-mérsékelt övig. A ?*Porites* genus jelenléte viszont azt jelenti, hogy itt a víz nem lehetett mélyebb húsz—harminc méternél. A rák fauna hasonlít a zátonyokéhoz, de kevesebb fajt tartalmaz. Érdekes a két *Ebalia* faj jelenléte, mert a zátonyokon egyetlen *Leucosiidae* fajt sem találtam. A *Portunidae* és *Dromioidea* fajok viszont hiányzanak a harminc példányból álló mintából.

A fenti négy típuson kívül egy ötödik körvonalait sejteti néhány szórványelelet. Több lelőhelyen került elő *Goneplax* faj, Oriakhovóban (Bulgária) *Miocyclus bulgaricus*-szal, Soossban (Baden mellett) „*Cancer*” szontaghi *vindobonensis*-szel együtt. Egy *Geryon* fajt adott a budapesti Z-2. sz. fúrás. Mind ezek agyagból vagy finom aleuritból származnak. A recens *Goneplax rhomboides* 40—760 m között, a *Geryon tridens* 770—1500 m között él az Adriában (PESTA 1918). *Ranidina rosaliae* két helyről került elő, Borbolyán „*Cancer*” *bittneri*-vel együtt. A *Raninidae* fajok zöme is mély szublitorális életet él. A felsorolt leletek tehát mélyebb szublitorális és sekélybatiális körülmenyekre utalnak.

Bár természetesen sírközösségeket vizsgáltunk, a rákok páncéljának törékenysége és az eredmények aránylag ellentmondás-mentes volta valószínűvé teszi, hogy ezek jól tükrözik az egykor élőhelyek közösségeinek összetételét.

A fejezetben ismertetett következtetések elég jól egyeznek más gerinctelenek vizsgálatával kapott eredményekkel (lásd a lelőhelyek leírását) és az üledéktani megfontolásokkal. Ilyen vizsgálatok még kevés helyen történtek, ezek eredményeit több cikk tárgyalja (MÜLLER 1974a, 1974b, 1976a, 1979a, KÓKAY—MIHÁLY—MÜLLER in press, LELKES—MÜLLER in press).

Szintén RAJCGY M. segítségével kaptuk meg a minták SHANNON-féle diverzitási („sokféleségi”) indexeit. Az egyes csoportokra vonatkozó szélső értéket  $H'$  log<sub>2</sub> formájában a következőkben adom:

MGY csoport:  $H' = 2,18 - 3,66$

MOE csoport:  $H' = 3,25 - 4,00$

MRZ csoport:  $H' = 2,10 - 3,63$

MB csoport:  $H' = 1,24 - 2,83$

Néhány árapályövi és sekély infralitorális tízlábú rák-közösség diverzitását ABELE (1974) vizsgálta. Bár ő a mintákat a meleg-mérsékeltövi és trópusi tengerekből vette — melyek hőmérséklete valószínűleg magasabb, mint amilyen a bádeni tengeré lehetett —, legmagasabb értéke ( $H' \log_2 = 4,346$ ) nem lényegesen nagyobb, mint a mi általunk kapott maximális adat. Igaz ugyan, hogy a fosszilis minták több élőhelyből összesen példányokat is tartalmazhatnak, ami megnövelheti a fajszámot, de ez a hatás a rákoknál valószínűleg elhanyagolható.

A diverzitási érték érdekes egyirányú változását figyelhettük meg a Tétényi-fennsíkon. Az MB csoport lelőhelyein belül a diverzitás észak felől (MM réteg,  $H' = 1,24$ ) dél felé (MBH réteg,  $H' = 2,93$ ) növekszik, majd még tovább dél felé, az MDG lelőhelyen még nagyobb ( $H' = 3,25$ ) értéket kaptunk. E rétegek valószínűleg egykorúak, ezért az is valószínű, hogy a  $H'$  növekedése a növekvő vízmélységet tükrözi, egy, a partra nagyjából merőleges szelvényben. Ez az eredmény nemileg ellentmond ABELE (1974) egyik megállapításának. Ő ugyanis az egyik legmagasabb értéket a sziklás árapályövre kapta, míg a mi esetünkben a sziklás árapályövi MM előhely értéke éppen a legalacsonyabb. ABELE viszont csupán egyetlen infralitorális együttest vizsgált, egy *Pocillopora*-asszociációt, melynek diverzitása is nagy volt ( $H' = 4,346$ ). Ha a bádeni együttesekre kapott értékek megbízhatónak bizonyulnak a kis elemszámú minták adta határon belül, akkor megállapíthatjuk, hogy nem a korallzátonyok rákjaira kaptuk a legmagasabb értékeket (MRZ csoport), hanem egy infralitorális csoportra, az MOE jelűre, mely erősen összetett aljzaton (mészhomokkő, osztriga-padok, ép kagylóhéjak, héjtörmelek) élt együtteseknek felel meg. ABELE végkövetkeztetése az volt, hogy a diverzitási érték legjobban az aljzat összetett voltával, komplexitásával korrelál. E tekintetben tehát eredményeink egybehangzanak az övéivel, mert az MOE típusú sokrétű aljzat komplexitás tekintetében egyenértékű lehet egy korallzátonnyal.



**DECAPOD CRUSTACEA OF THE BADENIAN**



## SUMMARY

120 species and subspecies of anomuran and brachyuran decapods have been found in Badenian (Middle Miocene, Central Paratethys, Central and Eastern Europe) layers of which one hundred were determined to the species level. Four new genera, 21 new species and one new subspecies are described. The new genera are: *Kromtitis* (*Dromiidae*), *Corystites* (*Corystidae*), *Rakosia*, *Miopipus* (*Portunidae*) and *Haydnella* (*Xanthidae*). The new species or subspecies are: '*Callianassa*' *szobensis* (*Callianassidae*), ?*Pagurus turcus*, '*Pylopagurus*' *leganyii*, *Diogenes matrensis*, *Paguristes cserhatensis* (*Paguridae*), *Petrolisthes magnus*, *P. haydni* (*Porcellanidae*), *Schizophrys visegradensis* (*Majidae*), *Parthenope tetenyensis* (*Parthenopidae*), *Portunus (Monomia) miocaenicus*, *Charybdis (Goniosupradens)* *mathiasi*, *Rakosia carupoides*, '*Xaiva*' *bachmayeri* (*Portunidae*), *Actaea turcocampestris*, *Chlorodiella mediterranea tetenyensis* (n. ssp.), *Chlorodiella loczyi*, *Ch. juglans*, *Haydnella steiningeri*, *Panopeus wronai*, *Pilumnopeus paratethyensis*, '*Pilumnopeus*' *tetenensis* (*Xanthidae*) and *Crossotonotus diosdensis* (*Palicidae*). The decapod associations were assigned to four stratigraphic zones which could be correlated tentatively to substages of the Badenian. Primarily by cluster-analyisis, these associations were divided into five clusters reflecting differing paleoecological conditions. Rocky intertidal, reefal, shallower and deeper infralittoral and circalittoral habitats were recognised.

## INTRODUCTION

The decapod crustacea represent a major and important element of marine biota. Many of these organisms have well calcified crusts. In spite of this the fossil record reflects their importance only very incompletely. In any of the recent tropical seas their species number several hundreds. Even in a warm-temperate sea, e.g. the Mediterranean some two hundred species are present, from which ninety belong to the group brachyura.

In contrast to this only few decapod localities, generally of tropical origin, yielded more than 20 or 30 fossil species. Such outstanding localities are mostly Mesozoic or Eocene. Most Neogene formations yielded only sporadic specimens. Rich associations are unfrequent. The Badenian of the Central Paratethys is one of the exceptional cases. The unusually high number of detected species, over hundred, may be attributed to the long tradition of study of fossil crabs in this area as well as to the good localities that are situated in the vicinity of the university towns, Budapest and Vienna. A need to revise previous publications and the availability of new material motivated our study.

The wealth of material permits the description of whole communities of Badenian decapods which can be correlated with paleoecological conditions. The factors controlling these associations will be tentatively characterized.

As far as this was possible the type- and other specimens of previous studies were located and revised. I visited several times some easily accessible and promising localities, e.g. Rákos, Órs vezér tere, Tétény, Gross-Höflein, which yielded rich and ecologically evaluable associations. Other, less promising or more remote localities, e.g. Mátraverebély-Szentkút, Zebegény, Szob, Deutsch-Altenburg or Grobie were visited only once or twice.

Inevitably, the material is somewhat one-sided. Upper Badenian localities yielded many times more specimens than Lower or Middle Badenian ones. In spite of this, a monographical evaluation of the whole known Badenian decapod fauna seemed to be warranted.

Several thousands of specimens (isolated carapaces, chelae, rarely entire tests) were studied and determined. 120 different forms were recognized, of which 102 were determined to species or even lower level. Most of the specimens were found in more or less rich associations. Four new genera, 21 new species and one new subspecies were recognized.

Our aim was the description or revision of all known Badenian decapods with the exception of isolated fingers the study of which is generally not very promising at the present level of our knowledge. To promote such studies in the future, however, I tried to identify isolated fingers too, if these were found in layers where carapaces and chelae were also present.

The associations, and in some cases the individual findings too, were evaluated ecologically, with the aid of cluster-analysis, taking into account the ecological requirements of recent forms allied to the Badenian ones. The Leythakalks, yielding the bulk of the decapods, were among the less studied Badenian deposits. In many cases it was only the crabs which gave evidence about paleo-ecological conditions. In some instances, the molluscan and foraminiferal faunas were elaborated by colleagues yielding further information about environments.

The associations were tentatively evaluated from paleogeographic and stratigraphic points of view as well.

### Acknowledgements

Before all, I owe thank all of the members of my family who tolerated with patience the mess and disruptions in our home for which my studies on fossils were responsible. My son, MIKLÓS helped with photography, my wife ÉVA, my brother MIKLÓS, my cousin, DR. GÁBOR SZIKLA, my aunt, MRS. ÉVA MARSOVSZKY supported my trips to foreign localities or museums.

In the Geological Institute, especially DR. GÉZA HÁMOR and GYÖRGY ZSILÁK supported this work in several ways. My colleagues, DR. SÁNDOR MIHÁLY, MRS. MARGIT PAPAJCSÍK and MR. PÉTER SOLT helped me either in field-work or by locating material in collection. I received similar assistance from DR. TIBOR KECSKEMÉTI, LÁSZLÓ

FORRÓ and HENRIK FARKAS of the Natural History Museum, and from DR. JENŐ BODA of the University of Budapest. I am indebted to many friends who brought me specimens of crabs from various seas, DRs. MARGIT H. DEÁK, ISTVÁN SÁRVÁRY, ÖDÖN RÁDAI, GYULA RADÓCZ, Mrs. ZSUZSA MAROS, MR. GÁBOR MAROS, and others. I express my gratitudes for advices and gift of material for experiments of casting technics to DRs. ATTILA Vörös and FERENC CSER and also for precious advices about photography to MR. ISTVÁN FÖRSÖNITS JUN. My friends and colleagues, DRs. JÓZSEF KÓKAY and GYÖRGY LELKES provided many unpublished results of their field so useful in this work. DR. JÓZSEF KÓKAY also gave me material he collected, in fact some of the most valuable crabs discussed were obtained through his courtesy. DR. MIKLÓS RAJCZY helped in a generous way by giving advice and carrying out numerical computations in cluster analysis and of diversity indices. I am especially indebted for the extremely careful review of my manuscript to DR. MIKLÓS MÜLLER, my brother and to DR. ENDRE DUDICH.

Sincere thanks are due to DR. FRITZ STEININGER for those precious days he devoted to helping my field-work in Burgenland, Austria. DR. FRIEDRICH BACHMAYER helped me in visiting a locality in Austria and the Natural History Museum of Vienna. DRs. LORETTE WITTIBSCHLÄGER and ORTWIN SCHULZ helped me to locate or to study type-specimens in Austrian collections. I am indebted to specialists for providing me with specimens of extant crabs or advices, especially to DRs. DANIELLE GUINOT and L. B. HOLTHUIS who both helped to elucidate systematical problems. Similarly I thank for sending specimens or casts to DRs. J. S. H. COLLINS, JOHN GARTH, ARIE JANSSEN, PIERRE MOISSETTE, JEAN-PAUL SAINT MARTIN, A. FENNINGER, G. E. GROOT and others. DRs. EMILIA KOJUMDGIEVA and WILHELM KRACH conveyed fine specimens for study from the Badenian of their countries. I am grateful to DRs. LUIS VOÀ BOADA, SYLVIE SECRETAN and R. V. INGLE for advices or other helps.

The Badenian stage is a regional stratigraphical unit. Its applicability is confined to the Central Paratethys. Both terms were defined in PAPP et al. 1978. The lower boundary of the Badenian coincides with the appearance of the planctonic foraminiferan *Praeorbulina Olson*. If this event was synchronous everywhere, the beginning of the Badenian should correspond to that of the Langhian, Mediterranean stage, a part of the Middle Miocene.

The upper boundary of the Badenian is less clearly defined because the Paratethys separated, probably gradually, from the oceans which led to enhanced endemism of the biota. Therefore this boundary was defined as the appearance of the highly endemic Sarmatian fauna and the extinction of the euhaline elements. The synchronicity of this event is in question and its global correlation is also problematic. According to PAPP, SENEŠ and STEININGER (in PAPP et al. 1978, p. 57) this event can be correlated to the Middle Serravallian, Mediterranean stage. It is still worth remembering that an erroneous correlation of the Badenian to the Tortonian was generally accepted in the period from 1928 to about 1968, introducing an error of some five or six millions of years. Although in the earlier literature there was a notable trend toward a more correct correlation (e.g. LÖRENTHEY 1909), SCHAFER (1927) placed the layers in question in the Tortonian stage without providing any evidence. Unfortunately, his view was generally accepted with only a few exceptions (e.g. STRAUSZ 1966, who continued to use the old term Upper Mediterranean Stage). This discrepancy made it difficult to correlate the Badenian to other European or Global stages for half a century. The term Tortonian (for Badenian layers) keeps appearing occasionally in recent papers.

The Badenian period extends from 13.3 to 16.5 million years B. P., according to VASS, BAGDASARYAN and STEININGER (in PAPP et al. 1978, pp. 35–46). These dates are based on local radiometric data and on their correlation to the global radiometric scale.

The Central Paratethys was an inland sea that occupied the foredeep and the inner-arc basin of the Carpathians, but also extended toward the Alps and the Dinarids and around the Northern slopes of the Balkan Mountains. The Paratethys was connected with the western Mediterranean by a molasse-deep north of the Alps. This connection has definitely closed before the beginning of the Badenian. A connection to the eastern Mediterranean through the Palaeo-Adriatic Sea might have existed until the end of the Badenian.

The Eastern Paratethys was an extended system of seas reaching as far as the area of the recent Balkhas Lake. It was probably connected with the Central Paratethys during the whole Badenian but striking differences in the facies and biota of these seas argue for their high grade of independence.

The oceanic salinity of the Badenian sea indicates that one or more open sea-ways to oceans permitted the outflow of the diluted or concentrated water of the sea and the simultaneous influx of oceanic water. The almost euhaline character of the Badenian Sea is supported by the diversity of corals, echinoids and many other organisms though a deviation of the salt-content within the limits of a few permilles was evident in the Middle Badenian and in the second part of the Late Badenian at least. The connections might correspond to one or more of the following:

1. to the Atlantic, through the Mediterranean and the Proto-Adriatic Sea,
2. to the Indian Ocean, through the Eastern Paratethys and through a hypothetical through running parallel to the Zagros-Chain,
3. to the Indian Ocean, through a highly hypothetical molasse-sea running along the Hellenids, Taurids and Zagros.

According to the correlation of Eastern Paratethyan stages to their western counterparts, as proposed by PAPP and SENEŠ (in PAPP et al. 1978, p. 53), the Tarkhanian of the Eastern sea corresponds to the Lower Badenian (Moravian). In Tarkhanian faunas an impoverishment of the biota indicates a slightly brachyhaline character of its water. During Tshokrakian and Karaganian, corresponding to the uppermost Moravian and the Middle Badenian (Wieliczian), this impoverishment is even more accentuated indicating a gradual closing of the connection with the oceans. During the

Konkian of the Eastern Paratethys, which might correspond to the Upper Badenian (Kosovian), a new but restricted marine invasion indicates a stabilised salinity which was not as high as that of the Upper Badenian Sea, however.

Based on the above mentioned hypothetical ideas it is highly probable that a firm euhaline connection of the Central Paratethys and the Mediterranean existed during the whole Badenian. The character of the biota of these seas is essentially the same. The possibility of an Indo-West-Pacific connection through a partly brachyhaline, partly hypersaline sea-way can not be ruled out for the Late Badenian, however (RÖGL-STEININGER-C. MÜLLER 1978). The persistence of a western link remains doubtless, however.

The Miocene Paratethys was in a crucial location during the separation of the Indo-West-Pacific and East-Atlantic-Mediterranean faunal provinces. The last massive faunal exchanges between the both realms might have taken place during the Early Miocene but some episodic and restricted connections might have occurred during the Late Middle Miocene, too.

Some of the earliest papers mentioning Badenian Crabs describe sporadic findings. The very first paper of this kind mentiones the occurrence of the Mediterranean species *Lobocarcinus sismondai* (= *Cancer s.*) in the Vienna Basin (A. E. REUSS 1859). This was followed by the description of *Microdium nodulosum* from Wieliczka (A. E. REUSS 1867). Some years later R. REUSS (1871) published a paper on *Phymatocarcinus speciosus* (= *Daira speciosa*), based on an internal mould of the crab from Baden Rauchstallbrunnengraben.

BITTNER's paper (1877) was the first to mention more than one species from a single locality, giving a detailed description of *Phymatocarcinus speciosus* from Gamlitz, Steiermark; he also mentiones *Carpilius* sp. and *Titanocarcinus* sp. (probably *Xantho*) but without description. From other localities he mentiones a chela of a portunid and *Macrophthalmus* sp. This latter came from a Carpathian layer.

In 1883 BROCCHI published a paper about the Rákos (now part of Budapest) fauna. He described six new species, *Pagurus priscus* (= *Petrochirus pr.*), *Calianassa munieri* (= *Callianassa m.*), *C. chalmasii*, *Calappa heberti*, *Matuta inermis* (= *M. brocchii*) and *Portunus pygmeus* (*Miopipus p.*) and also presented an illustration of a portunid (in French: Portunien) chela (= *Thalamita fragilis*). Unfortunately the Rákos section comprises several layers which are not indicated in BROCCHI's paper. As the type specimens are probably lost forever, we do not even have the matrices to indicate what layers contained the specimens. A year later BITTNER (1884) published a paper on crabs from strata of various geologic ages, including the following Badenian forms: *Cancer styriacus*, *C. illyricus* and *C. carniolicus*. BITTNER later (1893) described several crabs, including Badenian ones, from the Tertiary of Pannonia (meaning Hungary and Croatia together at that time). He mentioned *Neptunus* cf. *granulatus* (? = *Portunus monspeliensis*) from the vicinity of Kolozsvár, Transylvania. From Walbersdorf (Borbolya, now Burgenland, Austria) a new species, *Ranidina rosaliae* is described and specimens of *Cancer* cf. *illyricus* (? = "Cancer" bittneri) are enumerated.

IMRE LÖRENTHEY, Hungarian paleontologist and geologist (in papers published in German language he used his name's German version, EMERICH LÖRENTHEY) was one of the most successful investigators of Tertiary crabs. He focused his efforts, however, on the diverse and well preserved Upper Eocene fauna of Kissvábhégy (now Martinovics-hegy), Budapest and much less on the Badenian ("Obermediterran" in those times) decapods. Nevertheless, he redescribed BROCCHI's species and described a series of new ones, providing good indications of localities and layers, whenever this was possible. Also in this respect he was among the most up to date paleontologists of his time.

Early and sudden death disrupted his studies. He left an unfinished but voluminous manuscript which was published posthumously by K. BEURLEN several years later. Although BEURLEN did this in an excellent manner, unfortunately he did not check the drawings in the manuscript. LÖRENTHEY himself was in the habit to scrutinise the figures before publication. Consequently all figures published during his life are extremely reliable to their finest details and to the outlines as well. Several new species are described in this posthumous work but the figures of the type specimens are incorrect and practically useless. For example, the figure of the right chela of *Calappa heberti* (LÖRENTHEY and BEURLEN 1929, pl. 6, fig. 11), borrowed from LÖRENTHEY's former publications is extremely reliable, while that of the left chela (fig. 12) is incorrect (pl. XXXVIII, figs. 4, 6 in this volume).

LÖRENTHEY's first papers were published in parallel Hungarian and German versions with identical contents. He presented the same data in preliminary publications without figures (LÖRENTHEY 1897 and 1898a), and in definite papers with excellent figures and detailed descriptions (LÖRENTHEY 1898b, c). The short definitions of new species, given in the first two papers satisfy articles 11 and 12 of the International Code of Zoological Nomenclature (ICZN, STOLL et al. 1961). Consequently 1897 is the date of first publication of the species *Cancer szontaghii*, *Calianassa rákosiensis* (= *Callianassa rakosiensis*), *C. brocchii*, *Pilodius mediterraneus* (= *Pilumnus r.*). He also mentions *Lambrus* sp. (= *Parthenope* sp.) and *Neptunus* cf. *granulatus* (= *Portunus monspeliensis*) and rede-

scribes BROCCHI's species. The name of *Portunus pygmeus* is erroneously spelled as *P. pygmaeus*, an error which remained uncorrected in the subsequent literature. *Andorina elegans* was described again in two parallel papers (LŐRENTHEY 1901a, b).

LŐRENTHEY (1911, 1913) described some Neogene localities from the environs of Budapest. In both the Hungarian and German versions, he mentions and names several new Badenian decapods without providing any description or figures. Thus the following names do not satisfy article 12 and 16 of the ICZN (STOLL et al. 1961) and should be regarded as nomina nuda: *Dorippe margaretha*, *Portumnus tricarinatus*, *Portunus rákosiensis* (= *Liocarcinus rakosensis*), *Zozymus mediterraneus* (= *Chlorodiella mediterranea*), *Microcorystes latifrons* (= *Corystites l.*).

The mentioned species were described later (LŐRENTHEY-BEURLEN 1929) together with some other Badenian species: *Maia biaensis* (= *Maja b.*), *Maia hungarica* (*Micippa h.*), *Calappa praelata* and *Calianassa pseudorákosiensis* (= *Callianassa pseudorakosensis*). As mentioned above the figures of the new species are incorrect. The text reveals that the descriptions were written by LŐRENTHEY and were left unaltered by BEURLEN (LŐRENTHEY-BEURLEN 1929, p. 8). Consequently, according to article 50 of ICZN (STOLL et al. 1961) the names should be attributed to LŐRENTHEY alone.

In Austria, TOULA (1904), in connection with the description of a new species, *Cancer bittneri* (= "Cancer" b.), enumerates all Badenian forms known. After the First World War, GLAESSNER began the study of Badenian decapods with the aim of completeness. In his first paper on this topic (GLAESSNER 1924) he did describe only a Carpathian ("Helvet") form, *Macrophthalmus vindobonensis*. In a subsequent publication (GLAESSNER 1928) he presents a monographic account of all known Neogene species of Austria and introduces the following Badenian forms: *Callianassa norica*, *Necro-nectes schafferi*, *Titanocarcinus vulgaris* (= *Pilodius v.*), *Carpilius antiquus*. In a volume of Fossilium Catalogus (GLAESSNER 1929) he summarized our knowledge of systematics of fossil decapods. Later he had to leave Europe thus the study of Badenian crabs was interrupted for two decades.

GLAESSNER's work was followed by BACHMAYER. In his paper on injured fossil gastropods (in PAPP et al. 1947), he analysed a rich collection of isolated fingers determining them to family or genus level. We have to remember here an amateur collector, K. OROSZY, whose collection provided material for the description of the extremely rich Deutsch-Altenburg fauna (BACHMAYER 1953a). In the same year a paper on the Gross-Höflein fauna appeared (BACHMAYER-TOLLMANN 1953a). The authors concentrated mainly on the description of a new species, *Dromilites koberi* (= *Kromtitis k.*), but mentioned the occurrence of many other forms including new ones. In other publications BACHMAYER dealt with sporadic findings (BACHMAYER 1950, 1953b, BACHMAYER-KÜPPER 1952) and with the calappids of the Vienna Basin (BACHMAYER 1962).

As shown above, most studies until the 1950-s were confined to Hungary and Austria. This tradition was broken, however, in the seventies. Förster (1979a, b) obtained a rich material, mostly isolated fingers but also some carapaces, from Poland. The most interesting localities were Grobie and Korytnica. YANAKEVICH (1969, 1977) described a new species, *Medaeus moldavicus* (= *Xantho m.*) and published figures of excellently preserved specimens of *Daira speciosa*. Both species came from reefal structures of Moldavia.

I reported mainly on Hungarian material (MÜLLER 1974a, b, 1976a, b, 1979a) but also on a fine collection obtained from E. KOJUMDGIEVA, Bulgaria (MÜLLER 1979b).

Other papers (e.g. BEURLEN 1939, SCHOUPPÉ 1949, STANCU-ANDREESCU 1968) contributed by mentioning known Badenian species from new localities.

Two more papers must be mentioned (HOJNOS 1923, 1933), which describe a small decapod fauna from an area lying between the northern parts of the Cserhát and the Mátra mountains, but without stating the exact locality. The author of them enumerates *Calianassa desmarestiana*, *C. subterranea*, *Gonoplax* sp., *Calappa heberti*, *Maia miocaenica* and describes a new variety, *Calianassa spinosa* LŐRENTHEY var. *cserhatica*. Unfortunately this material was lost and the determinations can not be checked. The short description of *C. spinosa* var. *cserhatica*, without figure, does not permit any further speculations.

The tradition of the study of Badenian decapods dates back for more than a century. In spite of this and in contrast to the presence of some excellent localities, the papers dealing with this topic are not numerous, and even the number of the authors is surprisingly low.

In order to revise previous studies I made attempts to locate the type specimens. LŐRENTHEY's material is preserved in the Hungarian Geological Survey in good order (Magyar Állami Földtani Intézet, Budapest XIV., Népstadion út 14. H 1142 Pf. 106). The specimens he collected were originally kept at the Palaeontological Institute of the Pázmány Péter University, Budapest (now Eötvös Loránd Tudományegyetem), but later his material was transferred to the Survey. The loss of types was moderate although universities are usually unsuited for preservation of such material. The following losses occurred: *Microcorystes latifrons* (= *Corystites l.*) the holotype was broken and lost while LŐRENTHEY was still alive (LŐRENTHEY-BEURLEN 1929, p. 138), *Portumnus tricarinatus*, holotype, and *Maia biaensis* (= *Maia b.*), holotype, but the plaster cast of which is extant, and *Callianassa rakosiensis* holotype.

In contrast, the fate of BROCCHI's (1883) specimens is unknown. The material described by him was collected by HEBERT and MUNIER-CHALMAS, paleontologists of the Sorbonne. Their collection, which includes a high number of specimens from Hungary, is still held at the Université Pierre et Marie Curie, Institut de Macropaléontologie, Paris 7. 2 Place Jussieu, France. With the kind help of professor PAJOT, I unsuccessfully tried to locate BROCCHI's specimens in this collections. The material was not found in the Muséum National d'Histoire Naturelle, Paris, either. The specimens are most likely lost.

All of GLAESNER's and BACHMAYER's material is preserved in excellent order in the Naturhistorisches Museum in Vienna (Burgring 7 A. 1014 Wien, Austria). Professor BACHMAYER kindly permitted me to study a good portion of them. The type-specimen of *Phymatocarcinus speciosus* (= *Daira speciosa*) however, was lost (GLAESNER 1928, p. 191). The fate of *Microdium nodulosum* is unknown. Many of BITTNER's types were also lost. In the inventory of the Geologischer Bundesanstalt, Wien (Rasumofskigasse 23, 1031 Wien, Austria) only *Ranidina rosaliae* is listed as extant (STOJASPALE 1975), though according to BITTNER (1884, p. 30) some specimens of *Cancer carniolicus* and the holotype of *C. styriacus* should also be kept there\*. The holotype of *Cancer illyricus* BITTNER is kept at the Institut für Geologie und Paläontologie of the Universität Graz (Heinrichstrasse 26, A-8010 Graz, Austria) as professor A. FENNINGER kindly communicated.

The specimens described by FÖRSTER (1979a, b) are kept in the collection of the Warsaw University (Universitet Warszawski, Wydział Geologii, Aleja Zwirki i Wigury 93. 02-089 Warszawa, Poland). YANAKEVICH' specimens are kept in State Teachers College, Tiraspol (Tiraspolski Gosudarstvenni Pedagogitsheski Institut im. Shevchenko, Tiraspol, Moldavian S.S.R., exact address unknown).

Specimens collected by KOJUMDGIEVA (MÜLLER 1979b) are deposited in the Museum of Sofia University (boulevard Ruski 15. Sofia 1000. Bulgaria).

The specimens I collected are owned and supervised by the Palaeontological and Geological Department of the Hungarian Natural History Museum (Természettudományi Múzeum, Föld- és Öslénytár, Múzeum körút 14-16. H-1088 Budapest, Hungary). They are deposited either in the Natural History Museum or are kept by the author (PÁL MÜLLER, Káplár u. 11. H-1024 Budapest, Hungary).

The author possesses rubber moulds of all type-specimens he was able to study and indicated as "material seen" in the descriptive chapter. He is prepared to supply casts for scientific study. The only exceptions are *Maia oroszyi* (*Pisa o.*) and *Necroneutes schafferi* the preservation of which did not allow to make a mould.

\* See „Addendum”

Most of the specimens were collected in open-air exposures. Efforts were made to visit the good outcrops repeatedly, ten to hundred times, to obtain a possibly complete record of the communities. This effort was helped by the circumstance that many of such localities are situated within Budapest or in its close vicinity. If it was possible, I removed bigger blocks from walls of exposures. The blocks were broken with hammer and chisel to pieces of an average size of four to five cm. All pieces were scrutinised with the help of head-band magnifying glasses. All recognisable fragments of decapods were retained, including moulds.

The generally poor preservation of the shells often precluded their cleaning up. In such cases we did not attempt to remove the rock covering the shell, but cleaned the external mould of the remnants of the shell and filled it with silicone rubber. After setting, the cast was pulled out cautiously. This procedure permitted the study of both inner and outer surfaces of the same specimen. To prevent sticking, porous rocks were impregnated with diluted varnish before applying the rubber. If shells were well preserved and hard, the common methods of cleaning were used as vibro-tool, needle, chisels, hippers, etc. Dark or shiny specimens were coated with vapour of ammonium chloride before photography. Silicone rubber is highly suitable for making casts or duplicates. Fine-grained plaster of Paris offers many advantages over plastics for making casts. After setting and drying, the casts were impregnated with diluted colourless varnish. To avoid bubble formation both rubber and plaster were applied with fine but strong brushes.

In some cases, e.g. for specimens with thin or narrow parts, epoxy resins might be preferred because of their strength. A resin of low viscosity was used. Its components were mixed with titanium oxide powder to achieve an opaque white cast. A highly suitable resin, fabricated by the Research Institute for Industry of Plastics (Műanyagipari Kutatóintézet, Hungária körút 114. H-1143 Budapest, Hungary) was used. Its components are designated FM-4 and T-30.

Casting with silicone rubber is described by KELLY-MCLACHLAN (1980). This paper defines the terms "cast" and "mould" which are frequently confused, and introduces the idea that casts can be enlarged treating with toluene. This method was independently developed by us. A linear increase of 50% without any distortion can be obtained after a bath in toluene for several hours. Repeating this procedure a linear magnification of 5 to 8 can be obtained.

In some cases the rock was sufficiently friable and the shells well preserved to permit sieving. The use of magnifying glasses while sorting the sieved material assures recognition of fossils of all sizes. In comparing such samples with hand-picked ones, a bias-factor was used to correct the number or proportion of small specimens.

## DESCRIPTION OF THE LOCALITIES

I describe here those localities where I collected (Table 1.). Sections are given if outcrops were surveyed by us. In other cases I refer to literature. Localities yielding only one or few species are presented in the systematic part. The codes given for the individual layers of the sections are those used in our inventory system in which the first group of letters and numbers refers to the layer itself, regarded as a unit, the second group refers to the serial number of the given species, and the third one is the serial number of individuals.

Outcrops in Budapest or in his closest vicinity were studied most intensively. Most of these localities are Upper Badenian. In subsurface position Middle Badenian rocks occur also but these yielded no decapods so far (Fig. 1).

Tétény-plateau (Tétényi-fennsík, Fig. 2). This upland of moderate height is built up of Egerian and Neogene rocks. The Upper Badenian rocks are exposed in small patches and as a narrow strip near to the northern edge of the height. The total thickness of Badenian layers seldom exceeds six meters.

The roadcut of the Kamaraerdei út (MK on Fig. 2) is still accessible. In earlier literature it was known as Katona út or Militärstrasse (LÖRENTHEY 1911, 1913, LÖRENTHEY-BEURLEN 1929, STRAUSZ 1923). Stratum MK is exposed as a 1 meter layer (Fig. 3) containing moulds and recrystallized remnants of corals. At the southern end of the exposure, *Tarbellastraea* sp., *Porites* sp. and *Siderastraea* sp. colonies were found in removed position. A species similar in shape to *Tarbellastraea* but probably belonging to another genus was also found. About 10 meters apart, in the northern part of the cut a small *Porites*-reef occurred, mainly in living position. The matrix is a conglomerate of ill-sorted grains and blocks. The layer was mentioned by us (MÜLLER 1976a), in which paper KÓKAY (p. 507) enumerated and interpreted the Molluscs. KÓKAY argued for a well areated, agitated water of a depth much less than 30 meters. The salinity was near to the oceanic one. Decapods of the layer MK: *Dardanus hungaricus* (LÖRENTHEY), *Petrolisthes magnus* n. sp., *Charybdis (Goniosupradens)* mathiasi n. sp., *Xantho moldavicus* (YANAKEVICH), *Chlorodiella mediterranea* (LÖRENTHEY) and *Daira speciosa* (REUSS).

The coral-bearing layer is covered by a calcarenite with abundant remnants of *Cerithium* (layer MKF), by a faint unconformity. Macroscopically, this 1.5 meter layer is a microoncoidal-skeletal grainstone. Decapods: *Pisidia kokayi* (MÜLLER), *Pisa oroszyi* (BACHMAYER), *Xantho moldavicus* (YANAKEVICH), *Pilumnopeus tetensis* n. sp. and *Brachynotus februario* MÜLLER.

Roadcut of the streets Balatoni út and Szabadkai út (MF on Fig. 2). The exposure is accessible. A medium to coarse-grained dark grey sand (Eggenburgian) is unconformably overlain by Upper Badenian layers. The lowermost one is an ill-sorted gravel with blocks up to 20 cm with some bivalve boreholes. A sandy limestone (MFK) (Fig. 3) covers the preceeding layer which is an ill-sorted quartz-sand and gravel-bearing packstone and contains abundant moulds of a *Glycymeris* species and other molluscs. Its thickness is about 0.8 meter. Its fauna was studied by KÓKAY (quoted by MÜLLER 1976a, p. 506; for *Stylophora* sp. read *Tarbellastraea* sp.).

The following layer is an 1.6 m thick limestone comprising a lower, softer and an upper, harder part, both containing quartz grains. These layers were not studied in details, nevertheless samples were collected from the debris at the foot of the cut, containing blocks of all layers. This mixed material got the code (MFA). The uppermost layer of the cut was studied at another spot be described below (MM and MB). The sequence is illustrated on Fig. 3. A quartz-bearing sandstone appeared at the base of a pit, situated westward of the cut (MA). KÓKAY's quoted description fits to the units MA and MFA as well. He argued for a depository environment with an agitated water of a salinity similar to that of the oceans. The water-depth did not exceed 30 m but probably it was shallower. The size of quartz grains of these layers decrease toward south suggesting the presence of a shore north of the site built up of sands and gravels (probably coming from Carpathian layers) and a bottom gently sloping southwards.

## Index of Badenian decapod localities

Country	Locality Layer, Site	Zone	Page	Country	Locality Layer, Site	Zone	Page
Hungary	Budapest, Tétény-plateau			Hungary	Zebegény, quarry		
	MK	3	36		MZZ	?1	44
	MKF	4	36		Szob, Dámasdi patak		
	MF	3	36		MSZ	?1	44
	MFA	3	36		Mátraverebély-Szentkút		
	MFK	3	36		Szent László-forrás	1	44
	MA	3	36		Meszes-tető	1	44
	MB	4	39		Sámsonháza-Halastó-hegy		
	MM	4	39		“Bázishomok”	1	44
	MDG	4	39		Sámsonháza, Kis-Zagyva		
	MJL	4	39		völgye		
	MBH	4	39		“Csúcs-hegy alja”	1	44
	MUO	4	39		Nyirád		
	Diósd				MNU	1	44
	MDZ	3	39		Balatonakali, drilling		
	MDN	?4	40		BAK-40		
	Bia				BAK	3+4	45
	MNA	3	40		Kishajmás		
	MN	3	40		railway-cut	?1	45
	Budapest, Rákos			Austria	Gross-Höflein, Fenkquarry		
	MR	3	40		MNH	2	45
	MRZ	3	40		Deutsch-Altenburg, quarry		
	MR45	3	40		DAL	?4	47
	MR8	3	40		Oslip, sand-pit		
	MRL	3	41		MOS	?2	47
	MRO	3	41	Poland	Grobie		
	MRF	3	41		MGR	?1	47
	MRC	4	42		Węglinek		
	MRS	4	42		WEG	4	47
	Budapest, Kerepesi út			USSR (Moldavian SSR)	Koban, Shepteban, Varatik, Gorodishtshe, Bolotino		
	MKC	3	42			3	47
	Budapest, Gyakorló út			Bulgaria	Iasen	?1	47
	MGY	3	42		Gorna Mitropolia	?1	47
	MGF	4	43		Krivodol	?1	47
	Budapest, Örs vezér tere				Oriakhovo	?3	47
	MOH	3	42	Romania	Transsylvania		
	MOE	4	43		Tasádfő	?	47
	Budapest, Mező I. út drilling H-34			Jugoslavia	Slovenia		
	MH34	3	43		Zagor	?	47
	Budapest, Telepy utca						
	MTE	?3	43				
	Visegrád						
	MV	1	44				
	MVA	1	44				
	Nagymaros-Törökmező						
	MTH	1	44				
	MTZ	1	44				
	MTM	1	44				




Fig. 1. Localities in and around Budapest. Areas indicated with letters A and B are shown on Figs. 2, 4




Fig. 2. Sites on Tétény-plateau with the codes of decapod-bearing layers




Fig. 3. Simplified section of sites on the Tétény-plateau (horizontally not to scale)  
1. Codes of decapod-bearing layers, 2. boundary of biozones, 3. unconformity

Almost the same species were found in layers MA, MF and MFK: *Callianassa munieri* BROCCHEI, *C. pseudorakosensis* LÖRENTHEY, *C. kerepeziensis* MÜLLER, *Dardanus substriatiformis* (LÖRENTHEY), *Petrochirus priscus* (BROCCHEI), *Dromia eotvoesi* (MÜLLER), *Calappa heberti* BROCCHEI, *Matuta brocchii* GLAESNER, *Maja biaensis* LÖRENTHEY, *Xantho moldavicus* (YANAKEVICH), *Pilumnus mediterraneus* (LÖRENTHEY) and *Micippa hungarica* (LÖRENTHEY).

The layer covering the quartz-bearing limestones is pure calcarenite containing oncoids and stromatolites and it is almost totally depleted of quartz-grains (MB and MM). The small pits where these rocks were exposed are mostly filled now, but collecting is still possible on the surface or in the road-cut. The oncoids and stromatolites are of cryptalgal and foraminiferal origin (LELKES-MÜLLER in press). The matrix is microoncoidal grainstone which contains molluscs, predominantly Cerithium. Its thickness decreases toward WSW, with the sessile stromatolitic structures giving place to sub-globular oncoids which occur in decreasing number toward WSW. All this suggest a paleoenvironment with a shallow nearshore site where the bottom gently sloped toward south similarly to the presumed situation when the underlying layers were deposited. The molluscs were determined by KÓKAY (quoted by MÜLLER 1974a, p. 119) suggesting an environment of a shallow sublittoral sea of about 26 to 28‰ salinity. The abundance of Cerithium indicates a dense cover of algae or seagrasses.

The diversity of the decapod fauna increases toward south. At the northern part of the section (MM) four species were found: *Pisa oroszyi* (BACHMAYER), *Xantho moldavicus* (YANAKEVICH), *Pilumnus mediterraneus* (LÖRENTHEY) and *Brachynotus februarius* MÜLLER. Further south at MB, MUO, MJL and MBH the following species occurred: *Callianassa* sp., *Pisidia kokayi* (MÜLLER), *Pagurus albus* MÜLLER, *Anapagurus marginatus* MÜLLER, *Galathea weinfurteri* BACHMAYER, *Matuta brocchii* GLAESNER, 'Ebalia' hungarica MÜLLER, *Parthenope tetenyensis* n. sp., *Pisa oroszyi* (BACHMAYER), *Micippa hungarica* (LÖRENTHEY), *Liocarcinus kuehni* (BACHMAYER), *L. rakosensis* (LÖRENTHEY), *Miopipus pygmeus* (BROCCHEI), ?*Thalamita* sp., *Xantho moldavicus* (YANAKEVICH), *Pilumnus mediterraneus* (LÖRENTHEY), 'Pilumnopeus' tetenyensis n. sp., *Brachynotus februarius* MÜLLER and *Pachygrapsus hungaricus* MÜLLER.

At the southernmost part of the section the discussed layer is partly covered, partly replaced by a fine-grained ill-sorted calcarenite, probably a packstone, which contains no macrooncoids (MDG). The non-decapod fauna was determined by KÓKAY (quoted by MÜLLER 1974b, p. 275). According to him the depth of the water was shallower than 30 to 40 meter, the salinity was, at least temporarily, slightly instable, but close to the oceanic one. The rock is exposed in some still accessible shallow trenches. Decapods: *Callianassa* sp. (?= *Semiranina oroszyi* BACHMAYER), *Upogebia* sp., *Galathea weinfurteri* BACHMAYER, *Anapagurus marginatus* MÜLLER, *Paguridaram* div. sp., *Kerepesia viai* MÜLLER, *Mursia lienharti* (BACHMAYER), *Dorippe margaretha* LÖRENTHEY, 'Ebalia' hungarica MÜLLER, *Trachypirimela grippi* (MÜLLER), *Miopipus pygmeus* (BROCCHEI), *Liocarcinus rakosensis* (LÖRENTHEY), 'Lissocarcinus' szoeraenyiae (MÜLLER), *Parthenope szaboi* MÜLLER, *P. tetenyensis* n. sp., *Micippa hungarica* (LÖRENTHEY), *Pisa oroszyi* (BACHMAYER), *Xantho moldavicus* (YANAKEVICH), *Actumnus telegdii* (MÜLLER) and *Pilumnus mediterraneus* (LÖRENTHEY).

Localities at Diósd (Fig. 2). In the southern parts of the Tétény-plateau there is a small outcrop of Badenian rocks. These are exposed in an abandoned quarry and at the edge of a sandpit, along a fault, which tectonically connects Pannonian sands to Badenian and Sarmatian limestones. Site MDZ is situated along this fault on the upper part of the sand-pit that is still worked making the locality accessible. A compact hard limestone contains here abundant moulds of *Tarbellastraea* sp., *Porites* sp. and two other species of reefbuilding corals, mostly in removed position. The matrix is generally ill-sorted and abundant in crab-remnants. The locality was not yet mentioned in the literature. Decapods found: *Pisidia kokayi* (MÜLLER), *Petrolisthes magnus* n. sp., *Galathea weinfurteri* BACH-

MAYER, *Dromia neogenica* (MÜLLER), *Calappa heberti* BROCCHEI, *Rakosia carupoides* n. gen. n. sp., *Charybdis mathiasi* n. sp., 'Necronectes' schafferi GLAESSNER, *Micippa hungarica* (LÖRENTHEY), *Panopeus wronai* n. sp., *Xantho moldavicus* (YANAKEVICH), *Pilumnus mediterraneus* (LÖRENTHEY), *Chlorodiella mediterranea* (LÖRENTHEY), *Ch. juglans* n. sp., ?*Glabropilumnus* sp., *Crossotonotus diosdensis* n. sp. and *Cancer styriacus* BITTNER.

Elsewhere in the just mentioned quarry (Fig. 2) some blocks contained a fauna (MND) which is probably younger than that of MDZ but the contact of the two layers is covered. Decapods: *Callianassa* sp., *Paguridaram* sp., *Pagurus albus* MÜLLER, *Pagurus rakosensis* MÜLLER, *Mursia lienharti* (BACHMAYER), *Dorippe margaretha* LÖRENTHEY, *Parthenope szaboi* MÜLLER, 'Lissocarcinus' szaeraenii (MÜLLER), *Liocarcinus rakosensis* (LÖRENTHEY), *Miopipus pygmeus* (BROCCHEI), *Actumnus teledegii* (MÜLLER).

**Bia.** At the western edge of an upland connected to Tétény-plateau, Badenian and Sarmatian strata are exposed along a valley-side. The name of the site is "Nyakaskő", and it was described among others by STRAUSZ (1924) and by SCHAFARZIK-VENDL (1929). At the lower part of the section there is a silt-layer (MNA) (Table 1.). The silt contains deformed, abundant moulds of *Turritella* sp. and other molluscs. Decapods found: *Callianassa munieri* BROCCHEI, *Mursia lienharti* (BACHMAYER), *Maja biaensis* LÖRENTHEY, *Portunus* cf. *monspeliensis* (A. MILNE-EDWARDS). The silty layer is covered by four to five meter thick sandy limestone. This contains a rich mollusc fauna, mostly in the form of moulds (MN) (Table 1.). Decapods: *Callianassa munieri* BROCCHEI, *C. pseudorakosensis* LÖRENTHEY, *Dromia eotvoesi* (MÜLLER), *Calappa heberti* BROCCHEI, *Matuta brocchii* GLAESSNER, *Mursia lienharti* (BACHMAYER), *Maja biaensis* LÖRENTHEY, *Xantho moldavicus* (YANAKEVICH), *Pilumnus mediterraneus* (LÖRENTHEY).

A group of sites (Figs. 1, 4, 5), which yielded the most diverse and best preserved known crab fauna of the Paratethys, is situated in the eastern part of Budapest, in and around the classic site Rákos. The sequences in this area were described in several publications and was recently summarized by KÓKAY-MIHÁLY-MÜLLER (in press.).

**Rákos.** This exposure is a railway-cut excavated in the second half of the last century and subsequently deepened and widened several times. The still accessible exposure consists of several layers of Badenian and Sarmatian age (see section in Fig. 5). The section was described by SZABÓ (1879), FRANZENAU (1881), VADÁSZ (1906) and by SCHAFARZIK and VENDL (1929). BROCCHEI's (1883) fauna came from here. Layers below MRC (see section, Fig. 5) contain tuffaceous minerals which are partly reworked from Upper Carpathian dacitic tuffs erstwhile covering slopes and hills eastwards. Some eruptions might have occurred during the deposition of layer MRF, since the potassium-argon age of this layer is much younger ( $13.4 \pm 0.6$  MYBP) than that of a slightly lower one (layered tuffite below MR8, age:  $15.6 \pm 0.8$  MYBP, BALOGH-ÁRVA-SÓS-PÉCSKAY 1980). Most likely the biotite of the lower layer originated exclusively from older tuffs, while that of MRF got crystals from a new eruption as well.

A *Porites*-reef (MRZ) in the eastern part of the cut forms a 1.5–2.0 meter thick lens. It was a patch-reef built mainly by *Porites* sp. which was preserved as moulds, mostly in living position. Few specimens of *Tarbellastraea* was also found. The 5–10 cm thick and 0.5–1.0 m tall coral branches are embedded in a tuffaceous limestone. Some of the corals are encrusted by calcareous red algae, probably by *Neogoniolithon* sp. Occasionally bryozoans participated in the building. The *Porites* was earlier erroneously determined as *Millepora* (LÖRENTHEY-BEURLEN 1929, p. 35) but in fact no *Hydrozoa* remnants were found so far.

A more or less monospecific coral reef indicates a muddy, shallow environment. *Porites* is one of the corals resistant to fast sedimentation and to muddy water. The matrix, rich in fine particles, argues for a muddy water, too. Decapods (MRZ): *Callianassa munieri* BROCCHEI, *Pisidia kokayi* (MÜLLER), *Petrolisthes magnus* n. sp., *Galathea weinfurteri* BACHMAYER, *Dromia neogenica* MÜLLER, *Calappa heberti* BROCCHEI, *Matuta brocchii* GLAESSNER, *Pisa oroszyi* (BACHMAYER), *Rakosia carupoides* n. gen. n. sp., *Charybdis mathiasi* n. sp., *Thalamita* n? sp., *Daira speciosa* (REUSS), *Chlorodiella mediterranea* (LÖRENTHEY), *Ch. mediterranea tetenyensis* n. ssp., *Ch. loczyi* n. sp., *Pilumnopeus paratethyensis* n. sp., ?*Glabropilumnus* sp., *Pilumnus mediterraneus* (LÖRENTHEY), *Xantho moldavicus* (YANAKEVICH), ?'Necronectes' cf. *schafferi* GLAESSNER and *Pachygrapsus hungaricus* MÜLLER.

Sandstone layers MR45 and MR8. These sandstones partly replace, partly cover the just described reef. These rocks are medium-grained, strongly bioturbated, ill-layered and nodular. Most decapods are well preserved but disarticulated. In contrast, aragonite-shelled molluscs are rare, probably their moulds collapsed before the diagenesis of rock. Decapods: *Callianassa munieri* BROCCHEI, *C. chalmasii* BROCCHEI, *C. pseudorakosensis* LÖRENTHEY, *C. rakosiensis* LÖRENTHEY, *C. kerepesiensis* MÜLLER, *Petrochirus priscus* (BROCCHEI), *Pagurus rakosensis* MÜLLER, 'Anapagurus' miocenicus MÜLLER, *Upogebia* sp., *Albunea asymmetrica* (MÜLLER), *Dromia eotvoesi* (MÜLLER), *Calappa heberti* BROCCHEI, *Mursia lienharti* (BACHMAYER), 'Ebalia' hungarica MÜLLER, *Parthenope szaboi* MÜLLER, *P. tetenyensis* n. sp., *Thalamita fragilis* MÜLLER and *Pilumnus mediterraneus* (LÖRENTHEY).




Fig. 4. Localities near Rákos

MR8 limestone is overlain by a sequence of sandy limestone layers. The lower layer is characterized by the abundance of *Lingula columbella* (MRL), the upper one (MRO) contains oyster beds. The proportion of quartz grains and tuffigenous particles decreases upwards. The rock of the upper bed is a sandy foraminiferal-molluscan biosparite (LELKES pers. comm.). Decapods of the both layers: '*Callianassa*' *pseudorakosensis* LÖRENTHEY, '*C.*' *chalmasicii* BROCCHI, '*C.*' sp., *Upogebia* sp., *Dromia eotvoesi* (MÜLLER), *Calappa heberti* BROCCHI, *Matuta brocchii* GLAESSNER, '*Ebalia*' *hungarica* MÜLLER, *Petrochirus priscus* (BROCCHI), *Thalamita fragilis* MÜLLER, *Portunus (Monomia)* *miocaenicus* n. sp., and *Pilumnus mediterraneus* (LÖRENTHEY).

The thickest layer in the railway-cut is a fine-grained ill-sorted sandstone (MRF). Most of the grains are volcanic. The same layer is exposed also at Keresztúri út (MEG, Fig. 3). Big pectinids and irregular sea urchins are abundant. The ill-layered sandstone is characterized by dense Callianassid burrows (*Ophiomorpha*). Decapods: '*Callianassa*' *munieri* BROCCHI, '*C.*' *chalmasicii* BROCCHI, *Petrochirus priscus* (BROCCHI), *Dardanus* sp., *Pagurus concavus* MÜLLER, *P. rakosensis* MÜLLER, '*Anapagurus*' *miocenicus* MÜLLER, *Diogenes* cf. *pugilator* ROUX, *Albunea asymmetrica* (MÜLLER), *Dorippe margaretha* LÖRENTHEY, *Calappa heberti* BROCCHI, *Matuta brocchii* GLAESSNER, *Dromia eotvoesi* (MÜLLER), *Parthenope tetenyensis* n. sp., *P. szaboi* MÜLLER, *Mursia lienharti* (BACHMAYER), *Pisa* cf. *oroszyi* (BACHMAYER), *Liocarcinus rakosensis* (LÖRENTHEY), *Portunus monspeliensis* (A. MILNE-EDWARDS), *Miopipus pygmeus* (BROCCHI), *Pilumnus mediterraneus* (LÖRENTHEY) and *Actumnus telegdii* (MÜLLER).




Fig. 5. Simplified section along the localities near Rákos (horizontally not to scale)  
1. Code of decapod-bearing layers, 2. boundary of biozones, 3. unconformity

This sandstone is covered by a porous mollusc-bearing limestone virtually depleted of terrigenous material (MRC). Its most characteristic molluscs are *Cardium* and *Cerithium* species. Decapods: *Callianassa* cf. *kerepesiensis* MÜLLER, *Dromiidarum* sp. (?*Kerepesia*), *Matuta brocchii* GLAESSNER, *Pilumnus mediterraneus* LÖRENTHEY and *Xantho moldavicus* (YANAKEVICH).

The limestone is covered by a 0.1–0.2 m thick macrooncoidal-bearing layer (MRS); the oncoids are embedded in microoncoidal-oolitic grainstone matrix, containing decapods (LELKES-MÜLLER in press.). Decapods: *Parthenope tetenyensis* n. sp. 'Pilumnopeus' *tetenensis* n. sp., *Pachygrapsus hungaricus* MÜLLER and *Brachynotus febrarius* MÜLLER. The oncoidal layer is overlain by a crossbedded calcarenite which did not yield decapods yet.

Another group of localities is situated at a distance of about 1 km from Rákos, around the square Örs vezér tere (Fig. 3). Numerous trenches and pits were excavated here in the last twenty years. The sites are described by STRAUSZ (1927), BARTKÓ-KÓKAY (1966), MIHÁLY (1969), MÜLLER (1976b, 1979a) and by KÓKAY-MIHÁLY-MÜLLER (in press.). The most interesting exposures were in trenches along the Kerepesi út, Gyakorló út and in the construction-pit of the "Sugár" supermarket (Fig. 3). The lower layers were almost identical in all exposures, while the upper ones differed from place to place.

The lowest crab-bearing layer was a tuffaceous limestone which contained corals. Its fauna did not differ from that of MRZ in Rákos. A tuffaceous sandy limestone occurred at the corner of the streets Kerepesi út and Sarkantyú út (now Gyakorló köz). This 2 m limestone (MKC) corresponds to layers MRC and MRO of Rákos but is deposited from a more agitated water than it was the case at Rákos (BARTKÓ-KÓKAY 1966). Decapods: *Callianassa munieri* BROCHI, *C. kerepesiensis* MÜLLER, *Pagurinae* ?n. sp., *Petrochirus priscus* (BROCHI), *Dardanus substriatiformis* (LÖRENTHEY), *Galathea weinfurteri* BACHMAYER, *Dromia neogenica* MÜLLER, *D. eotvoesi* (MÜLLER), *Kerepesia viai* MÜLLER, *Calappa heberti* BROCHI, *Matuta brocchii* GLAESSNER, *Mursia lienharti* (BACHMAYER), 'Ebalia' *hungarica* MÜLLER, *Parthenope szaboi* MÜLLER, *P. tetenyensis* n. sp., *Thalamita fragilis* MÜLLER, *Portunus* sp., *Liocarcinus rakosensis* (LÖRENTHEY), *Miopipus pygmeus* (BROCHI), *Xantho moldavicus* (YANAKEVICH), *Pilumnus mediterraneus* (LÖRENTHEY), *Chlorodiella* sp., *Pisa oroszyi* (BACHMAYER), *Maja biaensis* LÖRENTHEY and *Micippa hungarica* (LÖRENTHEY).

The cover of the limestone, a four to five meter thick medium-grained sand, was exposed along the street Gyakorló út (MGY) and in the construction-pit of "Sugár" supermarket (MOH, Fig. 5). It had the appearance of quartz-sand, though its quartz-content was only 5%, the clear particles were mostly glasses. The CaCO<sub>3</sub> content was about 40%. Other crystalline minerals were mostly feldspars, mica and clay-minerals. The sand contained big *Pectinids* (*Flabellipecten leythaianus*, *Pecten aduncus*), sea urchins and foraminifers (*Borelis melo* dominated). The crabs were usually well preserved and only moderately fragile, thus they could be collected from sieved material. Decapods found: *Upogebia* sp., *Callianassa kerepesiensis* MÜLLER, *C. chalmensis* BROCHI, *C. rakosiensis* LÖRENTHEY, *C. pseudorakosiensis* LÖRENTHEY, *C. brocchii* LÖRENTHEY, *Dardanus substriatiformis* (LÖRENTHEY), *Diogenes* cf. *pugilator* ROUX, *Galathea weinfurteri* BACHMAYER, *Dromia eotvoesi* (MÜLLER), *Calappa heberti* BROCHI, *Matuta brocchii* GLAESSNER, *Mursia lienharti* (BACHMAYER), *Maja biaensis* LÖREN-

THEY, *Portunus neogenicus* MÜLLER, *Liocarcinus rakosensis* (LÖRENTHEY), *Miopipus pygmeus* (BROCCHI), *Mioxaiva psammophila* MÜLLER, *Xantho moldavicus* (YANAKEVICH), *Pilumnus mediterraneus* (LÖRENTHEY), *Actumnus telegidii* (MÜLLER) and ?*Ocypode* sp.

This tuffaceous sand is covered by a limestone-complex composed of different rocks at different places. At Gyakorló út (Fig. 5) the sand is covered by *Ervilia*-bearing limestone. Its thickness was about 0.4 m. It was covered by a 0.5 m thick *Cerithium*-limestone (MGF). This contained big oncoids with calcarenite-pebble or black pebble nuclei (LELKES-MÜLLER in press.). Decapods found: *Callianassa* sp., *Pisa oroszyi* (BACHMAYER), *Xantho moldavicus* (YANAKEVICH), *Pilumnus mediterraneus* (LÖRENTHEY), *Pachygrapsus hungaricus* MÜLLER.

At Örs vezér tere in the construction-pit of "Sugár" supermarket sand MOH is covered by a tuffaceous layer of moderate thickness (0.5 m). A calcarenite (MOE), which is probably contemporaneous with layer MGF at Gyakorló út, covers the tuffite. The thickness of this limestone is about 2.2 meters. The rock is skeletal grainstone with sparse microoncoids, characterised by small oysters, *Chlamys*, regular and spatangid sea urchins and *Lingula* sp. Due to extremely favourable conditions for preservation, its decapod fauna is the richest known from the Paratethys: *Callianassa kerepesiensis* MÜLLER, *C. sp.*, ?*Upogebia* sp., *Pagurus rakosensis* MÜLLER, *P. concavus* MÜLLER, *P. albus* MÜLLER, 'Anapagurus' *marginatus* MÜLLER, *Dardanus substriatiformis* (LÖRENTHEY), *Galathea weinfurteri* BACHMAYER, *Kerepesia viai* MÜLLER, *Matuta brocchii* GLAESSNER, *Mursia lienharti* (BACHMAYER), *Dorippe margaretha* LÖRENTHEY, *Palicus* n. sp., *Cancer styriacus* BITTNER, *Trachypirimela grippi* (MÜLLER), *Parthenope szaboi* MÜLLER, *P. tetenyensis* n. sp., *Maja biaensis* LÖRENTHEY, *Micippa hungarica* (LÖRENTHEY), *Pisa oroszyi* (BACHMAYER), *Achaeus magnus* MÜLLER, 'Lissocarcinus' *szoeraenyiae* (MÜLLER), 'Xaiva' *bachmayeri* n. sp., *Liocarcinus rakosensis* (LÖRENTHEY), *Miopipus pygmeus* (BROCCHI), *Ebalia oersi* MÜLLER, *Palaeomyra globulosa* (MÜLLER), *Xantho moldavicus* (YANAKEVICH), *Pilumnus mediterraneus* (LÖRENTHEY), *Actumnus telegidii* (MÜLLER).

Drilling H-34 in Mező Imre út, Budapest (MH34, Fig. 1). Below a layer of quaternary sand and gravel an alternating sequence of fine-grained silty sandstone and calcarenite is present (KÓKAY pers. comm.), which is underlain by a variegated clay-silt deposit of Middle Badenian. In 25.8–33.0 meter depth, the following decapods were found in Upper Badenian sandstones and limestones: *Callianassa* sp., *Calappa heberti* BROCCHI, *Liocarcinus rakosensis* (LÖRENTHEY), *Pilumnus mediterraneus* (LÖRENTHEY) and *Macrophthalmus vindobonensis* GLAESSNER.

At Telepy utca (MTE, Fig. 1) a trench was excavated at the beginning of the century. It exposed Badenian limestones, described by SCHAFARZIK (1903). The site yielded one decapod species: *Andorina elegans* LÖRENTHEY.

Visegrád (Fig. 6). There are poorly exposed layers on a hill-side (Fekete-hegy). The sequence was described by SCHAFARZIK—VENDL (1929), SCHOLZ (1970) and BÁLDI—KÓKAY (1970). Probably this is the type-locality of numerous hermatypic corals described by REUSS (1872) as Nagymaros specimens. Tuffaceous layers with blocks of andesite are covered by Lower Badenian marly limestones




Fig. 6. Localities in the Börzsöny and Pilis Mountains

which contain corals in the lower layers, and calcareous red algae in the upper ones. Fossils can be collected from blocks scattered on the surface. The lack of good exposures precludes exact surveying of the section. Moderately preserved crabs were found in the coral-bearing layer (MV) together with *Tarbellastraea* sp., *Porites* sp., *Stylophora subreticulata* REUSS etc: *Pisidia* aff. *kokayi* MÜLLER, *Galathea weinfurteri* BACHMAYER, *Kromtis koberi* (BACHMAYER), *Dynomene emiliae* MÜLLER, *Schizophrys visogradensis* n. sp., *Chlorodiella mediterranea tetenyensis* n. ssp., *Xantho* cf. *moldavicus* (YANAKEVICH), *Daira speciosa* (REUSS), *Trapezia glaessneri* MÜLLER, *Haydnella steiningeri* n. sp., *Portunus* cf. *monspeliensis* (A. MILNE-EDWARDS), *Carupa* cf. *tenuipes* DANA, *Charybdis mathiasi* n. sp. and *Carpilius antiquus* GLAESNER.

Some meters farther uphill a calcareous red algae bearing limestone (MVA) yielded the following decapods: *Paguridaram* sp., *Calappa praelata* LÖRENTHEY.

Törökmező (Fig. 6). A group of new exposures, not described yet, are situated near the hotel "Törökmező". The cut of a new road exposes a marly tuffaceous limestone containing the giant foraminifer *Heterostegina gigantaeformis* PAPP which is characteristic for Lower Badenian deposits (MTH). Decapods found: *Dardanus arrosor* (HERBST) and *Calappa praelata* LÖRENTHEY. 500 meter north a small quarry operated in the past. The visible thickness of the only exposed layer is only 0.5 meter (MTZ). This limestone contains abundant moulds of a diverse coral-fauna. The specimens were embedded in removed position. The fine-grained micritic limestone contains poorly preserved decapods: *Pagurus turcus* n. sp., 'Pylopagurus' sp., *Pisidia* aff. *kokayi* MÜLLER, *Dynomene emiliae* MÜLLER, *Dromia neogenica* MÜLLER, *Dromiidarum* ?n. sp., *Schizophrys visogradensis* n. sp., *Charybdis mathiasi* n. sp., *Trapezia* cf. *glaessneri* MÜLLER, *Daira speciosa* (REUSS), *Actaea turcocampestris* n. sp., *Panopeus wronai* n. sp., *Carpilius antiquus* GLAESNER, *Chlorodiella mediterranea* (LÖRENTHEY), *Ch. mediterranea tetenyensis* n. ssp., *Xantho moldavicus* (YANAKEVICH), *Pilumnus mediterraneus* (LÖRENTHEY), 'Glabropilumnus' ?n. sp., *Haydnella steiningeri* n. gen. n. sp.

A small pit further north is excavated in a marl-layer which contains molluscs (MTM). The visible thickness of the only layer is about 3 meter. The relation of this marl to the previously mentioned limestones is unknown because the exposures are small and separated. All layers belong to the Lower Badenian substage at Törökmező. This age is supported by studies on calcareous mannoflora (pers. comm. by NAGYMAROSI). The molluscs have not been studied yet. Decapods: *Ebalia* ?n. sp., ?*Calappa* sp.

Quarry at Zebegény (MZZ, Fig. 6). A huge quarry is under operation in the Southern Börzsöny Mountains, named as "Bakókút" or as "Bószob". A soft calcareous red algae bearing limestone is intercalated by a 4 m thick hard layer. This hard micritic calcarenite is full of moulds of a small, branching ahermatypic coral, resembling *Oculina*. Few moulds of *Porites* sp. were also found. Decapods: *Galathea* cf. *weinfurteri* BACHMAYER, *Petrolisthes haydni* n. sp., *Callianassa* cf. *munieri* BROCCHI, *Chlorodiella mediterranea* (LÖRENTHEY), *Xantho* cf. *moldavicus* (YANAKEVICH), 'Polidius' *vulgaris* (GLAESNER), *Actumnus* ?n. sp., *Ebalia* n. sp. 1 and n. sp. 2.

Szob (MSZ), a site well-known of its molluscs, yielded only few chelae and fingers. The section is situated above a small creek. The lowest layer is a grey and yellow sandstone (about 6 m thick) with sporadic molluscs. It is covered by a 0.1 m thick lumachella-like sand and by a 0.5 m thick brown tuffaceous silt. All layers yielded some decapods: 'Callianassa' *szobensis* n. sp., *C. cf. chalmasic* BROCCHI, *Calappa praelata* LÖRENTHEY, *Cancer* sp.

A group of Badenian outcrops, situated at the northwestern ends of Mátra Mountains, yielded mostly chelae and fingers. The area is described by NOSZKY (1940, pp. 102–113, 203–205) and by HÁMOR (in press).

Mátraverebély-Szentkút (Szent László-forrás, Fig. 7). The crab-bearing layers are sandstones. They cover a volcanic sequence. Decapods: *Callianassa pseudorakosensis* LÖRENTHEY, *C. kerepesiensis* MÜLLER, *Diogenes matrensis* n. sp., *Dardanus hungaricus* (LÖRENTHEY), 'Pylopagurus' cf. *leganyii* n. sp., ?*Paguristes* sp., *Calappa praelata* LÖRENTHEY, *Maja* cf. *biaensis* LÖRENTHEY, *Parthenope* sp., *Portunidaram* ind. sp., *Liocarcinus rakosensis* (LÖRENTHEY) and *Pilumnus mediterraneus* (LÖRENTHEY).

Two sites are situated near Sámsonháza (Fig. 7). The exact location of one of these, named on labels as "Halastóhegy-bázishomok" is unknown. The presumed position is shown on Fig. 7. Decapods collected by HARMAT: *Callianassa pseudorakosensis* LÖRENTHEY, *Paguristes cserhatensis* n. sp.

The other site is a small quarry in the valley of the creek Kis-Zagyva. This might be identical with HOJNOS' locality (1923, 1933). Decapods: 'Pylopagurus' *leganyii* n. sp., *Pilumnus* cf. *mediterraneus* LÖRENTHEY.

Nyirág (MNU Fig. 8). A roadcut is situated 500 meter west of the cemetery of the village Nyirág (Bakony Mountains) along the highway leading to Sümeg. A single 2.5 m layer of limestone is exposed, which is a coarse-grained calcarenite containing abundant moulds of big molluscs and some corals. The non-decapod fossils are not determined yet. Palaeogeographic considerations suggest a Lower Badenian age for the site (KÓKAY pers. comm.). Decapods found: *Calappa praelata* LÖRENTHEY, *Palaeomyra globulosa* (MÜLLER) and *Maja biaensis* LÖRENTHEY.




Fig. 7. Localities near Mátraverebély




Fig. 8. Sites at Nyirád and Balatonakali

**Balatonakali**, drilling BAK-40 (Fig. 8). The drilling explored Sarmatian and Upper Badenian layers. In Badenian limestones, the following decapods were found: at 35.0 m, *Matuta brocchii* GLAESNER, at 35.5 m, *Callianassa munieri* BROCCHI, and at 37.3 m, *Callianassa pseudorakosensis* LÖRENTHEY, *Pilumnus mediterraneus* (LÖRENTHEY) and *Actumnus telegdi* (MÜLLER).

**Kishajmás**, Mecsek Mountains. An exposure of (Lower?) Badenian sandstones is situated in a deep railway-cut between the stations Husztót and Kishajmás-Szatina, along the line Budapest–Pécs. Decapods: six specimens of *Portunus monspeliensis* (A. MILNE-EDWARDS) were collected by P. SOLT.

**Fertőrákos**, drilling FRK-21 (Fig. 9). A thick sequence of (Middle?) Badenian limestone, almost depleted of macroinvertebrates was explored (HÁMOR and BOHN-HAVAS pers. com.). A 20 cm thick reef limestone was found at 268 m, which contained decapods: *Galathea* cf. *weinfurteri* BACHMAYER, *Porcellanidarum* sp. (?*Pisidia*), *Actaea turcocampestris* n. sp. and *Haydnella steiningeri* n. gen. n. sp.

**Gross-Höflein** (Burgenland, Austria, in Hungarian: Nagyhöflány), quarry "Fenk" (MNH, Fig. 9). This quarry, situated in the Leitha-Mountains, was chosen as the facio-stratotype of Badenian limestones, the so-called "Leithakalks" (STEININGER and PAPP in PAPP et al. 1978, p. 194). The




Fig. 9. Localities in the Vienna-basin

sequence was studied and described in details by DULLO (1983). He observed four horizons with reef-building corals. The second horizon from below ("oberer Korallenkalk" of STEININGER and PAPP in PAPP et al. 1978, p. 197) contains a diverse and well-preserved decapod fauna. During a stay of two days, I collected some 200 remnants of crabs. Decapods are most frequent in the upper 0.5 meter of the 2 m thick reef. All species described by BACHMAYER-TOLLMANN (1953) probably originated also from this layer. Corals: *Tarbellastraea* sp. and *Porites* sp. (both in reef-building quantity), *Stylophora* cf. *subreticulata* REUSS, *Acanthastraea* sp. and a thin foliaceous form that occurs also at Törökmező. Decapods (my collection): *Petrolisthes haydni* n. sp., *Dardanus substratiiformis* (LÖRENTHEY), *Kromtisis koberi* (BACHMAYER and TOLLMANN), *Dromiidarum* ?n. gen. n. sp., *Dymomene emiliae* MÜLLER, ?*Maja* sp., *Rakosia carupoides* n. gen. n. sp., *Daira speciosa* (REUSS), ?*Carpilius* sp., *Xantho moldavicus* (YANAKEVICH), 'Pilodius' *vulgaris* (GLAESNER), *Haydnella steiningeri* n. gen. n. sp., *Panopeus wronai* n. sp. and *Chlorodiella mediterranea* (LÖRENTHEY).

Decapods, collected and described by BACHMAYER-TOLLMANN (1953): *Cullianassa* cf. *rakosensis*, *Galathea* sp. ind., *Dromilites koberi* n. sp. (= *Kromtisis k.*), *Maia* sp. ind., *Necronectes schafferi* GLAESNER, *Chlorodopsis mediterranea* LÖRENTHEY, *Daira speciosa* (REUSS) and *Titanocarcinus vulgaris* GLAESNER. Unfortunately it was not possible for me to check these determinations. I could study only the holotype-specimen of *Dromilites koberi*. The matrix of this specimen contained some carapaces and chelae of *Chlorodiella mediterranea* (LÖRENTHEY). For statistical reasons it seems probable, however, that the list of my collection covers most of the species of BACHMAYER and TOLLMANN.

**Deutsch-Altenburg**, Niederösterreich, Austria (DAL, Fig. 9). An abandoned quarry is situated on the Hundsheimer Berg (BACHMAYER 1953a). Over a basal breccia-conglomerate, which overlays Mesozoic carbonates, a thick layer (about 30 m) of calcarenite is situated, intercalated by a thin layer of bryozoa-marl. The exposure is still accessible. Decapods, described by BACHMAYER: *Callianassa* cf. *jahringensis* GLAESSNER, *C. cf. pseudorakosensis* LÖRENTHEY, *C. cf. rakosensis* LÖRENTHEY, *C. sp.* ind., *Pagurites substratiiformis* LÖRENTHEY (= *Dardanus s.*), *Galathea weinfurteri* BACHMAYER, *Ebalia van-straeleni* BACHMAYER, *Maia austriaca* n. sp. (= *Micippa hungarica*), *Maia oroszyi* n. sp. (*Pisa o.*), *Phrynolambrus weinfurteri* n. sp. (= *Micippa hungarica*), *Cancer sismondae* H. v. MEYER (? = *C. styriacus*), *Portunites kühni* n. sp. (= *Liocarcinus kuehni*), *Carcinides oroszyi* n. sp. (= *Liocarcinus o.*), *Carpilius* sp. ind. (= *Dromiidae* sp. ind.), *Daira speciosa* (REUSS), *Chlorodopsis mediterranea* LÖRENTHEY (= *Pilumnus mediterraneus*), *Eriphia* sp. ind. (? = *Necroneutes schafferi*), *Titanocarcinus vulgaris* GLAESSNER (= *Xantho moldavicus*) and *Pilumnoplax carnuntinus* n. sp.

**Oslip** (Burgenland, Austria, in Hungarian: Oszlop), sand-pit (Sandgrube Bauer, MOS, Fig. 9). A detailed description and paleoenvironmental analysis is given by DULLO (1983). Over a sand-layer, apparently depleted of fossils, a one meter thick *Ophiomorpha* (*Callianassa* burrows) sand contains many fragments of chelae. The sand is covered by a red-algae bearing limestone. Decapods found during a stay of an hour: *Calappa praelata* LÖRENTHEY, *Parthenope tetenyensis* n. sp. and *Liocarcinus cf. rakosensis* (LÖRENTHEY).

Other Austrian decapod-bearing sites are described by GLAESSNER 1928.

**Grobie**, Holy Cross Mountains, Southern Poland. Described by FÖRSTER (1979a). There is no exposure, blocks of red algal limestone are scattered on cultivated terrain. A remnant of a patch-reef of *Tarbellastraea* sp. yielded the crab-fossils (collected by R. WRONA), according to FÖRSTER: ?*Eriphia* sp. C. (= *Daira speciosa*), *Pilumnus* sp. (fingers ? = *Pilumnus* sp., carapaces = *Dynomene emiliae*), ?*Eriphia* sp. D. (= *Panopeus wronai*).

**Węglinek**, Eastern Poland. Described and collected by KRACH (1981). A small abandoned quarry yielded decapods, some others were found on cultivated terrain. Decapods: *Callianassa* sp., *Maia bivensis* LÖRENTHEY, *Xantho moldavicus* (YANAKEVICH), *Pilumnus mediterraneus* (LÖRENTHEY) and *Liocarcinus rakosensis* (LÖRENTHEY).

Other decapod-bearing localities in Poland are described by A. E. REUSS (1867) and by FÖRSTER (1979a, b).

YANAKEVICH (1969, 1977) gave a monographical account about Upper Badenian reefal limestones. In localities at villages **Koban**, **Shepteban**, **Varatik**, **Gorodishtshe** and **Bolotino** (Moldavian SSR.) he collected the following decapods: *Daira speciosa* (REUSS) and *Medaeus moldavicus* n. sp. (= *Xantho* m.).

Bulgarian sites (MÜLLER 1979b) are described by KOJUMDGIEVA (1975). These are: **Iasen** (yielded *Dynomene emiliae* MÜLLER), **Gorna Mitropolia** (*Ranidina rosaliae* BITTNER), **Krivodol** (*Liocarcinus* sp.) and **Oriakhovo** (*Miocyclus bulgaricus* MÜLLER, *Goneplax* cf. *gulderi* BACHMAYER).

**Tasádfő** (Transsylvania, now Romania) a site, described by LÖRENTHEY (1898b, c) yielded "Cancer" *szontaghii* LÖRENTHEY. No detailed description is available about the sequence of layers.

**Zagor** (Jugoslavia, Slovenia) a locality, which yielded *Cancer illyricus* BITTNER 1884, was not described in details yet.

I visited most of the above mentioned localities, while in other cases I could revise their material. Other, not visited localities, which yielded only sporadic specimens, are mentioned in the descriptive part.

Speciation of *Brachyura* is rather rapid. All known, well-described Pleistocene species are identical with recent ones, but the Pliocene record contains several extinct species (GLAESSNER 1929). The development of recent genera was also fast. Most of them probably originated in the Miocene or Pliocene (MÜLLER 1979b). Related Badenian and extant species often show defined morphological differences, e.g. *Pachygrapsus hungaricus* vs. *P. marmoratus*, or *Ebalia oersi* vs. *E. edwardsi*. Consequently identification of 12–15 million year old species with extant ones should be done with caution if no link is known from intermediary times. Therefore, even when Badenian forms have morphologically similar recent counterparts, new names were given. In certain cases this is clearly a temporary compromise.

The same approach is not possible on generic level thus we used the names of recent genera without comments if a miocene species falls within the range of variation of recent species of the genus. Single quotation marks ('.....') were used to denote collective genera i.e. when generic determination was not possible due to the incomplete preservation of fossil forms, but relationship with the nominate genus is probable. When no significant similarity with any recent or fossil genus was seen, new generic names were given to new species. Quotation marks (".....") were used for species which were erroneously placed in well-known genera but due to poor preservation or loss of types (e.g. "Cancer" *carniolicus*) could not be assigned to the correct genus.

Fingers are the most frequent remnants of fossil crabs. These functionally important organs are generally well calcified and robust. Attempts of their systematic study (BACHMAYER 1962, JANSSEN 1972, FÖRSTER 1979a, b) gave only moderate success. The reason for this was discussed in detail by FÖRSTER (1979b, p. 254). It should be mentioned that zoologists and paleontologists often neglect to provide detailed descriptions or figures of fingers, making the study of a specimen impossible if original specimens of related forms are not available. Fingers found in carapace-bearing layers, assuring their determination on a higher level of probability, were depicted only for a limited number of species (e.g. GLAESSNER 1928, LÖRENTHEY-BEURLEN 1929, BACHMAYER 1953a). One of the main purposes of my study was to attempt the identification of individual parts, including fingers, of disarticulated decapods. For this reason I compared them with related recent forms. The relative frequency of the pieces in a given layer gives indication about their relation, too. In some localities fingers were found attached to palms. Decoration of carapaces are often similar to that of palms of the same species, which permits determination of isolated chelae. If the mentioned goal was achieved to some extent, a step was made toward the utilization of individual fingers so frequently found in sites from which only molluses or few other invertebrates have been described so far. It is expected that information on decapods will significantly contribute to stratigraphic, palaeoecologic and palaeogeographic studies. The usefulness of such information comes from the mentioned fast evolution of some decapod groups and the ability for rapid migration of many species (e.g. "Lessepsian migration", POR 1978) due to mobility of adults to a rather prolonged pelagic larval stage of development, making even transoceanic migrations possible (GARTH 1974).

## SYSTEMATICAL PART

In the following part the species will be presented in the systematic order given by GLAESSNER (1969), for the sake of convenience. This or similar systems were used in most of the palaeontological works during the last decade, therefore a comparison will be eased using GLAESSNER's system. In spite of this the author is fully convinced that GUINOT (1977) and DE SAINT-LAURENT (1980a, b) showed the way to a better understanding of the systematics of *Brachyura*.

In the systematic part the synonym lists will enumerate works containing new data or any other emendations of the given taxon. Consequently, Fossilium Catalogus (GLAESSNER 1929) and Treatise on Invertebrate Paleontology (GLAESSNER 1969) will not be mentioned in spite of their fundamental importance unless they modify the taxonomic status of the species in question.

Abbreviations used in this part, for collections: NMÖ: Department of Natural History Museum of Budapest (p. 34), FI: Geological Survey of Hungary, CA: my collection, NHMW: Natural History Museum in Vienna. Other collections are not abbreviated. The inventory numbers of collection CA, refers to the codes as defined in chapter: "Description of localities" (Table 1).

Abbreviations for parts of crabs: ca.: carapace, ch.: chela, da.: dactylus, ff.: fixed finger.

The type-species of the genera are not given unless they are included in GLAESSNER (1969).

### Ordo DECAPODA LATREILLE, 1803

Subordo PLEOCYEMATA BURKENROAD, 1963

Infraordo A N O M U R A H. MILNE-EDWARDS, 1832

Superfamilia Thalassinidea LATREILLE, 1831

Familia Laomediidae BORRADAILE, 1903

Genus *Jaxea* NARDO, 1847

### *Jaxea kuemeli* BACHMAYER, 1954

1954. *Jaxea kümeli* n. sp. — BACHMAYER: 64—66, pl. 1, figs. 1, 2.

Material not seen: holotype, ca. 68/1954 ?ch. Martinsdorf: NHMW.

Description: According to BACHMAYER, the body is elongated, rather narrow. The cervical furrow is strong, and situated posteriorly, the linea thalassinica is visible. Some segments of the abdomen are preserved, these are elongated. A left chela, probably belonging here, bears small spines on its surface.

Remarks: According to BACHMAYER, the species is extremely close to *J. nocturna* NARDO. The recent species is more slender.

### Familia Callianassidae DANA, 1852

Remarks: In conventional classification (GLAESSNER 1969 pp. R477—R478) this family includes not only *Callianassa* and related forms but *Upogebia* as well. The convincing studies of DE SAINT-LAURENT-LE LOEFF (1979) argued, however, for the phylogenetic independence of the mentioned two genera.

**R e m a r k s :** The most abundant of all fossil decapods are remnants of isolated chelae similar to those of some recent *Callianassid*-genera, as *Callichirus* and *Callianassa* (DE SAINT-LAURENT-LE LOEUFF 1979). The chelae are highly adapted for digging in having sharp edges. The chelae bear many characteristic features, consequently species are easily distinguished from each other, but the carapace is far less characteristic and rarely preserved as fossil. Most of recent *Callianassids* are strongly heterochelous. Surprisingly, in the diverse Badenian material no evidence of heterochely was observed, as right and left chelae are found in same proportion and in same size as well, with the exception of *C. brocchii*.

Consequently the genus '*Callianassa*' is regarded as a collective taxon.

Recent *Callianassa* species occur in any depth from the littoral zone down to some hundreds of meters, so the presence of their chelae or burrows (*Ophiomorpha* LUNDGREN) are by no means indicators for intertidal conditions (WEIMER-HOYT 1964).

### '*Callianassa*' *munieri* BROCCHI, 1883

Pl. I, figs. 1-7, pl. II, figs. 1, 2.

- 1883. *Calianassa munieri* n. sp. — BROCCHI: 5, pl. 5, figs. 5, 6.
- 1897. *Calianassa munieri* BROCCHI — LÖRENTHEY: 150, 168.
- 1898a. *Calianassa munieri* BROCCHI — LÖRENTHEY: 93, 105, 114.
- 1898b. *Calianassa munieri* BROCCHI — LÖRENTHEY: 105, 132, 155.
- 1898c. *Calianassa munieri* BROCCHI — LÖRENTHEY: 104.
- 1904a. *Callianassa munieri* BROCCHI — LÖRENTHEY: 161.
- 1904b. *Callianassa munieri* BROCCHI — LÖRENTHEY: 30.
- 1911. *Calianassa munieri* BROCCHI — LÖRENTHEY: 522.
- 1913. *Calianassa munieri* BROCCHI — LÖRENTHEY: 322.
- 1929. *Calianassa munieri* BROCCHI — LÖRENTHEY: 33, 62, 63, 64, pl. 2, figs. 19-23, not: fig. 24.
- partim 1929. *Portunus rákosensis* n. sp. (merus only!) — LÖRENTHEY: 173, pl. 12, figs. 22, 23.
- 1979a. *Callianassa munieri* BROCCHI — MÜLLER: 274.

M a t e r i a l not seen, probably lost: syntypes, at least 2 ch. Rákos: ?Sorbonne.

M a t e r i a l seen: 14 ch. Rákos, Kamaraerdő, Bia: NMÖ. About 100 ch. Rákos, Budavárs, Bia: FI. 31 ch. MR-5, 15 ch. MR8-1, 1 ch. MR45-2, 18 ch. MRF-1, 2 ch. MKC-5, 4 ch. MEG-2, 12 ch. MFA-3, 1 ch. drilling BAK-40 m, 1 ch. MZZ-9: CA.

**D e s c r i p t i o n :** Both right and left chelae are found frequently. The palm is approximately square, slightly longer than high and converging distally. The upper margin is keeled on its proximal 1/3 and bears two small teeth on its distal part. The lower edge is sharp. Small pores are situated along its inner side. The palm is decorated with numerous tubercles confined to two areas on both sides. The dactylus is rounded on its upper margin while keeled on the lower one, where it bears a tooth-like, elongated elevation. At some specimens the dactylus is uncommonly high and stout. The carpus is decorated with a row of teeth on upper edge. Meri were not found in connection with other articles, but they are typically *Callianassa*-like and their size rules out any other species than *C. munieri*.

Stratigraphically it seemed to be restricted to Upper Badenian, zone 3. (see p. 103), but recently a specimen was found in Zebegény (code MZZ), in Lower Badenian stratum. Ecologically the species is bound to fine homogenous sands, it is rather rare or absent in medium-to-coarse grained calcarenites. Known from Budapest, Bia, Balatonakali and Zebegény.

### '*Callianassa*' *chalmasi* BROCCHI, 1883

Pl. II, figs. 3-7.

- 1883. *Calianassa chalmasi* n. sp. — BROCCHI: 6, 7, pl. 5, figs. 7, 8.
- 1897. *Calianassa chalmasi* BROCCHI — LÖRENTHEY: 150, 160, 168.
- 1898a. *Calianassa chalmasi* BROCCHI — LÖRENTHEY: 93, 105, 114.
- 1898b. *Calianassa chalmasi* BROCCHI — LÖRENTHEY: 9, 129, 130, 155.
- 1898c. *Calianassa chalmasi* BROCCHI — LÖRENTHEY 102.
- 1904a. *Callianassa chalmasi* BROCCHI — LÖRENTHEY: 161.
- 1913. *Calianassa chalmasi* BROCCHI — LÖRENTHEY: 322.
- 1929. *Calianassa chalmasi* BROCCHI — LÖRENTHEY in LÖRENTHEY-BEURLEN: 33, 65, pl. 2, figs. 15a-15b.
- 1961. *Callianassa espichelensis* n. sp. — VEIGA FERREIRA: 479-481, pl. 1, figs. 1-11.
- 1979a. *Callianassa chalmasi* BROCCHI — MÜLLER: 274, pl. 1, fig. 5.

Material not seen, probably lost: syntypes, at least 2 ch. Rákos: ?Sorbonne.

Material seen: about 100 ch. Rákos: FI. 12 ch. Rákos: NMÖ. 53 ch. MR-6, 19 ch. MR8-2, 4 ch. MR45-1, 44 ch. MRF-2, 1 ch. MRO-2, 1 ch. MEG-1, 5 ch. MOH-1, 20 ch. MGY-9: CA.

Description: Both right and left chelae are similarly frequent. The palm is converging distally. Both upper and lower edge sharp. On the inner side, parallel to the edges two rows of pores of setae are present. The upper ones are arranged perpendicular to the longitudinal axis of the limb as they are elongated. Their number is 8 to 9. The lower ones are more numerous as they extend to the immovable finger as well. On the outer side of the fixed finger there is a ridge leading toward the palm. On the cutting edge there is a tooth. The dactylus is moderately stout and bears characteristic pores along two rows.

The species is frequently found in coarser sands and calcarenites too, in contrast to *C. munieri*. Its stratigraphic range is Upper Badenian to Tortonian (in Portugal). Known from Budapest and Portugal.

### 'Callianassa' *rakosiensis* LÖRENTHEY, 1897

Pl. III, figs. 1, 2.

1897. *Calianassa rakosiensis* n. sp. — LÖRENTHEY: 160, 161, 168, 169.

1898a. *Calianassa rakosiensis* LÖRENTHEY — LÖRENTHEY: 105, 114, 115.

1898b. *Calianassa rakosiensis* LÖRENTHEY — LÖRENTHEY: 131, 132, pl. 9, fig. 4.

1898c. *Calianassa rakosiensis* LÖRENTHEY — LÖRENTHEY: 103–104, pl. 9, fig. 4.

1929. *Calianassa rakosiensis* (sic!) LÖRENTHEY — LÖRENTHEY in LÖRENTHEY—BEURLEN: 33, 66, 67, pl. 2, fig. 14.

1979a. *Callianassa rakosiensis* LÖRENTHEY — MÜLLER: 274, 276, pl. 1, fig. 4.

Material not seen, probably lost: holotype, 1 ch. Rákos: ?FI.

Material seen: 2 ch. MR-7, 2 ch. MR8-6, 3 ch. MGY-6: CA.

Description: The palm is converging distally. Its upper edge is almost straight, sharp. On the inner side there is a row of 9 to 10 pores. Its lower edge is slightly curved, proximally the curvature is stronger than distally. The lower edge is sharp and keeled. A ridge is situated on the outer surface of the fixed finger, which is curved and its portion on the palm is subparallel to the longitudinal axis, more proximally it is bended upwards. Around the proximal end of this ridge the surface is decorated by tiny tubercles.

Remarks: The species is remarkably similar to *C. chalmasi* in its general form, size and the distribution of pores. The form of the ridge on the finger and the position of the tooth is completely different in the two species and no trend was observed toward a transitional form. Thus their independence seems highly probable though a close relation is likely.

### *Callianassa brocchii* LÖRENTHEY, 1897

Pl. III, fig. 3.

1897. *Calianassa brocchii* n. sp. — LÖRENTHEY: 161, 168, 169.

1898a. *Calianassa brocchii* LÖRENTHEY — LÖRENTHEY: 106, 114, 115.

1898b. *Calianassa brocchii* LÖRENTHEY — LÖRENTHEY: 132–134, 155, pl. 9, fig. 5.

1898c. *Calianassa brocchii* LÖRENTHEY — LÖRENTHEY: 104–105, pl. 9, fig. 5.

1929. *Calianassa brocchii* LÖRENTHEY — LÖRENTHEY in LÖRENTHEY—BEURLEN: 33, 64, pl. 2, fig. 13.

1979a. *Callianassa brocchii* LÖRENTHEY — MÜLLER: 274.

Material seen: holotype, ch. Rákos, M2329: FI. 2 ch. Budafok: FI. 3 ch. MGY-24: CA.

Description: All known specimens are right chelae suggesting heterochely in this single Badenian species. The palm is rather convex on its outer surface while flat on the inner one. Its outline is close to a square. The upper edge is not sharp while the lower one is decorated with a characteristic saw-like row of small denticles. Just on the inner surface of this edge there is a row of pores situated densely on the palm sparcely on the finger. The cutting edge of the fixed finger is denticulated with sharp teeth. The dactylus is strong, rather stout.

Remarks: The species, because of its probable heterochely and by the form of its finger is closer to some recent forms (e.g. *Callianassa gilchristi*, *C. natalensis*, cf. BARNARD 1950, p. 510) than any of the other Badenian forms. The saw-like lower edge, however, separates it from the mentioned species.

**'Callianassa' pseudorakosensis LÖRENTHEY in LÖRENTHEY—BEURLEN 1929**

Pl. III, figs. 4, 5, pl. IV, figs. 1–6, pl. V, figs. 1–4.

1929. *Calianassa pseudorakosensis* n. sp. — LÖRENTHEY in LÖRENTHEY—BEURLEN: 33, 67, 68, 69, pl. 4, figs. 16, 17, 18.

?1953. *Callianassa* cf. *pseudorakosensis* LÖRENTHEY — BACHMAYER: 241.

1976a. *Callianassa pseudorakosensis* LÖRENTHEY — MÜLLER: 507.

Material seen: lectotype, designated herein: right ch. with dactylus, Rákos, M 139 pl. 3, figs. 4, 5 pl. 4, fig. 1. Paralectotype: ch. Felménes, Transsylvania, M 2334: FI.

Material not seen, probably lost: paralectotypes: 9 ch. Rákos, 1 ch. Budafok, 10 ch. Bia, 26 ch. Felménes, numerous ch. Mátraverebély-Szentkút (?Szent László-forrás).

Material seen: 33 ch. Mátraverebély-Szentkút, M83/164, 1 ch. Mátraverebély-Szentkút, M83/160, 13 ch. M83/152: NMÖ. 19 ch. MR-4, 6 ch. MR8-5, 3 ch. MRO-1, 6 ch. MGY-7, 22 ch. MFA-1, 22 ch. MFK-1, 7 ch. MA-3, 5 ch. Bántapuszta, Várpalota, Carpathian stage, ch. BAK-40, 37.3 m: CA.

**Description:** There are two distinct forms, occurring on both right or left chelae, most probably due to a dimorphism. The palm is the same at both forms. It is rectangular, rather thick. The upper edge is keeled on its proximal half while rounded on the distal one. The lower edge is sharp, straight or slightly curved. There is a row of pores along this edge. The index is short and stout. The two forms differ in the existence or absence of a tooth on the index. Anterior this tooth the cutting edge is finely denticulated. The dactylus is curved, stout, its cross-section is circular. It bears a strong basal tooth.

**Ecology:** The species is most abundant in rather coarse sands and occurs frequently also in limestones. Probably its robust chela fits for digging sediments containing big particles.

Stratigraphically the species is characterised by a long record since it is known from the Carpathian to the Upper Badenian, zone 3. The Carpathian and the Lower Badenian specimens are much bigger than the Upper Badenian ones.

**'Callianassa' kerepesiensis MÜLLER, 1979**

Pl. VI, figs. 1–6.

1976a. *Callianassa* sp. nov.? — MÜLLER: 508, pl. 1, figs. 1–4.

1976b. "Callianassa" *kerepesiensis* n. sp. — MÜLLER: 149, 150, 154, pl. 1, figs. 1–5.

?1979a. *Callianassa kerepesiensis* MÜLLER — FÖRSTER: 90, text-fig. 1, pl. fig. 5.

1979a. *Callianassa kerepesiensis* MÜLLER — MÜLLER: 274, 276, pl. 1, figs. 1–3.

Material seen: holotype, pl. VI, fig. 1–2, MKC-2-1: CA. Paratypes, 2 ch. MA-2, 100 ch. MKC-2: CA. 2 ch. MA-2, 50 ch. MOE-11, 30 ch. MGY-8: CA.

1 ch. Mátraverebély-Szentkút, M83/153: NMÖ.

**Description:** The palm is rectangular, slightly elongate at some specimens. Both lower and upper edges are sharp. There is a characteristic undulating region on the inner surface near the lower edge. The fixed finger is rather slender. It bears two teeth, one, almost rudimentary, is situated near the proximal end of the finger, the bigger one is near to its end. There is a denticulated cutting edge between the two teeth, at least this is observable on bigger specimens. The dactylus is slender and it bears two rows of pores.

Stratigraphically, the species was found near the beginning of the Badenian (Szentkút) as well as at its end (MOE). Ecologically it was a tolerant species as it was found in many different types of sediments.

**'Callianassa' cf. *kerepesiensis* MÜLLER 1979**

Pl. VII, figs. 1, 2.

1979a. *Callianassa kerepesiensis* MÜLLER — FÖRSTER: 90, text-fig. 1, pl. 1, fig. 5.

Material not seen: right ch. with carpus, Nawodzice: Coll. Univ. of Warsaw.

Material seen: ch. MOE-12: CA.

**Remarks:** As FÖRSTER stated, the species is remarkably similar to *C. kerepesiensis*, but its fixed finger bears only one, rudimentary tooth near its end, and the undulated region lacks on the inner side of the palm.

**'Callianassa' cf. jahringensis GLAESSNER 1928\***

1928. *Callianassa* cf. *jahringensis* n. sp. — GLAESSNER: 166–167.

M a t e r i a l not seen: ch., 3 da. Vöslau: NHMW.

R e m a r k s : According to GLAESSNER, the palm is square but the two edges are rather convex. On the other hand the specimen, as far as this is possible to be observed on figures, is not very much like *C. jahringensis*, Carpathian species. Most probably it belongs to a new species. Stratigraphically it is confined to Lower Badenian.

**'Callianassa' sismondae A. MILNE-EDWARDS 1860**

1928. *Callianassa sismondae* A. MILNE-EDWARDS — GLAESSNER: 168, giving the former synonyms.

M a t e r i a l not seen: 2 ch., da. Spielfeld: NHMW.

**'Callianassa' norica GLAESSNER 1928**

1928. *Callianassa norica* n. sp. — GLAESSNER: 169–170, text-fig. 1., pl. 3, figs. 3, 4.

M a t e r i a l not seen: 5 ch., 1 fixed finger, 8 da. Spielfeld, holotype 1860/V/8 b: NHMW.

R e m a r k s : The species is different from any Badenian forms as its palm is converging proximally.

**Callianassa sp. ind.**

Pl. V, figs. 5, 6.

1954. *Semiranina* n. gen. *oroszyi* n. sp. — BACHMAYER: 66, 67, pl. 1, figs. 4, 5.

1974b. *Callianassa oroszyi* (BACHMAYER) — MÜLLER: 276, pl. 1, fig. 3.

M a t e r i a l not seen: holotype of *Semiranina oroszyi*, ca. Deutsch-Altenburg, 69/1954: NHMW.

M a t e r i a l seen: 3 ca. MDG-1: CA.

D e s c r i p t i o n : The specimens represent the anterior parts of carapaces of Callianassa, being a semicylindrical, smooth part situated before the cervical groove.

R e m a r k s : The specimens do not offer any possibility for a determination beyond the generic level as any carapace of this kind is essentially the same.

**?Callianassa szobensis** n. sp.

Pl. VII, figs. 3, 4, ?figs. 5, 6.

M a t e r i a l seen: holotype, ch. MSZ-1-1. Other, ?da. MSZ-2-1: CA.

L o c u s t y p i c u s : Szob, Börzsöny Mountains (Fig. 6), Nagyfeltárás.

S t r a t u m t y p i c u m : Lower Badenian NN 5 Nannozone (pers. comm. A. NAGYMAROSI).

D e r i v a t i o n o m i n i s : From the village Szob, near the type-locality.

D i a g n o s i s : Long, flat, rectangular palm decorated by big granulae near the base of the fixed finger.

D e s c r i p t i o n : The palm of the left chela is twice as long as high, quadratic. The upper and the lower margins are sharp, moderately convex on the ends. On the inner side, along the margins there are rows of pores. Characteristic is the decoration on the outer side. Near the base of the fixed finger there are strong tubercles scattered around a horizontal line. This line can be followed up to the proximal 3/5 of the total length of the palm. The fixed finger is moderately long. There is a prominent tooth on it. Between the base of the finger and this tooth the cutting edge is denticulated. A finger, probably a dactylus was found in the same layer, characterised by rows of tubercles.

R e m a r k s : The peculiar decoration on the outer side does not suggest any closer relationship with recent or fossil *Callianassids*, though the species might belong to other families as well.

\* After closing the manuscript, I studied the material. The specimens belong to '*C. pseudorakosiensis* LŐRENTHEY.

Remarks : According to DE SAINT-LAURENT-LE LOEFF (1979), the groups of *Callianassa* and *Upogebia* should be regarded as different families as the similarities among them are due to convergency owing to their similar habits.

Genus *Upogebia* LEACH, 1814

**Upogebia** sp. (div.?)

Pl. VIII, fig. 1, ?figs. 2-6.

1974b. *Upogebia scabra* n. sp. — MÜLLER: 276, 283, pl. 1, figs. 1, 2.

1979a. *Upogebia radula* (error!) — MÜLLER: 274.

1979a. *Upogebia* sp. — MÜLLER: pl. 2, figs. 1-3.

Material seen: holotype of *Upogebia scabra*, MDG-3-1: NMÖ. 2 ch. MDG-3, ch. MRO-5, 3 ch. MOE-10, 6 ch. MGY-10: CA.

Description : The carapace is typically that of *Upogebia* but offers little if any possibility for a determination beyond generic level. Only the precervical part was calcified in an extent making possible fossilisation. The posterior half of this part is elliptical while the anterior is slightly diverging. Three furrows are arranged longitudinally. The surface between the furrows is rough.

Remarks : As in the case of *Callianassa*, the carapace of *Upogebia* is rather uniform making specific separations impossible. The chelae are probably better for determination but at present the scarce material does not allow descriptions. Possibly two or even more species occurred in Badenian seas.

Superfamilia Paguroidea LATREILLE, 1803

Familia Paguridae LATREILLE, 1802

Subfamilia Paguriinae LATREILLE, 1802

Genus *Pagurus* FABRICIUS, 1775

Remarks : This abundant and widespread genus was rarely recognised as a fossil but numerous species was described under its name, partly because the confusion of *Pagurus* and *Dardanus*, partly because the wellknown name was used as a collective one.

Ecology : *Pagurus*-species generally avoid shallow waters and intertidal regions. Most species live below 40 meters down to several hundreds of meters. Presumably most species are euhaline.

**Pagurus rakosensis** MÜLLER, 1979

Pl. X, figs. 1-5.

1979a. *Pagurus rakosensis* n. sp. — MÜLLER: 274, 277, 286, pl. 5, fig. 3, pl. 6, fig. 1.

Material seen: holotype, ch. MR-1-1. Paratypes, 3 ch. MGY-11, 14 ch. MOE-6. Others, 2 ch. MR8-6, ch. MEG-5: CA.

Description : The palm of the right chela is elliptical in cross-section. Its upper margin is straight, slightly flattened, decorated by scattered tubercles. On the median part of the outer surface there is a row of tubercles. Between the upper edge and the median line there is a slightly depressed area. The lower edge of the palm and fixed finger forms an arc the curvature of which is stronger near the proximal end. On this edge there are medium-sized tubercles. The other parts of the outer surface are covered with smaller tubercles. The inner side of the palm is decorated with short transverse ridges of squamae-like protuberances. The tip of the fixed finger is hard, strongly calcified.

**R e m a r k s :** The species is close to *P. bernhardus* (L.) but its decoration is less coarse. An other similar species, *P. variabilis* (A. MILNE-EDWARDS-BOUVIER) has an even finer granulation on surface.

**E c o l o g y :** The species is known from localities of finer or coarser quartz-sand or of calcarenite with shell-debris. It was not found either in deposits of shallow sublittoral environments, or in coral-reefs.

***Pagurus concavus* MÜLLER, 1979**

Pl. IX, figs. 1-6.

1979a. *Pagurus concavus* n. sp. — MÜLLER: 274, 277, 286, pl. 7, figs. 1-5.

**M a t e r i a l seen:** holotype, ch. MOE-1-1. Paratypes, 23 ch. MOE-1. Others, 130 ch. MOE-1, 1 ch. MEG-4, 2 ch. MOH-3: CA.

**D e s c r i p t i o n :** Both of the right and left chelae are known. Both are characterised by three strong finely denticulated carinae on the outer side of the palm. Consequently two depressed longitudinal areas are delimited. The upper margin of the palm of the right (crusher) chela is flattened. The upper and the median carinae are parallel to each other while the lower one is situated at the lower margin and is arched. The outer and upper surfaces are densely covered by small granulae. The fixed finger is triangular in cross-section. The left (nipper) chela is much smaller and more slender than the other. The upper carina is less developed. The decoration is the same as that of the right chela. The index is relatively long. The teeth of the cutting edges, according to their role, differ on both sides as shown in the illustrations.

**R e m a r k s :** The chelae are rather similar to those of *P. alatus*, a recent species, but the latter has an arched upper margin on the right palm. The depressions are much deeper in the recent species. *P. sculptimanus* has a strongly denticulated upper margin on the palm. The Badenian species is closely related to *P. alatooides* SECRETAN, Burdigalian species from Courennes, France (PHILIPPE-SECRETAN 1971, p. 15). The Burdigalian species has dentated carination on both chelae, on the upper and on the median carinae there are 5 to 6 prominent teeth.

Ecologically and stratigraphically *P. rakosensis* and *P. concavus* seem to be close to each other.

**'*Pagurus*' *albus* MÜLLER, 1979**

Pl. XI, figs. 1-3.

**M a t e r i a l seen:** holotype, ch. MOE-2-1. Paratypes, 5 ch. MOE-2. Others, ?ch. MR-3, ch. MJL-2: CA.

**D e s c r i p t i o n :** The right chela is rather stout. The outer surface is densely covered with tubercles which are spiniform on the upper margin and at the distal end of the lower one. The tuberculae are rather irregularly scattered on the surface. The lower margin is strongly curved, its outline is circular. The fixed finger is turned upwards.

**R e m a r k s :** The only species which reveals similarities with *P. albus* is *P. sculptimanus* (ZARIQUIEY ALVAREZ 1968, p. 234.) with similar decoration and outline. The deep furrows are absent from the Badenian species, however.

**Genus *Anapagurus* HENDERSON, 1886**

**R e m a r k s :** The recent genus includes several species having simple chelae with few diagnostically useful features. Unfortunately, good illustrations lack in the literature. Consequently the two Badenian species can not be compared properly with recent ones.

**'*Anapagurus*' *miocenicus* MÜLLER, 1979**

Pl. XI, figs. 4-9.

1979a. *Anapagurus miocenicus* n. sp. — MÜLLER: 274, 277-278, 286, pl. 6, figs. 2-5.

**M a t e r i a l seen:** holotype, ch. MGY-1-1. Paratypes, MGY-1. Others, 6 ch. MR8-9, ch. MEG-6: CA.

**Description:** Few characteristic features are on the chela. Even on SEM micrographs only a small protuberance was found on the proximal end of the palm. The teeth of the cutting edge are characteristically *Pagurinae*-like, bifurcating proximally. The palm is extremely stout. The lower edge is strongly curved.

**Remarks:** This featureless species is similar to its recent relatives, which have generally more slender chelae.

**'Anapagurus' marginatus MÜLLER, 1979**

Pl. XII, figs. 1-5.

1979a. *Anapagurus marginatus* n. sp. — MÜLLER: 274, 278, 286-287, pl. 4, figs. 1-3.

**Material seen:** holotype, ch. MOE-3-1. Paratypes, 8 ch. MOE-3, ch. MJL-3, 6 ch. MDG-5: CA.

**Description:** The palm of the species is rather variable in its general form, there are rather stout and more slender specimens. The lower margin is slightly curved and there is a carina on its outer part. The carinae consist of very fine tubercles. There is a proximal elongated tooth on the upper part of the palm. The outer surface is rather smooth, on some specimens a rudimentary granulation is visible. The teeth on the cutting edge are typically *Pagurinae*-like.

**Remarks:** The chelae are more characteristic than those of *A. miocenicus*, having a characteristic carina on the lower edge. Probably this feature separates this species from other *Pagurinae* species.

**?Pagurus turcus n. sp.**

Pl. XII, fig. 6, pl. XIII, figs. 1-4.

**Material seen:** holotype ch. MTZ-17-1. Paratype, 2 ch. MTZ-17: CA.

**Locus typicus:** Nagymaros-Törökmező, southern Börzsöny Mountains, small quarry, code MTZ.

**Stratum typicum:** Lower Badenian.

**Derivatio nominis:** The name of the locality, Törökmező, means "meadow of Turks", lat.: Turcus, campus.

**Diagnosis:** Right chelae decorated by randomly scattered tubercles.

**Description:** The right chelae are moderately stout. Their upper edge is straight, the lower one curved. There are no rows on the surface, instead the tubercles are apparently randomly distributed on the surface. Their size varies within a narrow range. The smaller ones are distributed among the others mainly at the base of the fixed finger. The inner surface is decorated with asymmetrical, bigger elevations resembling scales.

**Remarks:** The chela is unlike any other fossil Pagurids by its peculiar decoration. On the other hand it was by no means possible to compare it with all living species as most illustrations are diagrammatic, offering few possibilities for comparison with fossil species. No similar form lives in the Mediterranean.

**Genus *Pylopagurus* A. MILNE-EDWARDS-BOUVIER, 1891**

**'Pylopagurus' leganyii n. sp.**

Pl. XIV., figs. 1-3., ?figs. 4, 5.

**Material seen:** holotype, ch. M62/3253. Sámonháza-Csúcs-hegy alja: NMÖ. ?internal moulds of 3 ch. MTZ-18: CA.

**Locus typicus:** Sámonháza, Csúcs-hegy alja (Noszky 1940).

**Stratum typicum:** Lower Badenian.

**Derivatio nominis:** From F. LEGÁNYI, amateur palaeontologist, who collected the type-specimen.

**Diagnosis:** Stout right chela densely covered by blunt tubercles.

**Description:** The chela is very stout, but flattened near the lower distal part. The lower margin is strongly curved, especially at the fixed finger. The upper margin is straight, flattened. The

outer surface is decorated with an extremely dense array of blunt tubercles, scattered irregularly. The cutting edge of the fixed finger is very wide, having teeth on its both sides. The fixed finger is bended inward.

**R e m a r k s :** In its general form, the chela is like recent *Pylopagurus* species, as *P. liochele*, *P. unguilatus* but similarity fails in details.

**Pagurinae sp. n.?** div.

Pl. XV, figs. 1-5.

**M a t e r i a l seen:** ch. MKC-11, ch. MRF-12: CA.

**R e m a r k s :** Two right chelae, belonging to two different species certainly distinct from any other Badenian species by their peculiar decoration. Since no fixed fingers were found they will not be described here as new species yet.

**Subfamilia Diogeninae ORTMANN, 1892**

**Genus *Diogenes* DANA, 1851**

***Diogenes matrensis* n. sp.**

Pl. XVI, figs. 1-3.

**M a t e r i a l seen:** holotype, ch. Mátraverebél-Szentkút, M61/2633, leg. Streda: NMÖ.

**L o c u s t y p i c u s :** Mátraverebél-Szentkút (Fig. 7), probably the lowest layer at the spring "Szent László-forrás".

**S t r a t u m t y p i c u m :** Lower Badenian.

**D e r i v a t i o n o m i n i s :** From the Mátra Mountains situated in the vicinity of the type-locality.

**D e s c r i p t i o n :** The palm is flattened, high, short. The lower edge is gently curved, convex proximally, slightly concave distally. It is rather sharp, the keel is formed by a row of dense teeth. The upper edge is also slightly convex decorated with three rows of rather irregularly arranged tubercles. The outer side of the palm is tuberculated near to the upper edge but smooth on the rest of the surface. The fixed finger is remarkably short and stout. At the half of its length it bears a big tooth while smaller teeth are situated before and behind the big one.

**R e m a r k s :** The species is quite distinct from the Mediterranean species of the genus but similar to some African species. The similarity is most striking with *D. extictatus* STEBBING (BARNARD 1950, p. 441.) but fails in the length of the fixed finger.

***Diogenes* cf. *pugilator* (ROUX, 1828)**

Pl. XVI, figs. 4, 5.

**M a t e r i a l seen:** ch. drilling "Váci út 120" 17 m. Budapest, ch. MR8-21, 3 ch. MGY-26: CA.

**D e s c r i p t i o n :** The left chela is rather stout, convex on both sides. Both edges are more or less rounded. The upper edge is decorated with irregularly scattered granulae. The rest of the surface is rather smooth as far as it was possible to observe this on the corroded specimens. The finger is moderately long, directed downwards.

**R e m a r k s :** The species is remarkably similar to *Diogenes pugilator*, a recent Mediterranean species, but its decoration is much poorer. Probably, when better specimens are available a new species will be recognised in it.

**Genus *Dardanus* PAULSON, 1875**

**Dardanus arrosor (HERBST, 1796)**

Pl. XVII, fig. 5.

1803. *Pagurus striatus* n. sp. — LATREILLE: 163.

1861. *Pagurus substriatus* n. sp. — A. MILNE-EDWARDS in E. SISMONDA: 20.

1891. *Pagurus arrosor* (HERBST) — RISTORI: 23.

M a t e r i a l seen: moulds of ch. MTZ-5, ch. MTH-1: CA.

**D e s c r i p t i o n :** The chela is characterised by the peculiar sloping ridges which cover the whole outer surface.

**R e m a r k s :** As mentioned in the chapter dealing with taxonomic questions (p. 48.), the identification of a Badenian species with a recent one is risky if the species has no record from intermediate times. *Dardanus arrosor* is the only species recorded from all stages between Badenian and today. In contrast to MILNE-EDWARDS' opinion no observable change occurred within this period.

**Dardanus substriatiformis (LÖRENTHEY in LÖRENTHEY—BEURLEN, 1929)**

Pl. XVII, figs. 1-4.

1929. *Pagurites substriatiformis* n. sp. — LÖRENTHEY in LÖRENTHEY—BEURLEN: 34, 73, pl. 3, fig. 3.

1953. *Pagurites substriatiformis* LÖRENTHEY — BEURLEN—BACHMAYER: 242.

1976a. “*Paguristes*” *substriatiformis* (LÖRENTHEY) — MÜLLER: 508.

1979a. *Dardanus substriatiformis* (LÖRENTHEY) — MÜLLER: 274.

M a t e r i a l seen: lectotype designated herein, ch. M24: FI.

M a t e r i a l not seen, probably lost: 3 ch. Rákos and Bia.

M a t e r i a l seen: 4 ch. MOE-9, ch. MR-8, 9 ch. MKC-7, ch. MER-3, 2 ch. MFA-4, ch. MNH-16: CA.

**D e s c r i p t i o n :** The outer surface of the palm and fixed finger is densely striated. On the inner side the outer striae are prolonged but they do not reach the median part. The palm is moderately stout, the upper and the lower edges are slightly convex. The striae are perpendicular to the longitudinal axis and they are finely denticulated. Every second ridge is stronger and longer than the intermediate ones.

**Dardanus hungaricus (LÖRENTHEY in LÖRENTHEY—BEURLEN, 1929)**

Pl. XVIII, figs. 1-5.

1929. *Pagurites hungaricus* n. sp. — LÖRENTHEY in LÖRENTHEY—BEURLEN: 34, 72, pl. 3, fig. 4.

?1945. *Dardanus lauensis* n. sp. — RATHBUN: 373, pl. 54, figs. c-h.

1976a. “*Paguristes*” *hungaricus* (LÖRENTHEY) — MÜLLER: 508.

1979a. *Dardanus hungaricus* (LÖRENTHEY) — MÜLLER: 274.

M a t e r i a l seen: lectotype designated herein, ch. without fingers M32: FI. Others, 3 ch. MK-3, ch. MTZ-18: CA. Ch. Mátraverebély-Szentkút, M61/2569: NMÖ.

**D e s c r i p t i o n :** Only the chelae are known. The palm is rather stout, convex transversely, its height slightly decreases in proximal direction. The lower part of the palm is adorned with 6 or 7 smooth striae, almost perpendicular to the long axis and are separated from each other by deep furrows. On the proximal part the striae are longer. The upper part is decorated with tubercles which gradually fuse downward into the striae, which are almost perpendicularly arranged to the longitudinal axis.

**R e m a r k s :** No recent species is similar to *D. hungaricus*, but RATHBUN described *D. lauensis* from the Pliocene of Fiji, which is identical with the Badenian species in all details visible on RATHBUN's figure. Unfortunately a request for casts of RATHBUN's specimens was refused by the Curator of the material in order to protect the fragile specimens.

**S t r a t i g r a p h y :** Lower to Upper Badenian, ?Pliocene.

Ecologically the species is probably bound to coral-reefs.

**Paguristes cserhatensis n. sp.**

Pl. XIX, figs. 1-4.

**M a t e r i a l seen:** holotype, ch. Sámsonháza, Halastópuszta, M61/3907: NMÖ.

**L o c u s t y p i c u s :** Sámsonháza, Halastópuszta, "bázisrétegek" (Noszky 1940).

**S t r a t u m t y p i c u m :** Lower Badenian.

**D e r i v a t i o n o m i n i s :** The type locality is situated in the Cserhát Mountains.

**D i a g n o s i s :** The left chela is decorated with rows of tubercles decreasing in size upwards.

**D e s c r i p t i o n :** The only known specimen, a left chela, without dactylus. The palm is very short, stout, its outer side is densely covered with relatively big tubercles arranged in rows slightly irregularly. The size of the tubercles decreases upwards while the distance between the individual tubercles and between the rows increases. The fixed finger is short and strong. Its cutting edge is triangular, with small teeth on both sides. The upper edge is covered by a group of calcite crystals.

**R e m a r k s :** The species is remarkably similar to *P. rosaceus* BARNARD (BARNARD 1950, p. 419.), but the tubercles are larger in the recent species.

**Genus *Petrochirus* STIMPSON, 1859**

**R e m a r k s :** The two recent species of this genus are notable for their unusual size. The Caribbean species, *P. diogenes* (L.) was reported from depth of 23-53 meters (HOLTHUIS 1959, p. 151), but several specimens were collected in a depth of 2 or 3 meters by L. CSILLING (pers. comm.).

***Petrochirus priscus* (BROCCHI, 1883)**

Pl. XIX, fig. 5, pl. XX, figs. 1-5, pl. XXI, figs. 1-3.

1883. *Pagurus priscus* n. sp. — BROCCHI: 7, pl. 5, fig. 9.  
1897. *Pagurus priscus* BROCCHI — LÖRENTHEY: 150, 161, 168.  
1898a. *Pagurus priscus* BROCCHI — LÖRENTHEY: 93, 106, 114.  
1898b. *Pagurus priscus* BROCCHI — LÖRENTHEY: 109, 134-136, pl. 9, figs. 6, 7.  
1898c. *Pagurus priscus* BROCCHI — LÖRENTHEY: 106-107, pl. 9, figs. 6, 7.  
1928. *Petrochirus priscus* (BROCCHI) — GLAESSNER: 173, 206, text-fig. 2.  
1929. *Pagurus priscus* BROCCHI em. LÖRENTHEY — LÖRENTHEY in LÖRENTHEY-BEURLEN: 34, 70, 71, pl. 3, figs. 1, 2.  
?1965. *Petrochirus cf. priscus* BROCCHI — VEIGA FERREIRA: 6-7, pl. 2, figs. 7-9, 11-12.  
1968. *Daira* sp. — STANCU-ANDREESCU: 466, pl. 7, fig. 84.  
1976a. "*Pagurus*" *priscus* BROCCHI — MÜLLER: 508.  
1979a. *Petrochirus priscus* (BROCCHI) — MÜLLER: 274.

**M a t e r i a l not seen,** probably lost: syntypes, 2? ch. Rákos: ?Sorbonne.

**M a t e r i a l not seen:** 64 ff., 3 da. Pöls, Baden-Rauchstallbrunngraben, Vöslau, Kalksburg, Devinska Nova Ves, Oued Tiflout (Atlas): NHMW.

Da. Rugi, Romania.

2 ch., 1 index, ? ca. Portugal: Coll. Serv. Geol. Port.

**M a t e r i a l seen:** numerous ch. Rákos: FI.

5 ch. Rákos, M60/5961 and M83/134: NMÖ.

6 ch. MR-2, 3 ch. MR8-7, 10 ch. MRF-3, 2 ch. MKC-10, 1 ch. MRO-7, 1 ch. MEG-3, 5 ch. MFA-5: CA.

**D e s c r i p t i o n :** The palm of the giant chelae is elliptical in cross-section, more or less rectangular in lateral view. The whole chela is densely covered with tubercles of subequal size. On the upper margin there is an irregular row of bigger tubercles or teeth. The both fingers are extremely robust. The cutting edges are different on the right vs. left chelae. On the right chela there are strong teeth on the fingers in one row at the distal end, diverging proximally. On the left side the dactylus bears also one row of teeth but these are smaller, while the fixed finger is flat, triangular, with two rows of small teeth on the edges.

**R e m a r k s :** The species is surprisingly similar to *P. diogenes* (LINNAEUS) in its form and in the arrangement of the teeth as well. The decoration on the surface is slightly different. The tubercles on the recent form are compound and bigger, while simple and smaller on *P. priscus*.

**E c o l o g y :** The species was bound to the presence of empty shells of giant *Gastropods*, e.g. *Galeodes*, *Strombus* or *Murex* in Badenian waters, the remnants of which are restricted to some layers.

**Superfamilia Galatheoidea SAMOUELLE, 1819**

**Familia Galatheidae SAMOUELLE, 1819**

**Subfamilia Galathinae SAMOUELLE, 1819**

**Genus Galathea FABRICIUS, 1793**

**Galathea weinfurteri BACHMAYER, 1950**

Pl. XXI, figs. 4, 5, pl. XXII, figs. 1-5.

- ?1928. *Galathea* sp. — GLAESSNER: 164, 206, pl. 3, fig. 1.  
1950. *Galathea weinfurteri* n. sp. — BACHMAYER: 135-137, pl. 1, figs. 2-4.  
1953. *Galathea weinfurteri* BACHMAYER — BACHMAYER: 242, 243, pl. 5, figs. 3, 4, 6.  
1974b. *Galathea weinfurteri* BACHMAYER — MÜLLER: 276, pl. 1, fig. 4.  
1976b. *Galathea weinfurteri* BACHMAYER — MÜLLER: 516.  
1979a. *Galathea* cf. *weinfurteri* BACHMAYER — MÜLLER: 274.

**M a t e r i a l** not seen: holotype, Furth: ?NHMW.

**M a t e r i a l** seen: 18 ca., 1 sternum, 1 merus, Deutsch-Altenburg: NHMW.

5 ca., 2 ch. MOE-33, 3 ca. MDG-4, 3 ca. MJL-5, 6 ca. MB-8, ca., 2 ch. MKC-9, ca. MEG-20, ca. MFK-5, 18 ca., 2 ch. MRZ-1, 24 ca., 2 ch. MV-2, 8 ca., 2 ch. MTZ-2: CA.

**D e s c r i p t i o n :** In spite of some earlier doubts on the unity of the taxon it seems that all specimens belong to BACHMAYER's species. The carapace is elongated, striated. Its rostrum is relatively bigger on smaller specimens but its form is similar to that of the bigger ones. Its base is triangular covered by some granulae. The long spines of the rostrum diverge anteriorly. Behind the rostrum there is a spiny ridge but the spines are very small or even rudimentary on some specimens. A specimen was found infected by *Bopyrids* (pl. XXII, fig. 5).

**R e m a r k s :** The species is close to *G. strigosa*, a recent form, but the postfrontal ridge is much less spinose on *G. weinfurteri*.

**S t r a t i g r a p h y :** Known from the whole time-span of the Badenian. In Messinian limestones of Kriti, Greece, a similar form occurs (GEORGIADES-DIKEOULIA-MÜLLER in press.).

**Familia Porcellanidae HAWORTH, 1825**

**Genus *Petrolisthes* STIMPSON, 1858**

**S t r a t i g r a p h y :** As far only one fossil species was recorded from Fiji (RATHBUN 1945). In the shape of its carapace it resembles some Lower Cretaceous *Galatheids*, e.g. *Galatheites troyoi* VAN STRAELEN.

**E c o l o g y :** *Petrolisthes* species invariably live in shallow waters, some of them are restricted to reefs. At least some *P.* species have a bioerosional activity on reefs.

***Petrolisthes magnus* n. sp.**

Pl. XXIII, figs. 1-4, pl. XXIV, figs. 1-4, pl. XXV, figs. 4-5.

**M a t e r i a l** seen: holotype, ca. MRZ-2-1, pl. XXIII, figs. 1., 2. Paratypes, 98 ca., 59 ch. MRZ-2, 2 ca., 2 ch. MKK-2: CA.

**L o c u s t y p i c u s :** Budapest, railway-cut of Rákos.

**S t r a t u m t y p i c u m :** Upper Badenian.

**D e r i v a t i o n o m i n i s :** Magnus, big (Lat.), because the species is one of the biggest *Porcellanid* known.

**D e s c r i p t i o n :** The carapace is subcircular in outline, but smaller specimens have subparallel lateral margins. The rostrum is triangular, with two faint incisions. The orbital margins are

rather elevated. The whole surface is striated, the small ridges are generally short but on the mesogastric field there are some longer ones. On the lateral margins there are sharp, entire keel-like ridges ending in one hepatic spine. The carpus is adorned on its upper edge with teeth separated from each other by deep incisions. On the chelae there are characteristic granulations on both sides tending to be arranged in rows. No median ridge occurs on the chelae.

**R e m a r k s :** ORTMANN's key (1897) led to the group of *P. galathinus* when it was used in the determination of this Badenian species. The only similar species within this group is *P. boscii* AUDOIN, even if one considers the newly described species (HAIG 1981). The specimens of *P. boscii* kindly sent me by GUINOT and DE SAINT-LAURENT, are indeed very close to the Badenian species. As no good figure of *P. boscii* was found in the literature some photographs are included here (pl. XXIII, fig. 5, pl. XXV, figs. 1-3.) for comparison. The carapax is essentially the same in both species, probably that of the recent one is slightly more narrow, and the orbital margins are less elevated. The differences are more accentuated on the cheliped. The elevations on the carpus are saw-like in the recent species, while the decoration on the inner side of the chela is much coarser than in *P. magnus*.

### **Petrolisthes haydni n. sp.**

Pl. XXVI, figs. 1-5.

**M a t e r i a l seen:** holotype, ca. 1984/40/7, pl. 26 fig. 1: NHMW.

Paratypes, 10 ca., 5 ch., 2 carpus, MNH-1, ca. Fertőrákos, drilling FRK-21 268 m, 2 ca. 2 ch. MTZ-16, ch. MZZ-1: CA.

**L o c u s t y p i c u s :** Gross-Höflein (Hung. Nagyhőflány), near Eisenstadt, Burgenland, Austria.

**S t r a t u m t y p i c u m :** Middle Badenian or lowermost Upper Badenian.

**D e r i v a t i o n o m i n i s :** Named after the great componist, JOSEPH HAYDN who worked in Eisenstadt.

**D e s c r i p t i o n :** The carapace is rounded square with a prominent triangular rostrum. The edge of the front is denticulated, its surface is finely granulated. The anterior part of the lateral margin is uninterrupted, forming a continuous carina, while the posterior half of this edge is cut into three individual spiniform teeth. On the postfrontal region there is a steplike transversal ridge cut into two parts by a median notch. The two parts of the ridge form an angle of about 150°. The surface of the carapace is finely striated with short lines which tend to unify behind the postfrontal ridge and at a narrow area near the anterior parts of the lateral margin. No *Bopyrid* infected specimen was found.

The carpus of the cheliped is serrated on its anterior edge. 5 or 6 teeth constitute the serration, the upper edges of the teeth form an angle of about 20° with the longitudinal axis. The outer surface of the palm is granulated by rounded granulae. About the median line the tubercles fuse into a ridge-like elevation which is parallel to the lower edge. The inner surface is also granulated. Near the median part the tubercles fuse into numerous oblique short ridges.

**R e m a r k s :** The species is closely related to *P. boscii* and to the other Badenian species. The presence of three teeth on the lateral margin is a diagnostic feature helping in the distinction. This species is much smaller than *P. magnus*. The serration on the carpus is different from that of the two other species. The ridge on the outer surface of the palm lacks on the other species.

**E c o l o g y :** The species was found in remnants of coral reefs, but one specimen occurred in Zebegény, in a layer full of colony-forming but ahermatypic (*Oculina?*) corals.

**S t r a t i g r a p h y :** The species is known in Lower and Middle (or lowermost Upper) Badenian layers.

### **Genus *Pisidia* LEACH, 1820**

**T y p u s - s p e c i e s :** *Pisidia linneana* LEACH = *Pisidia longicornis* (LINNAEUS, 1767).

### ***Pisidia kokayi* (MÜLLER, 1974)**

Pl. XXVII, figs. 1-5.

1974a. *Porcellana* (*Pisidia*) *kokayi* n. sp. — MÜLLER: 121, 124, 126, pl. 1, figs. 1-4.

1979a. *Porcellana kokayi* MÜLLER — MÜLLER: 274.

**M a t e r i a l seen:** holotype, ca. MB-2-1: NMÖ.

Paratypes: ca. MKF-1, 3 ca. MB-2: CA.

Others: 8 ca. MJL-18, ca. MB-2, 3 ca. MRZ-18, 3 ca. MDZ-1, ?ca. MZZ-2: CA.

**Description:** The outline of the carapace is a rounded square. The trilobated front is not very prominent. The cervical furrow is moderately deep. On the branchial regions the decoration consists of more or less long weak ridges while the other parts are covered with short ridges or tubercles. The lateral margin bears four teeth. The postfrontal ridges are rather long.

**Remarks:** The species is close to *P. longicornis* and *P. longimana* but is characterised by its shape being closer to a square. *P. blutely* is different by its prominent and wide frontal region.

**Pisidia cf. kokayi (MÜLLER, 1974)**

Pl. XXVII, fig. 6, pl. XXVIII, figs. 1, 2, ?3.

**Material seen:** 8 ca. 2 ch. MTZ-1, 18 ca. MV-3: CA.

**Remarks:** The form is similar to *P. kokayi* but its frontal part is more prominent, the cervical furrow is deeper and the postfrontal ridges are shorter and almost perpendicular to the longitudinal axis. Unfortunately well-preserved specimens were not found, thus a detailed description would be premature.

**Superfamilia Hippoidea LATREILLE, 1825**

**Familia Albuneidae STIMPSON, 1858**

**Genus *Albunea* WEBER, 1795**

**Ecology:** Similarly to *Raninidae Albunea* species dig themselves into different kinds of sediments, predominantly into sands. A striking grade of convergence occurs to *Raninids* owing to similar habits.

**Albunea asymmetrica (MÜLLER, 1979)**

Pl. XXVIII, figs. 4-6.

1979a. *Mioranina asymmetrica* n. gen. n. sp. — MÜLLER: 274, 278, 279, 287, pl. 9, 10.

**Material seen:** holotype, ca. MGY-2-1: CA.

Paratype, 8 ca., carpus, ch. MGY-2: CA.

Others, 5 ca. MOH-4, 4 ca., ch. MR8-16: CA.

**Description:** The carapace is subhexagonal in outline, convex in cross-section, nearly straight in lateral view. The fronto-orbital margin is very narrow, deeply notched. The margins outside of the orbits are ornamented with characteristic spines forming a comb-like feature. The lateral borders are nearly straight and parallel. The anterolateral borders converge posteriorly. The surface is decorated by step-like ridges forming a striking pattern. The decoration is slightly asymmetrical. The palm of the chela is triangular, decorated by ridges similar to those of the carapace. The carpus is covered by tubercles.

**Remarks:** Unfortunately the remnants were attributed to a new *Raninid* genus, *Mioranina*. The species, regarding its frontal decoration, is close to *A. steinitzi*, a recent form (SERÈNE 1965). The decoration of the recent form is composed of teeth of uniform size while at *A. asymmetrica* the size of the individual teeth varies, there is even a double tooth on it. *Albunea carabus* has less spines which are shorter than those of *A. asymmetrica*.

**Ecology:** The species is apparently confined to sediments composed of sands without many shell-debris or other coarse particles which could hinder burrowing. It is absent from limestones.

**Infraordo BRACHYURA LATREILLE, 1803**

**Sectio DROMIACEA DE HAAN, 1833**

**Superfamilia Dromioidea DE HAAN, 1833**

**Familia Dromiidae DE HAAN, 1833**

**Genus *Dromia* WEBER, 1795**

**E c o l o g y :** Most *D.* species are sublittoral, some were found in depth down to 150 meters (SAKAI 1976, p. 8).

**S t r a t i g r a p h y :** Some Eocene species attributed to *Dromilites* may belong here.

### **Dromia neogenica MÜLLER, 1979**

Pl. XXIX, figs. 1-6.

1979a. *Dromia neogenica* n. sp. — MÜLLER: 274, 278, 287, pl. 8, fig. 1.

**M a t e r i a l seen:** holotype, internal mould of ca. MKC-4-1: CA. Others, ch. MRZ-16, 5 ca., 2 ch. MDZ-3: CA. Ch. Souk el Khemis, Algeria, Orania, Messinian, leg. SAINT-MARTIN: coll. SAINT-MARTIN.

**D e s c r i p t i o n :** One of the biggest Badenian crabs. The carapace is strongly convex in both directions. The surface is smooth. The front consists of three short teeth, the median one is situated in a lower level. The orbits are of medium size. The supraorbital tooth is weak. The extraorbital tooth is well developed, situated on an upper level than the anterolateral teeth. The cervical groove is faint. The two extraorbital teeth situated before the last one are close to each other though with increasing age this distance increases as well.

The palm is rather long but its upper edge is short and it is adorned with faintly visible tubercles arranged in a row.

**R e m a r k s :** The species is similar to *D. personata* (LINNAEUS), but its supraocular teeth are rudimentary while in the recent species they are developed. At *D. personata* a well-developed cardiac furrow is present, while this furrow is visible only on the medium part of the carapace of the Badenian species. The arrangement of the anterolateral teeth and the shape of the chela are also different at the two forms.

**S t r a t i g r a p h y :** Known from the Lower Badenian to the Messinian.

**E c o l o g y :** All but one specimen were found in remnants of reefs.

### **Dromia eotvoesi (MÜLLER, 1976)**

Pl. XXIX, figs. 8, 9, pl. XXX, figs. 1-7.

1976a. *Dromilites eotvoesi* n. sp. — MÜLLER: 508, 509, 512, pl. 1, fig. 5, pl. 2, figs. 1, 4.

1976c. *Dromilites eotvoesi* MÜLLER — MÜLLER: 150, pl. 2, figs. 1, 2.

1979a. *Dromilites eotvoesi* MÜLLER — MÜLLER: 274.

**M a t e r i a l seen:** holotype, ca. MFA-1-1: CA.

Paratypes, ca. MA-4, ca. MKC-12: CA.

Others, 17 ca. MKC-12, 3 ca. MA-4, 7 ca., 2 ch. MR8-11, ch. MRC-2, ch. MRO-3, 7 ca. MEG-7, 2 ca. MOH-6, 12 ca., 2 ch. MGY-12, ca. Bia, unknown layer: CA.

Ca. Rákos, leg. STOCZEK, 3 ca. Kerepesi út, leg. MIHÁLY: FI.

**D e s c r i p t i o n :** The carapace is semiglobular. On the front there are two prominent spines on both sides. The extraorbital spine is directed below the orbit. The mesogastric region bears a rounded elevation, which is divided by a median notch. On the inner anterior part of the branchial field there are elevations on both sides. The other parts of the carapace are rather smooth. The branchiocardiac groove is shallow.

The palm of the chela is rather long, its upper edge is ornamented with four or five small teeth. The fixed finger is bent downwards. It bears 4 teeth. The distal end of the finger is spoon-like. The dactylus is strongly curved and longer than the index.

**R e m a r k s :** The species was originally placed in the genus *Dromilites* H. MILNE-EDWARDS. It is, however, closer to some recent *Dromia*-species than to *Dromilites bucklandi*, the type species of its genus, since there is no additional groove among the cervical and branchiocardical ones at the Badenian species. This feature might be of high diagnostic value. On the other hand, *D. eotvoesi* is close to *D. intermedia* LAURIE (SAKAI 1976, pl. 1, fig. 3.). The frontal teeth are much longer on the Hungarian species. The difference of the two Badenian species is striking.

**E c o l o g y :** The species is absent from reefal structures. Generally occurs together with *Calappa heberti*.

**S t r a t i g r a p h y :** Known only from the lower part of the Upper Badenian layers in Budapest.

**Typus-species:** *Dromilites koberi* BACHMAYER — TOLLMANN, 1953 by monotypy.

**Derivation nominis:** Phantasy name, a variation of *Dromilites* caused by a printing error. Used because it occurred in former literature as nomen nudum in STEININGER — PAPP (in PAPP et al. 1978, p. 199).

**Diagnosis:** The outline of the carapace is circular. The urogastric, mesogastric and epibranchial regions are adorned with elevations. The protogastric, hepatic, meso- und metabranchial regions are covered by small tubercles.

**Remarks:** The species and consequently the genus is rather different from any recent or fossil *Dromids*. *Dromilites lamarcki* is similarly strongly decorated but its overall form is different. Also, some recent *Petalomera* species, as *P. nodosa* SAKAI or *Cryptodromia areolata* have similar decoration but the outline of their carapace is dissimilar.

### **Kromtisis koberi (BACHMAYER — TOLLMANN, 1953)**

Pl. XXXI, figs. 1–4.

1953. *Dromilites koberi* n. sp. — BACHMAYER — TOLLMANN: 312, 313, pl. 1, figs. 2, 2a.

1976b. *Dromilites koberi* BACHMAYER — TOLLMANN — MÜLLER: 520.

1978. *Kromtisis koberi* BACHMAYER — TOLLMANN — STEININGER — PAPP in PAPP et al. 1978: 199.

**Material seen:** holotype, ca. Gross-Höflein, 61/1953: NHMW.

Others, 6 ca. MV-5, 3 ca. MNH-3: CA.

**Description:** The carapace is subcircular in outline. The anterolateral margin bears 6 teeth, one of which is situated on the posterolateral margin if properly designated. On the posterolateral margin there are two big tubercles near to the posterior margin. The surface is densely covered with bigger and smaller tubercles described in the diagnosis of the genus.

**Remarks:** As mentioned in the description of the genus, no similar *Dromid* species is known. Its place among *Dromids* is beyond any doubt, although some similarly decorated *Trachycarcinus* are also known.

**Ecology:** The species was found only in fossil coral-reefs.

**Stratigraphy:** Lower and Middle (or lowermost Upper) Badenian.

### **Genus Kerepesia MÜLLER, 1976**

**Typus-species:** *Kerepesia viai* MÜLLER, 1976 by monotypy.

**Remarks:** The genus was introduced as a Dromioidea incertae sedis. Actually it is a *Dromid*. Similar patterns of the anterolateral margin was observed at *Dromidiopsis excavata*, for instance. Its dromid affinities were confirmed by R. V. INGLE (pers. comm.), too.

### **Kerepesia viai MÜLLER, 1976**

Pl. XXXI, figs. 5–7, pl. XXXII, figs. 1–2, ? 3, 4.

1976c. *Kerepesia* n. gen. *viae* n. sp. — MÜLLER: 150, 151, 155, pl. 3, figs. 1–3., pl. 2, fig. 4.

1979a. *Kerepesia viai* MÜLLER — MÜLLER: 274.

**Material seen:** holotype, MKC-3-1: CA.

Paratypes, 3 ca. MKC-3: CA.

Others, ca. MKC-3, 2 ca. MDG-7, ca., ch. MOE-30: CA.

**Description:** Only the anterior parts of the carapace, including small regions of the branchial regions, are known. Probably other parts were less calcified. The anterior parts are moderately convex in both directions. The surface is smooth. The sides strongly converge in front of the last anterolateral tooth. Behind this tooth the sides are subparallel to each other. There are two frontal and two supraorbital spines. The anterolateral margin, consisting of four spiny teeth is hidden below the surface of the carapace, because above the genuine margin a false margin appears formed by the carapace that is strongly bended downwards in the plane of the orbits. Consequently the row of the anterolateral teeth is directed below the orbits.

On the locality MOE a typical *Dromid* chela was found. It is very much like the hand of *Cryptodromia*, *Petalomera*, *Hypoconcha*. All other Upper Badenian *Dromids* have different chelae. Consequently the illustrated hand can be attributed to *Kerepesia*.

**E c o l o g y :** The species was found in layers formed in infralitoral environments not very close to surface. The bottom was sandy with abundant shells. The species most probably used these shells as a shield much like its presumed recent relatives do.

**Dromiidarum n? gen. n. sp.**

Pl. XXXII, figs. 5-7.

M a t e r i a l seen: 2 ca. MTZ-23, ca. MNH-2: CA.

**R e m a r k s :** Only fragments of the median part and frontal region were found thus the species cannot be described. It does not belong to any of the Badenian crabs. The frontal region is typically *Dromid*-like, the presence of two deep parallel furrows (cervical and branchiocardioc) is an additional proof of the systematical position among *Dromioidea*.

**Familia Dynomenidae ORTMANN, 1892**

Genus *Dynomene* LATREILLE in DESMAREST, 1825

**R e m a r k s :** This frequent and widespread recent genus remained unrecognized in fossil form because of the similarity of its carapace to some *Xanthids*. Its front and the peculiar linear structure on the medium part of the cervical furrow are very characteristic features of this genus which is so important in phylogenetic speculations.

**Dynomene emiliae MÜLLER, 1979**

Pl. XXXIII, figs. 1-6.

- ?1945. *Xantho acutispinis* n. sp. — RATHBUN: 382, pl. 62, A.  
1979. *Pilumnus* sp. — FÖRSTER: 95, 103, text-fig. 7, pl. 2, fig. 2.  
1979b. *Dynomene emiliae* n. sp. — MÜLLER: 4, pl. 1, fig. 1.

M a t e r i a l seen: holotype, ca. Iasen, Bulgaria, leg. KOJUMDGIEVA, coll. University Sofia, Nt 531.

M a t e r i a l seen as cast: ca. Grobie, Poland, leg. WRONA, coll. University Warsaw.

M a t e r i a l seen: 3 ca. MV-4, 2 ca. MTZ-4, 3 ca. MNH-4: CA.

**D e s c r i p t i o n :** Only the carapace is known. It is moderately convex in both directions. The orbits are elongated, the upper orbital margins touch medially, forming a triangular, slightly rounded extension of the carapace in place of a front. There are five anterolateral teeth, since the outer orbital margin does not form any tooth or spine. The mesogastric region is well delimited posteriorly by a V shaped cervical groove. Immediately behind the cervical groove there is a rudiment of the postcervical groove. On partly decorticated specimens the carapace is finely granulated but the upper layer is smooth with fine pores.

**R e m a r k s :** The species is similar to *D. pilumnoides* though it is wider than the recent form. It is also close to *D. hispida* but the surface of the miocene species is smooth and the recent one displays only some indistinct tubercles. The posterior margin is wider at *D. emiliae* than on its recent relatives. The most similar form is *Dynomene acutispinis* (RATHBUN) (= *Xantho acutispinis*) from the Pliocene of Fiji. Unfortunately a direct comparison was impossible just as mentioned for *Dardanus hungaricus*.

**E c o l o g y :** All specimens but the Iasen one were found in coral-reefs, but in Iasen, corals occur in the close vicinity of the locality.

**Sectio OXYSTOMATA H. MILNE-EDWARDS, 1834**

Superfamilia *Dorippoidae* DE HAAN, 1841

Familia *Dorippidae* DE HAAN, 1841

Subfamilia *Dorippinae* DE HAAN, 1841

Genus *Dorippe* WEBER, 1795

**Dorippe margaretha LÖRENTHEY in LÖRENTHEY—BEURLEN, 1929**

Pl. XXXIV, figs. 1–5.

1911. *Dorippe margaretha* nom. nud. — LÖRENTHEY: 528.  
 1913. *Dorippe margaretha* nom., nud. — LÖRENTHEY: 326.  
 1929. *Dorippe margaretha* n. sp. — LÖRENTHEY in LÖRENTHEY—BEURLEN: 34, 135, 136, pl. 6, fig. 1, a rather inaccurate drawing.  
 ?1965. *Dorippe aff. lanata* LINNAEUS — VEIGA FERREIRA: 10, pl. 1, figs. 5, 6.  
 1979a. *Dorippe margaretha* LÖRENTHEY — MÜLLER: 279, 287, pl. 13, fig. 4.

**M a t e r i a l seen:** holotype, ca. (internal mould), Rákos, M25: FI.  
 Others, 2 ca. MEG-12, 12 ca. MOE-22, ca. MDG-14: CA.

**D e s c r i p t i o n :** Only the carapace is known. It is rather wide, pyriform. The surface is divided by deep furrows. There are spiniform teeth on the branchial and on the hepatic border (broken down on the illustrated specimens). The rather smooth branchial region is wide and is decorated with a tubercle. On some specimens three faint ridges leave this tubercle. On the cardiac region there are two ridges diverging anteriorly, forming an angle of about 50–60°.

**R e m a r k s :** The species is close to *Dorippe lanata*. The recent form is slightly wider than the Badenian one. Consequently the mentioned ridges on the cardiac region form a rectangle at *D. lanata*. Also the decoration on the branchial region is stronger, but rather variable. The Indo-West-Pacific species *D. frascone* is unlike *D. margaretha* at all.

**?Dorippe carpathica (FÖRSTER, 1969)**

- 1969a. *Dorippe? carpathica* n. sp. — FÖRSTER: 91–93, text. fig. 3, pl. 2, fig. 3.

**M a t e r i a l not seen:** holotype, ca. Niskowa, Poland; coll. Univ. Warsaw.

**D e s c r i p t i o n :** According to FÖRSTER, the carapace is subquadrangular. The posterior margin is bilobated. The regions are well-defined. On the meso/protogastric region there are three protuberances. Epibranchial and hepatic regions are modified by longitudinal grooves and depressions, and are separated from meso- and metabranchial regions by a groove. There is a prominent ridge on meso/metabranchial region, which is subparallel to the groove.

**R e m a r k s :** The species reveals similarities to *Dorippids*, but, as FÖRSTER pointed out, the distinct grooves are absent on the epibranchial and hepatic regions of any *Dorippid*. Similar grooves are present on some *Cymonomus* (and even at some *Tymolus*) species though there are obvious differences between the mentioned recent and Badenian forms.

**Superfamilia Calappoidea DE HAAN, 1833**

**Familia Calappidae DE HAAN, 1833**

**Subfamilia Calappinae DE HAAN, 1833**

**Genus Calappa WEBER, 1795**

**Calappa praelata LÖRENTHEY in LÖRENTHEY—BEURLEN, 1929**

Pl. XXXV, figs. 1, 2, 7, ?figs. 3–6, ?pl. XXXVI, fig. 6.

- partim? 1928. *Calappa heberti* BROCCHEI — GLAESNER: 174–175.  
 1929. *Calappa praelata* n. sp. — LÖRENTHEY in LÖRENTHEY—BEURLEN: 132, 133, pl. 6, fig. 3.  
 ?1958. *Calappa heberti* BROCCHEI — VEIGA FERREIRA: 203–207, text-fig. 1–3, pl. 1, figs. 1–3.  
 ?1959. *Calappa cf. heberti* BROCCHEI — GALOPIM DA CARVALHO: 79, pl. 2, figs. 4–6.  
 1961. *Calappa heberti* sp. 2. — BACHMAYER: 41, pl. 2, fig. 1.  
 ?1961. *Calappa aff. granulata* LINNÉ — BACHMAYER: 42.  
 1979b. *Calappa aff. heberti* BROCCHEI — FÖRSTER: 255, 257, pl. 1, figs. 2, 4.

**M a t e r i a l seen:** holotype, ca. Mátraverebély-Szentkút, Szent László-forrás, leg. Noszky, M27: FI.

Others, da. Korytnica, Poland, leg. BALUK, 3 da. MSZ-2, ch. Szob-Kérkhegy, ch., ca. MTH-2, ca. MVA-1, ca. MNU-1, 3 da. Oslip, sand-pit, ca. Mátraverebély-Szentkút, Meszestető: CA.

**D e s c r i p t i o n :** The carapace is strongly convex, subpentagonal in outline. The median regions (mesogastric, urogastric, cardiac) are markedly delimited from other regions by deep longi-

tudinal furrows. This longitudinal elevation is ornamented with 5 or 6 bigger and numerous smaller tubercles. On the anterior part of the branchial region an area is covered with 7-8 big, densely packed tubercles. On other parts of the carapace there are numerous regularly arranged small tubercles. The posterolateral angle is decorated with angular lobes, each of which are medially carinated by ridges. The widest part of the carapace is characteristically situated at its anterior one-third, but this feature is not as striking as it appears on LÖRENTHEY's idealized figure.

The chela is known as fragments. The dactylus is typical *Calappa*-like, the only diagnostic feature on it is the proximal lobe situated on its almost symmetrical upper margin.

**R e m a r k s :** The species is highly similar to *C. granulata* LINNAEUS. Since few illustration exist in the literature showing the sculpture, photographs are presented to show the carapace, chela and right dactylus (leg. P. SOLT, Sicily) pl. XXXVI, figs. 1-5. The similarity is indeed striking. The anterolateral lobe is more extended, however, in *C. granulata*, consequently it bears 6 big lobes instead of 4. The proximal lobe on the right dactylus is bent in proximal direction. As it can be observed on the figures, *C. heberti* differs from the other species.

**S t r a t i g r a p h y :** The few data available suggest that the species does not occur in Upper Badenian layers. The species lived longer in the Mediterranean where it was perhaps the ancestor of *C. granulata*.

**E c o l o g y :** In contrast to *C. heberti*, this species has not been found in reefs yet.

### ***Calappa heberti* BROCCHI, 1883**

Pl. XXXVII, figs. 1-7, pl. XXXVIII, figs. 1-6.

1883. *Calappa heberti* n. sp. — BROCCHI: 2, pl. 4, fig. 3.  
1897. *Calappa heberti* BROCCHI — LÖRENTHEY: 150, 159, 166.  
1898a. *Calappa heberti* BROCCHI — LÖRENTHEY: 93, 103, 104, 112.  
1898b. *Calappa heberti* BROCCHI — LÖRENTHEY: 109, 113-116, 151, pl. 8, figs. 1-3, pl. 9, fig. 1.  
1898c. *Calappa heberti* BROCCHI — LÖRENTHEY: 88-90, pl. 8, figs. 1-3, pl. 9, fig. 1.  
1911. *Calappa heberti* BROCCHI — LÖRENTHEY: 523.  
1913. *Calappa heberti* BROCCHI — LÖRENTHEY: 318.  
partim? 1928. *Calappa heberti* BROCCHI — GLAESSNER: 174-175.  
1929. *Calappa heberti* BROCCHI — LÖRENTHEY in LÖRENTHEY-BEURLEN: 34, 130-132, pl. 6, figs. 2, 9-12.  
1976a. *Calappa heberti* BROCCHI — MÜLLER: 509.  
1976c. *Calappa heberti* BROCCHI — MÜLLER: 152.  
1979a. *Calappa heberti* BROCCHI — MÜLLER: 274.

**M a t e r i a l** not seen, probably lost: syntypes, ca., ch. Rákos: ?Sorbonne.

**M a t e r i a l** seen: about 100 ca., ch., merus, carpus, Rákos: FI.

4 ca. M59/2369, 2 ca. M61/5790-91, Rákos: NMÖ. 8 ca. MR8-12, 4 ca. MR45-4, 3 ca., 10 ch. MR-10, 3 ca. MRF-5, 38 ca. MKC-14, 4 ca. MOH-7, 5 ca. MGY-14, 2 ca., ch. MRL-3, 2 ca. MEG-19, 4 ca. MFA-6, 3 ca., 5 ch. MFK-4, 10 ca. MA-6, ca. MRZ-3, 2 ca., da. MER-5, ca. MDZ-5, ca. MH34: CA.

**D e s c r i p t i o n :** The subpentagonal carapace is convex in both directions. A longitudinal median region is delimited by two deep furrows. The anterior part of the carapace is adorned with tubercles, the posterior with short ridges. There are transitional forms of these decorations in intermediate areas. Near the posterolateral angle the margin is composed of lobes which are not very prominent. The chelae are characteristically decorated with rows and groups of tubercles. The dactylus bears a proximal lobe directed distally.

**R e m a r k s :** The species is highly characteristic and differs from all living species of *Calappa*. *C. hepatica* (LINNAEUS) and *C. gallus* (HERBST) have similar ornaments on their carapace but the outline of them is unlike that of *C. heberti*. It can be hardly understood why this characteristic species was so often reported from the Neogene of the Mediterranean since all figures published are dissimilar from the Badenian form (cf. literature given at *C. praelata*).

**E c o l o g y :** The species was found in various types of sediments, fine and coarse calcarenites, conglomerates, tuffites, further in remnants of Coral-reefs.

**S t r a t i g r a p h y :** The species seems to be confined to the lower level of Upper Badenian layers.

Genus *Calappilia* A. MILNE-EDWARDS 1873 (= *Paracyclois* Miers 1866)

**Calappilia matzkei (BACHMAYER, 1961)**

1961. *Calappa matzkei* n. sp. — BACHMAYER: 43–45, text-fig. 1, pl. 3, fig. 1.

1979c. *Calappilia matzkei* (BACHMAYER) — MÜLLER: 866.

M a t e r i a l not seen: holotype, ca. (internal mould), Brunn, Austria: coll. ?Matzke\*.

D e s c r i p t i o n : According to BACHMAYER, the internal mould of the only known carapace is wider than long, its greatest width is in the middle. The lateral margins are adorned with numerous spines, or spiniform tubercles. The outline of the carapace is ovoid. On the hinder part of the anterolateral border there were spines which were broken off.

R e m a r k s : According to BACHMAYER the species is similar to *Calappa praelata* LÖRENTHEY. The general form of the carapace, its decoration and the spines on the anterolateral border all reveal close similarities to *Calappilia dacica* BITTNER, however.

S t r a t i g r a p h y : The presence of a *Calappilia*-species in the Badenian stage is not surprising since, according to GLAESNER (1969, p. R494), the still living Indo-West-Pacific *Paracyclois milneedwardsi* MIERS should be ranged among *Calappilia*-species.

**Genus *Mursia* DESMAREST, 1823**

***Mursia lienharti* (BACHMAYER, 1961)**

Pl. XXXIX, figs. 1–6, pl. XL, figs. 1–3, 5–6.

1961. (?)*Calappa lienharti* n. sp. — BACHMAYER: 41–42, pl. 2, fig. 2.

?1964. *Calappa* sp. — GRIPP: 55, text-fig. 1.

1972. *Calappa lienharti* BACHMAYER — JANSSEN: 93, text-fig. 39.

1974b. *Parthenope loczyi* n. sp. — MÜLLER: 277, 281, 283, pl. 1, fig. 7.

1976b. *Mursia loczyi* (MÜLLER) — MÜLLER: 149, 151, 152, 155, pl. 4, figs. 3–4, pl. 5, figs. 1–4.

1979a. *Mursia loczyi* (MÜLLER) — MÜLLER: 274.

In press *Mursia lienharti* (BACHMAYER) — JANSSEN—MÜLLER.

M a t e r i a l not seen: holotype, da. Nussdorf, Austria 456/1962: NHMW.

M a t e r i a l seen: holotype of *Parthenope loczyi*, ca. MKC-1-1: NMÖ.

Others, da. MOE-33, 2 ca., 5 ch., 4 da. MR8-20, ca. MRF-7, 17 ca., 2 ch. MKC-1, ca. MGY-17, 2 ca., 2 ch. MFK-2, 3 ca. MNA-5, ca. MN-5: CA.

D e s c r i p t i o n : The anterior part of the carapace is circular in outline, the posterior one converges posteriorly. The front is prominent, trilobated. The orbits are deep, their upper margin is cut by two notches. The anterolateral margin is adorned with 9 to 10 blunt lobes. The row of these lobes is closed posteriorly by a lateral spine of moderate size which is directed slightly backwards. The posterolateral margin is slightly concave. The posterior margin is convex, delimited by two corners which are prominent, tooth-like in large specimens. The whole surface is decorated by tubercles. On the branchial and hepatic regions there are two rows of big tubercles on both sides. The inner rows are subparallel to the longitudinal axis, while the outer ones diverge posteriorly.

The palm is flattened, rather high. The outer surface is adorned with rows of tubercles.

The dactylus described by BACHMAYER as *Calappa lienharti* is a typical *Mursia* member, but the stridulating ridge is absent on the inner side. In its general form and adornment it is like *Mursia armata* (pl. XL, fig. 4.). As the dactylus and other remnants were found in the same group of layers, in the absence of other candidate species, very likely they belong to the same species.

R e m a r k s : The Badenian species is close to *M. hawaiiensis*, but posterolateral margins bear no teeth at *M. lienharti*. Recently beautiful specimens undoubtedly of the same species were collected in Ramsel, Belgium (JANSSEN—MÜLLER in press.).

S t r a t i g r a p h y : Though the species was found only in Upper Badenian layers in Hungary, the Ramsel locality is Hemmoorian in age (JANSSEN personal comm.), thus it is older than Lower Badenian.

**Subfamilia *Matutinae* MCLEAY, 1838**

**Genus *Matuta* FABRICIUS, 1798**

\* Holotype kept in the Heimatmuseum, Neunkirchen (Niederösterreich. BACHMAYER, pers. comm.).

**Matuta brocchii GLAESSNER, 1969**

Pl. XLI, figs. 1–6., pl. XLII, figs. 1–5.

- non 1882. *Matuta inermis* n. sp. — MIERS: 256, pl. 26, fig. C.  
 1883. *Matuta inermis* n. sp. — BROCCHI: 5, pl. 4, fig. 1.  
 1897. *Matuta inermis* BROCCHI — LÖRENTHEY: 150, 159, 166.  
 1898a. *Matuta inermis* BROCCHI — LÖRENTHEY: 93, 104, 112.  
 1898b. *Matuta inermis* BROCCHI — LÖRENTHEY: 109, 116, 152.  
 1898c. *Matuta inermis* BROCCHI — LÖRENTHEY: 91.  
 1904a. *Matuta inermis* BROCCHI — LÖRENTHEY: 161.  
 1904b. *Matuta inermis* BROCCHI — LÖRENTHEY: 30.  
 1913. *Matuta inermis* BROCCHI — LÖRENTHEY: 322.  
 1929. *Matuta inermis* BROCCHI — LÖRENTHEY in LÖRENTHEY — BEURLEN: 34, 134, 135, pl. 6, figs. 14–18.  
 1969. *Matuta brocchii* nom. nov. — GLAESSNER: R495.  
 1979a. *Matuta brocchii* GLAESSNER — MÜLLER: 274.

Material not seen, probably lost: syntypes or holotype, ca. Rákos: ?Sorbonne.

Material seen: others, about 100 ca. Rákos, Budapest: FI. 2 ch. M83/136, ca. M83/125, Rákos: NMÖ. 2 ca. MOE-23, ca. MGF-3, ca. MB-10, 1 ca., 5 ch. MR-11, 8 ca., 10 ch. MR8-15, ca., 2 ch. MR45-2, 3 ca., 3 ch. MRF-6, 132 ca. MKC-13, 2 ca. MRC-4, 5 ca. MRO-4, 16 ca., ch. MEG-8, 9 ch. MOH-8, 6 ca., ch. MER4, numerous ca., ch. MGY-13, ca. MFA-10, 4 ca. MFK-3, ch. MA-8, ca., ch. MRZ-15, ca. BAK-40: CA.

**Description:** The carapace is subhexagonal, rather smooth. The epigastric, mesogastric, urogastric and branchial regions are decorated with big tubercles, the relative size of which decreases with increasing size of the animal. The palm is decorated by big tubercles. On the inner side of it there are stridulating ridges.

**Remarks:** The only known fossil species of the genus. It is remarkably similar to *M. inermis* MIERS, but its carapace is wider (although narrow carapace occurs also in Badenian material) and the denticulation on the anterior 1/3 of the anterolateral border is finer in *M. brocchii*. *Matuta curtispina* SAKAI is also similar but its anterolateral denticulation is different.

**Familia Leucosiidae SAMOUELLE, 1819**

**Genus *Ebalia* LEACH, 1817**

***Ebalia vanstraeleni* BACHMAYER, 1953**

Pl. XLIII, figs. 1, 2.

1953. *Ebalia van-straeleni* n. sp. — BACHMAYER: 243, 244, pl. 2, figs. 1, 4, 4a.  
 1979a. *Ebalia vanstraeleni* BACHMAYER — MÜLLER: 279, 288.

Material seen: holotype, ca. (internal mould), Deutsch-Altenburg: NHMW

Material not seen: paratypes, 2 ca. (internal moulds), Deutsch-Altenburg: NHMW.

**Description:** The carapace (internal moulds) is flat, subhexagonal. According to BACHMAYER, the median regions are well developed. The mesogastric region bears two elevations. The urogastric region is adorned with two small tubercles. On the metagastric region there is a strong tubercle, while the branchial region bears two of them.

**Remarks:** The species is similar to some European *E.* species. Probably *E. granulosa* is the closest form but the strong ornament of the branchial region and the extended hepatic region are distinctive at *E. vanstraeleni*.

***Ebalia oersi* MÜLLER, 1979**

Pl. XLIII, figs. 3–7. ?fig. 8.

- 1979a. *Ebalia oersi* n. sp. — MÜLLER: 279, 288, pl. 13, figs. 1–3.

Material seen: holotype, ca. MOE-4-1: CA.

Paratypes: 2 ca. MOE-4: CA.

Others: 18 ca. MOE-4; ?ca. drilling "Váci út 120": CA.

**Description:** The carapace is flat, shield-like, subpentagonal. The whole surface is covered with granulation which is rather even but in the median parts slightly coarser. The lateral and posterior borders have sharp edges. The median regions are elevated, especially the cardiac one, which

is round. On the branchial region there is one, not striking, smooth elevation. The posterior margin bears two lobes but at some specimens the margin is rather straight.

**R e m a r k s :** The species is most probably closest to *E. edwardsi* COSTA, but is much less wide. From *E. vanstraeleni* it differs by the weaker decoration and by the less extended hepatic regions.

**Ebalia sp. div., nov.?**

Pl. XLIV, figs. 1-4.

**M a t e r i a l seen:** ca. MTM-1, 2 different ca. MZZ-2 and MZZ-3: CA.

The specimens are rather incomplete or poorly preserved. They belong to three different species and they differ from any other Badenian species, though the Törökmező specimen is close to *E. oersi*.

**‘Ebalia’ hungarica MÜLLER, 1974**

Pl. XLIV, fig. 5, pl. XLV, figs. 1-5.

1974a. *Ebalia hungarica* n. sp. — MÜLLER: 121, 126, pl. 1, figs. 5-7.

1974b. *Ebalia hungarica* MÜLLER — MÜLLER: 276.

1979a. *Ebalia hungarica* MÜLLER — MÜLLER: 274.

**M a t e r i a l seen:** holotype, ca. MB-3-1: NMÖ.

Paratypes, 2 ca. MB-3: CA.

Others, 10 ca. MDG-12, 14 ca. MJL-11, 3 ca. MBH-5, 6 ca. MB-3, ca. MUO-1, ca. MBK-2, 2 ca. MKC-24, 2 ca. MRO-6, ca. MR8-18: CA.

**D e s c r i p t i o n :** The carapace is close to an elongated hexagon, convex laterally and longitudinally. The whole surface is densely tuberculated. The median regions (gastric and cardiac) are elevated. All regions are well delimited by depressions. There are some bigger tubercles near the lateral angle. The posterior margin is bilobated, but the lobes fuse in some specimens. All transitions exist between the fused and bilobated forms.

**R e m a r k s :** Though the species was originally compared with *E. tuberculosa* H. MILNE-EDWARDS, it is highly peculiar and no similar form occurs either in the fossil or in the zoological record. Most probably the species belongs to a new genus.

**Genus *Andorina* LÖRENTHEY, 1901**

***Andorina elegans* LÖRENTHEY, 1901**

Pl. XLVI, figs. 1-4.

1901a. *Andorina elegans* n. gen. n. sp. — LÖRENTHEY: 833-835, pl. 1, fig. 1.

non 1901a. *Andorina?* *elegans?* n. sp. — LÖRENTHEY: pl. 1, fig. 2. (Parthenope!)

1901b. *Andorina elegans* LÖRENTHEY — LÖRENTHEY: 330-332, pl. 1, fig. 1.

non 1901b. *Andorina?* *elegans?* — LÖRENTHEY: pl. 1, fig. 2.

1929. *Andorina elegans* LÖRENTHEY — LÖRENTHEY in LÖRENTHEY-BEURLEN: 34, 139-140, pl. 8, fig. 2.  
Non! fig. 3.

**M a t e r i a l seen:** holotype, ca. Telepy utca, Budapest, M30: FI.

**D e s c r i p t i o n :** The carapax is elongated, elliptical in its outline, since posterior margin is absent. The surface is densely adorned with tubercles predominantly of two different sizes. On the postfrontal and hepatic regions only the tubercles of smaller size are present. The hepatic region is rather extended. The cardiac region is elevated.

**R e m a r k s :** Though described as an *Oxyrhynch*, the species is a typical leucosid (MÜLLER 1979a, p. 279, 287). No similar form was found in the zoological literature, however.

**Genus *Palaeomyra* A. MILNE-EDWARDS 1861**

1979a. *Ebalites* n. gen., typus-species *Ebalia globulosa* MÜLLER — MÜLLER: 279, 287.

**'Palaeomyra' globulosa (MÜLLER, 1976)**

Pl. XLVII, figs. 1-7.

1976a. *Ebalia globulosa* n. sp. — MÜLLER: 509, 510, 512, pl. 2, figs. 2, 3, 5.

1979a. *Ebalites globulosus* (MÜLLER) — MÜLLER: 279, 287, pl. 14.

M a t e r i a l seen : holotype, ca. MA-1-1: CA.

Others, 27 ca., ch. MOE-14, ca. MRF-14, ca. MNU-2: CA.

**D e s c r i p t i o n :** The carapace is elongated, subglobose, all but the cardiac regions are ill-delimited. The whole surface is granulated except the smooth postfrontal region. The lateral margins are completely circular, but a slight variation occurs in their curvature. There are two characteristic spiniform extensions on the posterior margin which are either flattened or rather narrow on different individuals. The chela found in the same layer where numerous carapaces occurred, is elongated. The slender fixed finger is directed downwards.

**R e m a r k s :** The species is similar to *Palaeomyra bispinosa* A. MILNE-EDWARDS, but the median ridge is absent from the Badenian form. The posterior spines are also different, much wider in *P. globulosa* than on the figure of MILNE-EDWARDS. Many old drawings are idealised, and the Italian specimen was an internal mould. Unless its type will not be found, the relationship of the two species remains doubtful.

**Superfamilia Raninoidae DE HAAN, 1841**

**Familia Raninidae DE HAAN, 1841**

**Genus Ranidina BITTNER, 1893**

**Ranidina rosaliae BITTNER, 1893\***

Pl. XLVI, fig. 6.

1893. *Ranidina* n. gen. *rosaliae* n. sp. — BITTNER: 33-46, pl. 2, fig. 2.

1929. *Ranidina rosaliae* BITTNER — LÖRENTHEY in LÖRENTHEY-BEURLEN: 121-122, pl. 5, fig. 8.

1975. *Ranidina rosaliae* BITTNER — STOJASPALE: A190.

1979b. *Ranidina rosaliae* BITTNER — MÜLLER: 4-5, pl. 3, fig. 2.

M a t e r i a l not seen: syntypes, 5 ca. Walbersdorf: Geolog. Bundesanstalt, Wien.

M a t e r i a l seen: ca. Gorna Mitropolia, Bulgaria, leg. KOJUMDGIEVA: coll. Univ. Sofia.

**D e s c r i p t i o n :** The carapace is elongate, smooth, hexagonal in outline. It is convex transversally while less so longitudinally. There are two lateral spines, two extraorbital and one frontal spine on the carapace. The two parallel ridges present on BITTNER's reconstruction were not observable on the Bulgarian specimen. Anterolateral margins are convex while the posterolateral ones are straight. The upper margin of the chela is spinose.

**R e m a r k s :** The position of the species within the recent *Raninids* remains doubtful until better specimens will not be found. Its independence is likely, however.

**Sectio OXYRHYNCHA LATREILLE, 1803**

**Familia Majidae SAMOUELLE, 1819**

**Subfamilia M a j i n a e SAMOUELLE, 1819**

**Genus Maja LAMARCK, 1801**

**Maja biaensis LÖRENTHEY in LÖRENTHEY-BEURLEN, 1929**

Pl. XLVIII, figs. 1-6, pl. XLIX, figs. 1-3.

?1854. *Maja orbigniana* nom. nud. — MILLET: 152.

?1865. *Maja orbigniana* n. sp. — MILLET: 1.

?1873. *Maja* sp. — WOODWARD: 325.

\* See Addendum.

- ?1907. *Maja miocenica* n. sp. — LÖRENTHEY: 237.  
 ?1908. *Maja orbigniana* MILLET — COUFFON: 2, 3, pl. 1, figs. 1–4.  
 ?1910. *Maja orbigniana* MILLET — COUFFON: 130, fig. 4.  
 1929. *Maja biaensis* n. sp. — LÖRENTHEY IN LÖRENTHEY — BEURLEN: 34, 148–150, pl. 7, fig. 1. (idealised!)  
 1976a. *Maja biaensis* LÖRENTHEY — MÜLLER: 510, pl. 3, fig. 2.  
 1979a. *Maja biaensis* LÖRENTHEY — MÜLLER: 274.
- M a t e r i a l** seen: plaster cast of holotype, ca. Bia: FI.  
 Others: ca. (fragment) WEG, (leg. KRACH), ca. (fragment) MNU-3, 1 ca. MNA-1, 2 ca., ch., carpus MFA-7, ca. MKC-23, 2 ca., ch. MOE-20, 6 ca. MOH-9, 3 ca. MGY-23, 2 ca. MN-1: CA.

**D e s c r i p t i o n :** The carapace is elongated, pyriform, strongly tuberculated and spinose. The frontal spines are particularly long and the lateral ones are more slender and sharp. On the median line there are some big tubercles, while on the other parts there are groups of bigger or smaller tubercles.

**R e m a r k s :** *Maja orbigniana* MILLET is very similar to our form. The only difference is that the tubercles are slightly bigger and closer to each other on the French form. Whether this is due to individual variability cannot be decided at moment. Also, *M. miocaenica* LÖRENTHEY is similar to the Budapest specimens. WOODWARD's Maltese specimen is still preserved in the British Museum of Natural History (pers. comm. of DR. COLLINS) though LÖRENTHEY's Sardinian specimen was most probably lost. Probably all the Miocene *Maja* species of this restricted area were closely related or even identical with each other. Comparing the Miocene species with *M. squinado*, it can be stated that the median spines are bigger in the Miocene form. Seen from the side *M. squinado* has an elliptical outline, while *M. biaensis* (and *M. orbigniana* also) is depressed between the meso- and urogastric fields.

#### Genus *Schizophrys* WHITE, 1848

##### *Schizophrys visegradensis* n. sp.

Pl. XLIX, figs. 4, 5, pl. L, figs. 1–3.

1976b. *Libinia?* sp. — MÜLLER: 516.

**M a t e r i a l** seen: holotype, ca. MTZ-20-1: CA.  
 Paratypes, 2 ca. MTZ-20, 10 ca. MV-8: CA.

**L o c u s t y p i c u s :** Nagymáros, Törökmező.

**S t r a t u m t y p i c u m :** Lower Badenian.

**D e r i v a t i o n n o m i n i s :** From the first known locality, Visegrád.

**D i a g n o s i s :** Rather flat, pyriform carapace with short frontal spines.

**D e s c r i p t i o n :** The pyriform carapace is rather flat. The decoration on the surface is sparse. The cardiac region is well delimited. The cervical groove is moderately deep. The two simple frontal spines are rather short. On the anterolateral margin there are spines. The two well preserved are double. The other spines were short and simple. On the mesobranchial region there are five tubercles in a transverse row. On the branchial region there are also tubercles, their total number is about 15 on each side. Between the tubercles the surface is smooth.

**R e m a r k s :** The species is close to *Schizophrys aspera*, recent Indo-West-Pacific species. The anterolateral teeth are simple in the recent form, while the frontal ones are compound and longer than that of the Miocene form.

#### Subfamilia *Inachinae* MCLEAY, 1838

#### Genus *Achaeus* LEACH, 1815

##### *Achaeus magnus* MÜLLER, 1979

Pl. L, figs. 4–6, pl. LI, figs. 1, 2.

1979a. *Achaeus magnus* n. sp. — MÜLLER: 274, 279, 280, 288, pl. 15.

**M a t e r i a l** seen: holotype, ca. MOE-5-1: CA.  
 Paratypes: 2 ca., 2 ch. MOE-5: CA.  
 Others, 12 ca. MOE-5: CA.

**Description:** The outline of the carapace is triangular, it is convex transversely. There are two short rostral spines with slightly upwards directed tips. Behind the orbits there is a neck-like narrow part. The height of the carapace decreases gradually posteriorly. On the mesogastric region there is an erect spine. The mesogastric and cardiac regions are well delimited laterally by deep furrows. On the cardiac region there are two smooth, rounded protuberances near each other.

**Remarks:** The palm of the chela is elongated but rather convex. The index is long, narrow, inward directed. The chela is similar to that of the modern *Achaeus* species. Other *Majids* of the locality have different chelae.

**Remarks:** The Mediterranean-East-Atlantic species of *Achaeus* are quite different, generally smaller. Among the Indo-West-Pacific forms *A. tuberculatus* has similar erect spines, but two instead of one.

#### Subfamilia *Pisinae* DANA, 1852

##### Genus *Pisa* LEACH, 1814

###### '*Pisa*' *oroszyi* (BACHMAYER, 1953)

Pl. LI, figs. 3–6, pl. LII, figs. 1–3.

1953. *Maia oroszyi* n. sp. — BACHMAYER: 245–247, pl. 2, fig. 3.

1974a. *Pisa oroszyi* (BACHMAYER) — MÜLLER: 122, 124, pl. 2, figs. 1–3.

1979a. *Pisa oroszyi* (BACHMAYER) — MÜLLER: 274, 274, 275.

**Material seen:** holotype, ca. Deutsch-Altenburg, leg. K. OROSZY: NHMW.

Others: 21 ca. MM-1, 46 ca., ch. MB-12, 2 ca. MDG-20, 8 ca. MGF-2, ca. MRZ-17, ca. MOE-13, 2 ca. MKF-2, 2 ca. MRS-3, 45 ca. MJL-13, 11 ca. MBH-4, 4 ca. MBK-3, ca. MKC-22, ca. MEG-14: CA.

**Description:** The carapace is subtriangular, its widest part is at the posterior 1/3, if the rostrum is omitted. The carapace is convex transversely, moderately so longitudinally. The two, strongly diverging rostral spines are almost as long as one half of the carapace. On the hepatic margin there are two spiny teeth. A pointed supraocular spine is also present. The branchial regions are adorned with three tubercles. The regions are well delimited. The cervical groove is the deepest one, but the hepatic region is delimited from the protogastric one by a deep furrow.

**Remarks:** It was not an easy decision to range this species among *Pisa* forms, where V shaped rostral spines are uncommon. Instead, spines of this type are present in *Rochinia* and various other forms. All other features, including shape the chela, and the well developed praeorbital spine are typical for *Pisa*. Perhaps this form is close to a hypothetical common ancestor of *Pisa* species.

**Ecology:** The species is most abundant where *Pachygrapsus* and *Brachynotus* occurs and where stromatolites or oncoïdes are present. All this indicates a shallow infralittoral or even an eulittoral environment. In contrast recent *Pisa* species live generally in a slightly deeper water, down to 100 meters, though *P. tetraodon* was reported from shallow waters too (1 meter, INGLE 1980, p. 139).

#### Subfamilia *Mithracinae* BALSS, 1929

##### Genus *Micippa* LEACH, 1817

###### *Micippa hungarica* (LÖRENTHEY in LÖRENTHEY—BEURLEN, 1929)

Pl. LII, figs. 4–6, pl. LIII, figs. 1–4.

1929. *Maia hungarica* n. sp. — LÖRENTHEY in LÖRENTHEY—BEURLEN: 34, 150, 151, pl. 7, fig. 2. (highly inaccurate!)

1953. *Maia austriaca* n. sp. — BACHMAYER: 244–245, pl. 2, fig. 3.

1953. *Phrynolambrus weinfurteri* n. sp. — BACHMAYER: 247–248, pl. 3, fig. 4.

1974b. *Micippa austriaca* (BACHMAYER) — MÜLLER: 277, pl. 1, figs. 5, 6.

1979a. *Micippa hungarica* (LÖRENTHEY) — MÜLLER: 274.

**Material seen:** holotype, Bia M 2330: FI.

Others, ca. Rákos, leg. FRANZENAU, M60/5964: NMÖ. 32 ca. MOE-17, 19 ca. MDG-19, 3 ca. MKC-21, 5 ca. M8-11, 2 ca. MBH-3, ca. MFK-6, 2 ca. MDZ-14: CA.

**Description:** The carapace is extremely convex in both directions. It is stout, not much longer than wide. The frontal part is directed downwards. There is a median longitudinal furrow on this part. The rostrum is subterminally bifurcated and terminates in two diverging short spines. The orbits are rather big, spiny, directed outwards. On all margins there are acute spines. The surface is strongly and almost evenly tuberculated. At the mesobranchial region there is an elevation decorated by an upward directed spiniform tubercle. A specimen from Rákos (pl. LII, figs. 5, 6) is apparently smoother than the others, but this can be attributed to a different preservation.

**Remarks:** With the very kind help of DR. GUINOT it was possible to compare the species with recent forms. The Badenian form is beyond any doubt quite similar to *M. thalia*, but the tubercles are closer on the surface of *M. thalia* HERBST, and individual variations might be even larger. The only argument against the fusion of the two species is that there are no links from intermediate geologic times (p. 48).

**Ecology:** The species is rare or absent in layers deposited in extremely shallow waters or on reefs. This is in accord with the fact that *M. thalia* lives in depth of about 20–100 meters (SAKAI 1976, p. 257) though other *M.* species prefer shallower waters or reefs.

### Familia Parthenopidae MCLEAY, 1838

#### Subfamilia Parthenopinae MCLEAY, 1838

#### Genus Parthenope WEBER, 1795

#### Parthenope szaboi MÜLLER, 1974

Pl. LIII, fig. 5, pl. LIV, figs. 1–9.

- ?1895. *Lambrus* sp. ind. — CREMA: 674, fig. 14.  
partim 1928. *Lambrus* sp. — GLAESSNER: 198.  
1974b. *Parthenope szaboi* n. sp. — MÜLLER: 277, 283, pl. 1, figs. 8, 9.  
1976c. *Parthenope szaboi* MÜLLER — MÜLLER: 152, pl. 4, fig. 2.  
1979a. *Parthenope szaboi* MÜLLER — MÜLLER: 274.

**Material seen:** holotype, ca. MDG-9-1 (= MG-2-1): NMÖ.

Paratypes, 4 ca. MDG-9: CA.

Others, 22 ca., 22 ch. MOE-15, 2 ch. MR-16, 16 ch. MR8-13, 4 ch. MRF-8, 2 ca., 6 ch. MKC-8, 5 ch. MGY-19: CA.

**Description:** The outline of the carapace is rhomboidal, rounded. The front consists of a trilobated rostrum. The anterolateral border is double. Between the hepatic and branchial parts there is a notch. This margin is adorned with even teeth, only the last one is slightly bigger. The median regions are elevated, the cardiac one is hump-like covered with numerous tubercles. Smaller and bigger tubercles are scattered on the whole surface, only the front is smooth.

The cross-section of the chela is triangular, all edges are covered with teeth or tubercles. The dactylus is curved. On its upper proximal part there is a flattened region delimited by spines.

**Remarks:** The species is extremely close to *P. angulifrons* extant form. Until intermediate forms are known their fusion would be risky.

**Ecology:** The species occurs in various rocks, but is rare in those deposited in shallow water. It was never found in reefal formations.

#### Parthenope tetenyensis n. sp.

Pl. LV, figs. 1–7, pl. LVI, figs. 1–7.

1897. *Lambrus* sp. ind. — LÖRENTHEY: 161.  
1898a. *Lambrus* sp. ind. — LÖRENTHEY: 106.  
1898b. *Lambrus* sp. ind. — LÖRENTHEY: 117, 152.  
1898c. *Lambrus* sp. ind. — LÖRENTHEY: 91.  
partim 1901a. *Andorina elegans?* — pl. 1, fig. 2.  
partim 1901b. *Andorina elegans?* — pl. 1, fig. 2.  
partim 1928. *Lambrus* sp. — GLAESSNER: 128.  
1929. *Andorina elegans* LÖRENTHEY — LÖRENTHEY in LÖRENTHEY—BEURLEN: pl. 8, fig. 3.  
1979a. *Parthenope* sp. nov? — MÜLLER: pl. 11, fig. 3, pl. 12, fig. 3.

**Material seen:** holotype, ca. MOE-16-1: CA.

Paratypes: 14 ca., 31 ch. MOE-16, 14 ca. MDG-8, ca., ch. MJL-6, 3 ch. MR-17, ca. MGY-18, ch. MR8-14, ca. MBH-11, 2 ca. MUO-2, 9 ch. MRF-9, 3 ca. MEG-11, da. MOS: CA.

**Locus typicus**: Budapest, Örs vezér tere, construction-pit.

**Stratum typicum**: Upper Badenian.

**Derivatio nominis**: Its first known occurrence was on the Tétény-plateau.

**Diagnosis**: Subtriangular carapace, strongly elevated median regions.

**Description**: The outline of the carapace is subtriangular, rounded at the lateral edges. The frontal part is rather wide, directed downwards. There is a deep median furrow on it. The median regions are highly elevated forming a structure resembling to the number 8. The branchial regions are also strongly elevated. The lateral margin is bordered by compound, flat lobe-like teeth. The species is strongly heterochelous, the right, crusher one is high, the other elongated. Their cross-section is four-sided.

The dactylus is flattened on its upper proximal part. The flat region is surrounded by tubercles.

**Remarks**: The species is similar to *P. massena*, a highly variable extant species, which has a tendency to being smooth, depleted of tubercles. No similar trend was observed at *P. tetenyensis*, nevertheless the range of their variability overlap. This species can be distinguished from *P. szaboi* even on the basis of a fragment of a chela, the cross-section of which is triangular at the former, four-sided at the other.

**Stratigraphy**: The species is known from the Upper Badenian. As the age of Oslip-locality is probably Middle Badenian, the species' range is longer. Perhaps the form is near to the ancestor of *P. massena*, extant species.

### Sectio CANCRIDEA LATREILLE, 1803

#### Familia *Corystidae* SAMOUELLE, 1819

Genus *Corystites* nom. nov., pro *Microcorystes* LÖRENTHEY in LÖRENTHEY-BEURLEN 1929, nom. praecocc.

non 1893. *Microcorystes* n. gen. — FRITSCH.

1929. *Microcorystes* n. gen. — LÖRENTHEY in LÖRENTHEY-BEURLEN: 137, typus-species: *M. latifrons* n. sp., by monotypy.

1969. *Microcorystes* LÖRENTHEY, non FRITSCH — GLAESSNER: R532.

#### *Corystites latifrons* (LÖRENTHEY in LÖRENTHEY-BEURLEN, 1929)

1904a. *Microcorystes latifrons* nom. nud. — LÖRENTHEY: 161.

1904b. *Microcorystes latifrons* nom. nud. — LÖRENTHEY: 30.

1929. *Microcorystes latifrons* n. gen. n. sp. — LÖRENTHEY in LÖRENTHEY-BEURLEN: 137–138, pl. 8, fig. 4.

Material not seen, broken and lost: ca. Budafok.

**Description**: LÖRENTHEY's figure shows a carapace which is elliptical, convex in both directions. The cardiac, branchial and mesogastric regions are well delimited from each other. The anterolateral margin bears 5 or 6 teeth. The rostrum is bilobed.

**Remarks**: As the specimen was found as early as 1904 or even earlier, one can hope that it was drawn by the artist who completed LÖRENTHEY's first figures which are all correct.

Indeed the shape of the species reveals similarities with *Micromithrax holsaticus* NÖTLING (GRIPP 1967, pl. 1, fig. 2, and GRIPP 1964, p. 123, pl. 21, fig. 8.) on one hand and with JONAS, extant genus, on the other.

### Familia *Canceridae* LATREILLE, 1803

#### Subfamilia *Cancriniae* LATREILLE, 1803

Genus *Cancer* LINNAEUS, 1758

#### *Cancer sismondae* (MEYER, 1843)

1859. *Lobocarcinus sismondae* MEYER — A. E. REUSS: 41, pl. 9, figs. 1–2.

?1884. *Cancer illyricus* n. sp. — BITTNER: 26–27, pl. 1, fig. 7.

?1905. „Hand eines Cancer” — TOULA: 163.

1924. *Cancer sismondae* MEYER — GLAESSNER: 115.

1928. *Cancer sismondae* v. MEY. — GLAESSNER: 176–177.

?1928. *Cancer cf. sismondai* v. MEY. — GLAESSNER: 177–178.

?1929. *Cancer* sp. — LÖRENTHEY IN LÖRENTHEY—BEURLEN: 162, pl. 8, fig. 7.

1982. *Cancer sismondai* MEYER — BONFIGLIO—DONADEO: 255–296, pl. 36–44. (exhaustive literature).

M a t e r i a l seen: almost entire specimen. Bruck: NHMW. ?ch. (lost?) Dévényújfalu, Devinska Nova Ves: ?coll.

R e m a r k s : The species is described in great detail by BONFIGLIO and DONADEO (1982). REUSS' specimen belongs most probably to this wide-spread and frequent Mediterranean species known from Miocene and Pliocene layers. An interesting point is, however, its rarity in the Paratethys. An exchange of label by REUSS, an extremely careful naturalist is by no means possible.

### Cancer styriacus BITTNER, 1884

Pl. LVII, figs. 1–4, pl. LVIII, figs. 1–2.

1884. *Cancer styriacus* n. sp. — BITTNER: 25–26, pl. 1, fig. 6.

1928. *Cancer styriacus* BITTNER — GLAESSNER: 178.

?1953. *Cancer sismondai* H. v. MEYER — BACHMAYER: 248, 249.

M a t e r i a l not seen, probably lost: holotype, ca. Aigen bei Fehring, Austria: ?Geol. Bundesanst.

M a t e r i a l seen: ca., fragment of ca., 2 ch. MOE-27, 4 ca. MDZ-4: CA.

D e s c r i p t i o n : The carapace is rather narrow, convex. The frontal region is narrow, there are three teeth on it between the intraorbital ones. The median region is depressed. The anterolateral margin consists of nine finely denticulated lobes. The regions are moderately well delimited, the deepest furrow is located on the sides of the urogastric and cardiac regions. The posterolateral margin is straight or slightly concave. The chela is stout, and its ridges consisting of small tubercles are low.

R e m a r k s : The Hungarian specimens are identical to BITTNER's figure except that the concavity of the anterolateral margin was most probably exaggerated by BITTNER's artist. This feature is variable, however. The form differs from all extant *Cancers* by its extreme narrowness. Probably *Cancer amphioctetus* is similarly narrow (RATHBUN 1930), but it is strongly ornamented.

E c o l o g y : The recent *C.* species live predominantly in moderately deep waters (some tens of meters). The species is frequent on MDZ locality of reef origin.

### Cancer illyricus BITTNER, 1883

Pl. LVIII, fig. 3.

1884. *Cancer illyricus* n. sp. — BITTNER: 26–27, pl. 1, fig. 7.

non 1893. *Cancer cf. illyricus* BITTNER — BITTNER: 32, 33.

M a t e r i a l seen as cast: holotype, ca. Zagor, Yugoslavia: coll. Univ. Graz.

R e m a r k s : The probably only specimen is slightly compressed laterally. Disregarding this distortion, the specimen is remarkably similar to small specimens of *C. sismondai* H. v. MEYER. (BONFIGLIO—DONATEO 1982, pl. 41, fig. 3N.) In small specimens a tubercular ornamentation of the protogastric region is clearly visible but is absent from the bigger ones. In this and other aspects there is no difference between the two forms of *Cancer*.

## Genus *Microdium* A. E. REUSS, 1867

T y p u s - s p e c i e s : *Microdium nodulosum* REUSS, 1867, by monotypy.

1929. *Cancer* — GLAESSNER: 104, 256.

### *Microdium nodulosum* A. E. REUSS, 1867

1867. *Microdium nodulosum* n. gen. n. sp. — A. E. REUSS: 173–176, pl. 8, figs. 7–8.

1929. *Cancer nodulosus* (REUSS) — GLAESSNER: 104.

?1929. *Cancer* sp. ind. — LÖRENTHEY IN LÖRENTHEY—BEURLEN 1929: 161, pl. 8, fig. 6.

M a t e r i a l not seen: holotype, ca. Wieliczka: NHMW.

Others: ?ch. Wieliczka: coll.?

?ch. Felménes: lost.

**R e m a r k s :** REUSS' description and figures well represent the species but it is prudent not to fuse the genera *Microdium* and *Cancer* in spite of all similarities until the holotype or new specimens are found.

**Genus *Miocycclus* MÜLLER, 1979**

**T y p u s - s p e c i e s :** *Miocycclus bulgaricus* MÜLLER, 1979, by monotypy.

***Miocycclus bulgaricus* MÜLLER, 1979**

Pl. LVIII, figs. 4–6, Fig. 10.

1979b. *Miocycclus bulgaricus* n. sp. — MÜLLER: 5, pl. 2, figs. 1–3, pl. 3, fig. 1.

**M a t e r i a l seen:** holotype, Oriakhovo, Bulgaria, leg. KOJUMDGIEVA, Nt533/a: coll. Univ. Sofia.  
Paratype, same data, Nt533/b.

**D e s c r i p t i o n :** The carapace is flat, ornamented with small evenly distributed tubercles on the surface. The anterolateral margin is decorated with eight undistinct lobes of different sizes. Instead of the last, ninth lobe there is a spine. The front is wide, cut into six lobes. The mesogastric region is wide, the furrows delimiting this and the cervical furrow are shallow but distinct.

**R e m a r k s :** The species is a quite peculiar representative either of the *Cancrids*, which is more likely, or the *Atelecyclids*. The wide mesogastric region is characteristic of the *Cancroidea*. The flat smooth surface rules out its identity with any of the Badenian *Cancroidea* species which are usually more adorned. The rather smooth surface of the *Cancer* carapaces is slightly similar to that of *M. bulgaricus*, but the wide frontal region rules out any closer relationship.




Fig. 10. *Miocycclus bulgaricus*  
I X Oriakhovo

**Genus “*Cancer*” as nomen collectivum\*.**

**R e m a r k s :** The name was widely used for fossil species as it was among the best known genus of recent Crabs. Having no better solution, it will be used for a variety of species described by three authors. Some of these species were transferred later to other, partly new genera.

**“*Cancer*” szontaghii LÖRENTHEY, 1897\***

Pl. LIX, figs. 1–5, pl. LX, figs. 1, 2.

1897. *Cancer Szontaghii* n. sp. — LÖRENTHEY: 159, 160, 167, 168.

1898a. *Cancer Szontaghii* LÖRENTHEY — LÖRENTHEY: 104, 113, 115.

1898b. *Cancer Szontaghii* LÖRENTHEY — LÖRENTHEY: 120–126, 153, pl. 8, fig. 4.

1898c. *Cancer Szontaghi* LÖRENTHEY — LÖRENTHEY: 94–99, pl. 8, fig. 4.

1929. *Atelecyclus szontaghi* LÖRENTHEY — LÖRENTHEY in LÖRENTHEY—BEURLEN: 156–158, pl. 8, fig. 9.

1958. *Cyclocancer szontaghi* (LÖRENTHEY) — BEURLEN: 15.

1969. *Atelecyclus szontaghi* LÖRENTHEY — GRIPP: 86, 87, pl. 3, fig. 1.

In press *Tasadia szontaghii* (LÖRENTHEY) — JANSSEN—MÜLLER.

**M a t e r i a l seen:** holotype, complete specimen, Tasádfő, M35: FI.

Others, a specimen in the same piece of rock than the holotype: FI.

2 ca. Ramsel, Hemmorian, Belgium (loan from A. JANSSEN).

**M a t e r i a l not seen:** ca. Segrahn, B. R. D.

**D e s c r i p t i o n :** The carapace is densely ornamented. All regions are delimited from each other by deep furrows. The front was probably decorated with four spiniform tubercles (including the

\* See Addendum.

intraorbital teeth), but the presence of a median tooth can not be ruled out. The teeth of the antero-lateral borders are composed of smaller tubercles. The ornament of the surface consists of bigger, compound tubercles and smaller, simple ones. The chelae are covered with rows of tubercles. The basis-ischium of the third maxilliped is *Cancer*-like.

**R e m a r k s :** The species was found in great number in Ramsel, Belgium. DR. A. JANSSEN kindly offered a joint study of this material which will probably elucidate the systematical position of this interesting species (JANSSEN-MÜLLER in press.). The third maxilliped is clearly *Cancer*-like, unlike that of *Atelecyclus*, thus a transitional position between the two genera is ruled out.

#### **“Cancer” szontaghii vindobonensis (BACHMAYER—KÜPPER 1952)\***

1952. *Atelecyclus szontaghi vindobonensis* nov. subspec. — BACHMAYER—KÜPPER: 2–6, text-figs. 1, 2.

**M a t e r i a l** seen: holotype, ca. Baden-Sooss: NHMW.

**R e m a r k s :** The subspecies is described in details by the two authors. The species is much less decorated than LÓRENTHEY’s. The outlines of the carapace are unknown. The problem of the Badenian “*Cancer*” species will be discussed later.

#### **“Cancer” carniolicus BITTNER, 1884\***

**M a t e r i a l** not seen, probably lost: syntypes, 4 ca. Stein, Yugoslavia, ca. Plesko, Yugoslavia: ?Geol. Bundesanstalt.

**R e m a r k :** There is no reason to repeat BITTNER’s description until some specimens will be found again.

#### **“Cancer” bittneri TOULA, 1905\***

?1893. *Cancer cf. illyricus* BITTNER — BITTNER: 32, 33.

1905. *Cancer Bittneri* n. sp. — TOULA: 161–168, text-figs. 1–5.

**M a t e r i a l** not seen, probably lost; holotype, Kalksburg: ?Technische Hochschule, Wien.

Others: ca. Sooss at Baden, numerous ca. Walbersdorf (Borbolya): NHMW and Geol. Bundesanstalt, Wien.

**R e m a r k s :** Though TOULA’s figures seem reliable, no definitive conclusions can be drawn. General remarks about the “*Cancer*” species\*. Four different “taxa” were presented under this dubious generic name. Unfortunately most specimens were lost or at least unavailable to me. Others are poorly preserved. Only “*C.*” *szontaghii* has a well-preserved extant holotype-specimen. BITTNER had serious doubts about the independence of his two species *C. illyricus* and *C. carniolicus* as early as in 1893. Later GLAESSNER assigned the walbersdorf specimens, described by BITTNER as *C. cf. illyricus*, to *C. bittneri*. All authors pointed out the similarities in superficial ornament of all these species. As the matter stands, it is almost impossible to draw any conclusions concerning these “species”. In fact it is not likely that a limited space and time-span could contain so many species from the same group.

**Familia Pirimelidae ALCOCK, 1899**

**Genus *Pirimela* LEACH, 1816**

#### ***Pirimela lorentheyi* (MÜLLER, 1974)**

Pl. LX, figs. 3–5.

1974a. “*Carcinus*” *lorentheyi* n. sp. — MÜLLER: 122, 126, pl. 2, figs. 4, 5.

1979a. *Pirimela lorentheyi* MÜLLER — MÜLLER: 274.

**M a t e r i a l** seen: holotype, ca. MB-4-1: NMÖ.

\* See Addendum.

**Description:** The carapace is rather narrow, convex transversely and longitudinally. Its surface is almost smooth. The mesogastric and cardiac regions are fused but well delimited from the other ones. The cervical furrow is also developed. On the hepatic and epibranchial regions there are elevations. The protogastric region is limited anteriorly by a step-like inclination. The front is trilobated, not counting the intraorbital lobes. The anterolateral teeth are not preserved.

**Remarks:** On my inquiry DR. L. B. HOLTHUIS suggested the transfer of the species from the *Portunidae* to *Pirimela*. Comparison with specimens of *P. denticulata* and the literature (INGLE 1980, pl. 15, fig. b.) provided convincing reasons for this move. The posterolateral margin of *P. lorentheyi* is straight while concave in the extant form. The frontal lobes are much more acute in *P. denticulata*. According to the article 31 c/i of the ICBN (STOLL et al. 1961) the original Hungarian spelling LÖRENTHEY should be modified by omitting the accents.

#### Genus *Trachypirimela* MÜLLER, 1974

**Type-species:** *Trachypirimela radula* MÜLLER, 1974 (= *Micromithrax grippi* MÜLLER, 1974).

#### *Trachypirimela grippi* (MÜLLER, 1974)

Pl. LXI, figs. 1–6.

1974a. *Trachypirimela radula* n. sp. — MÜLLER: 278, 279, 283, 284, pl. 2, figs. 5–6.

1974b. *Micromithrax grippi* n. sp. — MÜLLER: 279, 284, pl. 2, figs. 1–2.

1979a. *Trachypirimela grippi* (MÜLLER) — MÜLLER: 274.

**Material seen:** holotype, ca. MA-1-1 (= MDG-15-1): NMÖ.

Holotype of *Trachypirimela radula*, ca. MDG-15-2 (= MG-3-1): NMÖ.

Paratypes, 3 ca. MDG-15: CA.

Others: 4 ca. MDG-15, ca. MJL-12, ca. MUO-3, 2 ca. MOE-28: CA.

**Description:** The carapace is pentagonal, rather flat at young specimens while convex at older ones. The regions are well defined. The mesogastric and cardiac fields are well delimited laterally. The cervical groove is rather deep. The front is not wide, trilobated, but all specimens are damaged at this area. The anterolateral margin bears five unequal teeth. The extraorbital tooth is double. The surface is finely striated, and some bigger tubercles occur on the branchial region.

**Remarks:** The allometric growth of the species and the poor preservation of the type-specimen of *Micromithrax grippi* led to the introduction of two species which were even attributed to separated genera. Transitional specimens do occur, however. The species is similar to *Sirpus*, especially to *S. monodi* GORDON but it has acute frontal spines. No allometric growth occurs in *Sirpus*.

#### Sectio BRACHYRHYNCHA BORRADAILE, 1907

#### Superfamilia Portunoidea RAFINESQUE, 1815

#### Familia Portunidae RAFINESQUE, 1815

#### Subfamilia Portuninae RAFINESQUE, 1815

#### Genus *Portunus* WEBER, 1795

#### *Portunus monspeliensis* (A. MILNE-EDWARDS, 1860)

Pl. LXII, figs. 1, 2.

non 1834. *Lupea granulata* H. MILNE EDWARDS (= *Portunus granulatus* H. MILNE-EDWARDS)

1860. *Neptunus monspeliensis* n. sp. — A. MILNE-EDWARDS: 232.

1860. *Neptunus monspeliensis* n. sp. — A. MILNE-EDWARDS: pl. 4, fig. 1, pl. 5, fig. 1.

1860. *Neptunus granulatus* n. sp. — A. MILNE-EDWARDS: 241, pl. 3, fig. 1, pl. 7, fig. 2.

?1893. *Neptunus* cf. *granulatus* A. MILNE-EDWARDS — BITTNER: 11–12.

1897. *Neptunus* cf. *granulatus* A. MILNE-EDWARDS — LÖRENTHEY: 159.

1898a. *Neptunus* cf. *granulatus* A. MILNE-EDWARDS — LÖRENTHEY: 94, 113.

1898b. *Neptunus* cf. *granulatus* A. MILNE-EDWARDS — LÖRENTHEY: 110, 153, pl. 9, figs. 2–3.

1898c. *Neptunus* cf. *granulatus* A. MILNE-EDWARDS — LÖRENTHEY: 92, pl. 9, figs. 2–3.

1928. *Neptunus granulatus* A. MILNE-EDWARDS — GLAESSNER: 183, 184.

1929. *Neptunus granulatus* A. MILNE-EDWARDS — GLAESSNER: 267, references.

1929. *Neptunus granulatus* A. MILNE-EDWARDS — LÖRENTHEY in LÖRENTHEY—BEURLEN: 188, pl. 13, figs. 3–4, pl. 14, figs. 1, 4.  
 1968. *Neptunus granulatus* A. MILNE-EDWARDS — STANCU—ANDREESCU: 466, pl. 7, fig. 85.  
 1979a. *Portunus granulatus* A. MILNE-EDWARDS — FÜRSTER: 94, further references from the Paratethys.  
 1979a. *Portunus monspelliensis* A. MILNE-EDWARDS — MÜLLER: 274, 280, 288, pl. 18.

**M a t e r i a l seen:** holotype, Montpellier: Muséum Natl. Hist. Nat. Paris.  
 Others, 2 ca., sternum Mátraverebély-Szentkút, M61/8686 and M62/3152: NMÖ.  
 Ca. Rákos: FI.  
 6 ca., partly with sternum, Kishajmás, Mesek (MKS-1), 2 ch. MRF-13, 2 ch. MR-12: CA.

**D e s c r i p t i o n :** A detailed description of this well known and wide-spread form is unnecessary. Two important features deserve mention. The carapace is moderately wide, consequently the anterolateral resp. the posterolateral margins enclose a rectangle. The shape of the front is characteristic (PHILIPPE—SECRETAN 1971, p. 9), with its six subequal teeth separated by similarly subequal notches.

**R e m a r k s :** The species should be separated from *Portunus viai* SECRETAN in PHILIPPE—SECRETAN, a Burdigalian form which occurs in the Hungarian Eggenburgian as well in Budafok, Budapest, and was determined by LÖRENTHEY as *Neptunus granulatus* (LÖRENTHEY—BEURLEN 1929, p. 189, 7th paragraph).

### **Portunus neogenicus MÜLLER, 1979**

Pl. LXII, figs. 3, 4.

1979a. *Portunus neogenicus* n. sp. — MÜLLER: 280, 281, 288, pl. 19.

**M a t e r i a l seen:** holotype, ca. MGY-3-1: CA.

**D e s c r i p t i o n :** Though the specimen is fragmented, diagnostic features are present. The carapace is extremely wide, consequently the lateral margins form an angle of about 60°. The last lateral tooth is long, spiniform, directed forwards. The other teeth are arranged radially i.e. the anterior ones are directed more and more anteriorly. The shape of the teeth is characteristic.

**R e m a r k s :** The species is similar to *P. sanguinolentus*, but the lateral spine is directed forwards unlike in the extant form. Furthermore, there is only a rudimentary ridge on its carapace at the last tooth, while this ridge is developed in the Indo-West-Pacific form. No other similar *Portunid* was described.

### **Subgenus Monomia GISTEL, 1848**

#### **Portunus (Monomia) miocaenicus n. sp.**

Pl. LXII, fig. 5.

**M a t e r i a l seen:** holotype, ca. MRO-10-1.

**L o c u s t y p i c u s :** Budapest, railway-cut Rákos.

**S t r a t u m t y p i c u m :** Upper Badenian.

**D e r i v a t i o n o m i n i s :** From the age Miocene.

**D i a g n o s i s :** Rather wide front with two lobes between the intraorbital ones.

**D e s c r i p t i o n :** The carapace is hexagonal, moderately wide. The front is bilobated between the intraorbital lobes. The orbits are well-developed, the orbital margins are elevated. The antero-lateral margin is almost straight. The last tooth is long, directed outwards. A long ridge leaves this tooth which is bended forwards. Two more ridges are on the epigastric regions.

**R e m a r k s :** The form is a characteristic Monomia species. This subgenus has not been found yet as fossil. Considering the overall shape of the carapace, *P. (M.) gladiator* is the closest extant species. Its front is, however, trilobated and the lobes are more acute than those of the new form. The shape of the front is close to that of *P. (M.) brocki*, its general form is quite different, however. The surface of the carapace is rather smooth in contrast of most *M.* species.

### **Genus Charybdis DE HAAN, 1833**

#### **Subgenus Goniosupradens LEENE, 1938**

**Charybdis (Goniosupradens) mathiasi n. sp.**

Pl. LXIII, figs. 1–4, pl. LXIV, figs. 1–5, Pl. LXVII, figs. 3–5.

1976b. *Charybdis* n. sp. — MÜLLER: 518.

Material seen: holotype, ca. MRZ-5-1; CA.

Paratypes, 12 ca., 8 ch. MRZ-5, ca. MV-7, 3 ca., ch. MDZ-6: CA.

Locus typicus: Budapest, railway-cut Rákos.

Stratum typicum: Upper Badenian.

Derivatio nominis: From the Hungarian king MATHIAS, whose favorite spot was Visegrád, the first known locality of the species.

Diagnosis: Five big and two small anterolateral teeth. On the front there are six densely packed lobes.

Description: The carapace is octogonal, slightly convex transversely. At the proximity of the margins it is flattened. The front is broad, consists of 8 lobes including the extraorbital ones, which are situated higher than the others. The outer two of the six inner lobes are semicircular, the inner lobes are square and the intermediate, two are transitional in their shape. The orbits are big, their upper margin is denticulated. The anterolateral margin bears 7 teeth, two of them, the 2nd and 4th, are small, almost rudimentary, fused to the base of the preceding ones. The posterolateral margin is slightly concave. The branchial ridges are gently curved. They reach the border of the mesogastric region. The protogastric ridges are straight. The postfrontal ridges are replaced by ill-delimited elevated areas with clusters of small granulae. Similarly ornamented areas are situated on the outer anterior parts of the protogastric areas.

Chelae were found attached to a carapace. This male specimen displayed a marked heterochely, uncommon in *Charybdis* species: one chela had a slender palm and fingers, while the fingers of the other one were robust. Whether heterochely is confined to males or not, can not be proven now (pl. LXIV figs. 3–5, pl. LXVII figs. 3–5).

Remarks: The new species is rather distinct from most extant *Ch.* forms. *Ch. (G.) obtusifrons* LEENE is the closest extant form, but its frontal lobes are arranged less densely to each other.

Ecology: The species was found exclusively in fossil Coral-reefs. In Rákos it is rather abundant.

Stratigraphy: The species occurs in Lower and in Upper Badenian strata.

**Genus Thalamita LATREILLE, 1829**

**Thalamita fragilis MÜLLER, 1979**

Pl. LXV, figs. 1–2, 4–5, ?pl. LXV, fig. 3, pl. LXVI, figs. 1–7.

1883. *Portunien* sp. (= *Portunidae*) — BROCCHI: pl. 5, fig. 4b.

partim 1929. *Portunus rákosensis* n. sp. — LÖRENTHEY in LÖRENTHEY—BEURLEN: pl. 12, figs. 20, 21, pl. 15, fig. 4.

1976a. *Charybdis?* sp. — MÜLLER: 510, pl. 3, fig. 3.

1979a. *Thalamita fragilis* n. sp. — MÜLLER: 274, 281, 289, pl. 17, figs. 1–3.

Material seen: holotype, ca. MGY-4-1: CA.

Paratypes, 2 ca. MGY-4, 5 ca. MKC-15, ca. MA-5: CA.

Others, 8 ch. MR-15, ca., 2 ch. MR-8-17, ca., 4 ch. MRF-10, 2 ca. MKC-15, ch. MRL-6, ch. MRO-8, 6 ca., ch. MEG-10, 2 ch. MGY-4, 2 ch. MFA-9, 3 ca. MA-5: CA. ch. Rákos, two fixed fingers MOH-11, M83/162: NMÖ.

Description: The carapace is hexagonal, moderately broad, rather flat. The front is eight-lobed, between the big intraorbital lobes there are six densely arranged blunt lobes. The anterolateral margin consists of 5 teeth which are generally subequal but in one specimen the 4th teeth were much smaller than the others. The anterolateral margin is rather oblique thus the front is not very wide. There are well-developed epibranchial, metagastric and protogastric ridges (STEPHENSON—HUDSON 1957). The chelae are slender. On the outer side of the palm there are ridges which are almost invisible on some specimens. The upper edge of the palm is spinose.

Remarks: The species is closer to *Th. crenata* (LATREILLE) than to any other extant form (no other fossil forms are known). Nevertheless the protogastric ridges are longer in *Th. fragilis*. The front of the Miocene species is comparatively shorter therefore the anterolateral margins are more oblique. One specimen (pl. LXV, fig. 5) differs from the others by its shorter 4th anterolateral tooth and by its less oblique margin.

**Thalamita n? sp.**  
Pl. LXVII, figs. 1-2.

Material seen: ca. (internal mould) MRZ-13: CA.

Remarks: One carapace was found, which is similarly ornamented than that of *Th. fragilis*. It is however, much more convex and its epigastric ridge is cut into two parts by a notch.

Genus *Rakosia* n. gen.

Type-species: *Rakosia carupoides* n. sp., by monotypy.

Derivation nominis: From one of its localities, Rákos, Budapest.

Diagnosis: A wide smooth carapace with lobated front, 8 anterolateral teeth and an epibranchial ridge.

Remarks: No similar *Portunid* is known. The form is highly peculiar by its general shape resembling *Charybdis*, but its chela is much unlike it. Some features are similar to those of *Carupa* but the general shape is quite different suggesting the systematic position of the Miocene form among *Portuninae* rather, than among *Catoptrinae*. *Necronectes* differs by a spinose front, though probably is related.

***Rakosia carupoides* n. gen. n. sp.**

Pl. LXVIII, figs. 1-7.

Material seen: holotype, ca. 1984/40/2: NHMV.

Paratypes: 4 ca. MNH-6, 15 ca., 10 ch. MRZ-14, ca. MDZ-7: CA.

Locus typicus: Gross-Höflein (Nagyhöflány), Burgenland, Austria, quarry "Fenk".

Stratum typicum: Middle (or lowermost Upper) Badenian.

Derivation nominis: The species resembles *Carupa*, Indo-West-Pacific genus of *Portunidae*.

Diagnosis: See the generic diagnosis.

Description: The carapace is hexagonal, rather flat, smooth. The front is bilobated, the lobes are wide and undulated. The intraorbital lobe is separated from the frontal ones by a notch. The orbits are big, circular, their upper margin is cut by two notches each. The oblique anterolateral margin is ornamented with eight teeth. The last tooth is not much bigger than the others. An epibranchial ridge originates at this tooth. The posterolateral margin is almost straight. The chelae are slender, extremely smooth. The cross-section of the palm is elliptical. The fingers are adorned with furrows. The dactylus bears a big proximal (crusher) tooth on one side at least.

Remarks: As mentioned in the description of the genus the species is quite peculiar. Probably some fragments were earlier determined as *Necronectes* but its front is quite different.

Genus *Necronectes* A. MILNE-EDWARDS, 1881

***Necronectes schafferi* GLAESSNER, 1928**  
Pl. LXVI, figs. 8, 9, ? pl. LXIX, fig. 1.

1928. *Necronectes schafferi* n. sp. — GLAESSNER: 179–182, text-fig. 4, pl. 3, fig. 6.

Material seen: holotype, ca. Müllendorf (Szárazvám), Austria (probably the quarry at Äusseren Berg of "Müllendorfer Kreide A. G.", cf. DULLO 1983), No 1927 I 1: NHMW.  
Others, ca. MDZ-10, 13 ch. MRZ-19: CA.

Remarks: In addition to GLAESSNER's detailed description and figures it can be stated that the frontal spines illustrated by dotted lines on the original drawing are indeed present on the Diósd specimen, they are acute and directed slightly outwards.

**R e m a r k s :** The old genus *Macropipus* (= *Portunus* FABRICIUS, non WEBER) was recently split into two genera: *M.* and *Liocarcinus*. According to HOLTHUIS (pers. comm.) the latter is still heterogenous. The miocene species attributed to *Macropipus* actually belong to *Liocarcinus* and to a new genus described below.

The recent *L.* species live mostly in the East-Atlantic-Mediterranean realm, one species is widespread as occurs in the Indo-West-Pacific area, too. The species of the genus live generally in shallow waters, mostly from five to hundred meters, but some occur almost intertidally (ZARIQUIEY ALVAREZ 1968, PESTA 1918). Some of them are euryhaline.

**Liocarcinus rakosiensis (LÖRENTHEY in LÖRENTHEY—BEURLEN, 1929)**

Pl. LXIX, figs. 2–6, pl. LXX, figs. 1–8.

1911. *Portunus rakosiensis* nom. nud. — LÖRENTHEY: 529.  
1913. *Portunus rakosiensis* nom. nud. — LÖRENTHEY: 326.  
1929. *Portunus rakosiensis* n. sp. — LÖRENTHEY in LÖRENTHEY—BEURLEN: 171–174, pl. 13, fig. 1. (false reconstruction).  
non 1929. *Portunus rakosiensis* n. sp. — LÖRENTHEY in LÖRENTHEY—BEURLEN: 172–173, pl. 12, figs. 20–21. (= *Thalamita fragilis*), pl. 12, figs. 22–23. (= *Callianassa munieri*).  
?1968. *Portunus rakosiensis* LÖRENTHEY — STANCU—ANDREESCU: 466.  
1974b. *Macropipus rakosiensis* (LÖRENTHEY) — MÜLLER: 280, pl. 3, figs. 2–6.  
1979a. *Portunidae* gen. sp. ind. — FÖRSTER: 94–95, text-fig. 6, pl. 3, fig. 4.  
1979b. *Portunidae* gen. sp. ind. — FÖRSTER: 257–259, pl. 2, figs. 1, 5, pl. 3, figs. 1, 2.  
1979a. *Macropipus rakosiensis* (LÖRENTHEY) — MÜLLER: 274.

**M a t e r i a l seen:** lectotype designated herein, ca. Rákos, M2328/a: FI.

Paratype, ca. Rákos, M2328/b: FI.

Others, da. Węglerek, leg. KRACH, 20 ca., 20 ch. MOE-21, 25 ca., 10 ch. MDG-17, 10 ca. MJL-10, 5 ca. MB-15, 4 ca. MKC-16, 5 ca., 2 ch. MEG-15, ch. MOH-10, 3 ca., 2 ch. MGY-16, ca. Mátraverebély-Szentkút, Szent László-forrás: CA.

**M a t e r i a l not seen:** ?ca. Rugi, Transsylvania: ?coll.

**D e s c r i p t i o n :** The carapace is hexagonal, strongly decorated, slightly convex. The front is finely denticulate, almost straight but on most specimens with a faint median lobe. The antero-lateral margin consists of five teeth (in contrast to LÖRENTHEY's figure), these are more or less sharp-pointed depending on the molt-cycle. The regions are extremely well defined, densely covered with short ridges. There is a characteristic T shaped protuberance on the cardiac region. The palm bears 5 ridges on the outer and upper surfaces, the lowest of these runs to the fixed finger. The fixed finger and the dactylus are pentagonal in cross-section. There is a heterochely visible on the teeth of the cutting edges. Generally, but not invariably, the right chela is the crusher.

**R e m a r k s :** The species is close to *L. corrugatus* but its front is not trilobated. Other ornamented species, e.a. *L. depurator*, *L. puber* differ in their front.

**E c o l o g y :** The species occurs in most localities except of those of reefal or extremely shallow-water-origin.

**S t r a t i g r a p h y :** Occurs in Lower and Upper Badenian rocks.

**Liocarcinus oroszyi (BACHMAYER, 1953)**

Pl. LXXI, figs. 1, 2.

1953. *Carcinides oroszyi* n. sp. — BACHMAYER: 251–252, pl. 2. figs. 5, 5a.

**M a t e r i a l seen:** holotype, ca. Deutsch-Altenburg, leg. Oroszy: NHMW.

**D e s c r i p t i o n :** The carapace is slightly convex in two directions. It was probably subhexagonal, rather broad, but all specimens are incomplete. The front is absent on all specimens. The anterolateral margin is slightly arched, bears five teeth with forward directed points. The surface seems almost smooth to bare eyes, but actually it is adorned with small tubercles. The anterior median parts of the branchial region, near to the cervical groove, are strongly elevated.

**R e m a r k s :** The species is closer to *Liocarcinus* than to *Carcinus*. The anterior extension of the mesogastric region is invisible in *Carcinus* species but is developed in *L. oroszyi*. The arrangement

of all regions coincides with that of *L. arcuatus*. The elevated branchial region, however is the same as in *L. depurator*.

*Carcinus* is probably a Pliocene immigrant to its recent areal in the East-Atlantic—Mediterranean realms. Its former presence here has not been proven (HOLTHUIS 1949), known for a certainty from Icenian, Lower Pleistocene, but was reported from the Coralline Crag (Pliocene) of England.

**Liocarcinus kuehni (BACHMAYER, 1953)**

Pl. LXXI, figs. 3, 4.

1953. *Portunites kühni* n. sp. — BACHMAYER: 249–251, pl. 2, fig. 2.

M a t e r i a l seen: holotype, ca. Deutsch-Altenburg, leg. OROSZY: NHMW.

Others: 2 ca. MJL-9: CA.

D e s c r i p t i o n : The carapace is hexagonal, moderately convex. The front lacks on all specimens. The anterolateral margin is also damaged but the presence of five teeth is doubtless. The regions are well delimited. The whole surface is densely covered with ridges.

R e m a r k s : The species is close to *L. depurator* and to other *L.* species. Its dense adornment is a characteristic feature. From *L. rakosensis* differs by a denser decoration and by a more convex carapace. The specimens from MJL (Tétény) are even more convex than the holotype.

Genus *Miopipus* n. gen.

T y p u s - s p e c i e s : *Portunus pygmaeus* BROCCHE, 1883.

D e r i v a t i o n o f n o m i n i s : Composition of the name of the age Miocene and of *Macropipus*, a related genus.

D i a g n o s i s : Small elongated carapace, strongly elevated mesogastric and cardiac region, smooth surface.

R e m a r k s : The genus is much unlike *Liocarcinus* or *Macropipus* in its strongly elevated, rounded, smooth regions, the prominent frontal part is also peculiar. No similar form was found in the zoological literature. In fact the surface resembles that of *Psammocarcinus* as it was pointed out by BEURLEN or LÖRENTHEY (LÖRENTHEY-BEURLEN 1929, p. 174, last paragraph).

***Miopipus pygmaeus* (BROCCHE, 1883)**

Pl. LXXII, figs. 1–4.

1883. *Portunus pygmaeus* (sic!) n. sp. — BROCCHE: 1., pl. 5, fig. 4.

1897. *Portunus pygmaeus* (sic!) BROCCHE — LÖRENTHEY: 150.

1898a. *Portunus pygmaeus* BROCCHE — LÖRENTHEY: 93, 104.

1898b. *Portunus pygmaeus* BROCCHE — LÖRENTHEY: 109, 117.

1898c. *Portunus pygmaeus* BROCCHE — LÖRENTHEY: 91, 92.

non 1924. *Portunus pygmaeus* BROCCHE — GLAESSNER: 115–116.

1929. *Portunus pygmaeus* BROCCHE — LÖRENTHEY — BEURLEN 1929: 174–175, pl. 15, fig. 1.

1974b. *Macropipus pygmaeus* (BROCCHE) — MÜLLER: 279, pl. 3, fig. 1.

1979a. *Macropipus pygmaeus* (BROCCHE) — MÜLLER: 274.

M a t e r i a l not seen, probably lost: holotype, ca. Rákos: ?Sorbonne.

M a t e r i a l seen: 2 ca. MOE-24, 7 ca. ?6 ch. MDG-16, ca. MJL-7, ca. MBH-8, 6 ca. MKC-17, 5 ca. MEG-16, ca. MGY-15: CA.

D e s c r i p t i o n : The carapace is rather narrow, its outline is close to a pentagon. The trilobate front is narrow, prominent. The orbits are large, there are two undistinct notches on their upper margin. There are five distinct, sharp anterolateral teeth. The anterolateral margin is long, but the posterolateral one is even longer. The posterior margin is short. The surface is strongly adorned with the presence of the distinct areas but it is smooth. The mesogastric and cardiac areas are very convex.

Subfamilia *Carciininae* McLEAY, 1838

Genus *Portumnus* LEACH, 1814

1911. *Portumnus tricarinatus* nom. nud. — LÖRENTHEY: 529.  
1913. *Portumnus tricarinatus* nom. nud. — LÖRENTHEY: 326.  
1929. *Portumnus tricarinatus* n. sp. — LÖRENTHEY in LÖRENTHEY — BEURLEN: 167–168, pl. 15, fig. 3.  
1979a. *Portumnus tricarinatus* LÖRENTHEY — MÜLLER: 274.

Material not seen, lost: holotype, ca. Rékos: ?Univ. Budapest.

Remarks: LÖRENTHEY gave a detailed description of this species, his figure, however, possibly is not reliable (p. 32) but reveals some similarities with '*Xaiva*' *bachmayeri* (see below), regarding the shape of the carapace.

### Genus *Xaiva* MCLEAY, 1838

Remark: The genus is regarded here as a nomen collectivum.

#### '*Xaiva*' *bachmayeri* n. sp.

Pl. LXXII, figs. 5–7, ?fig. 8.

Material seen: holotype, ca. MOE-26-1: CA.  
Paratype, ca. MOE-26-2: CA.

Locus typicus: Budapest, Örs vezér tere.

Stratum typicum: Upper Badenian.

Derivatio nominis: From PROF. FRIEDRICH BACHMAYER, Vienna, researcher of fossil decapods.

Diagnosis: Subpentagonal carapace, strongly delimited mesobranchial and cardiac regions.

Description: The outline of the carapace is subpentagonal. The anterior part is more rounded than the posterior one. The carapace is moderately convex in both directions. The front is rather short, decorated with three acute teeth of equal size, situated at a line anterior to the intraorbital angles. The orbits are semicircular, moderately deep. There are two shallow notches on their upper margin. The anterolateral margin bears five teeth. The first, extraorbital and the third teeth are bigger than the others. The fifth is slightly damaged probably it was also big. The posterolateral margin is slightly concave at its anterior 1/3 length, while the posterior part is straight. A *Xaiva*-like chela was found in the same locality.

Remarks: '*Xaiva*' *bachmayeri*, as far as it is possible to judge from a carapace, is close to extant *Xaiva* species. *Xaiva macleayi* is similar in respect of its anterolateral teeth (the third one bigger than the adjacent ones), but its shape is different due to its more prominent frontal region. It must be added that the extant *X.* species are rather dissimilar. '*X.*' *bachmayeri* differs from LÖRENTHEY's figure of *Portumnus tricarinatus* in his longer front (but LÖRENTHEY remarked that the front of his specimen was damaged) and more importantly, in the shape of the furrows delimiting the mesogastric and cardiac regions. One can hardly imagine an artist making such errors.

### Genus *Mioxava* MÜLLER, 1979

Type species: *Mioxava psammophila* MÜLLER, 1979, by monotypy.

Remarks: The position of the genus among *Carcininae* is doubtful. The genus is probably closer related to other, primitive genera of *Portunidae* than to *Xaiva*. The question remains open, however, since the single damaged carapace available proves only the independence of the genus, since no similar front is known among the members of the family.

**Mioxaiva psammophila MÜLLER, 1979**

Pl. LXXIII, figs. 1-3.

1979a. *Mioxaiva* n. gen. *psammophila* n. sp. — MÜLLER: 274, 281, 289, pl. 16, figs. 1-3.

Material seen: holotype, ca. MGY-5-1: CA.

Description: The right anterior part of a carapace is preserved, broken behind the last anterolateral tooth. The outline of the frontal and anterolateral parts is almost subcircular. The surface is flat, only the branchial regions are deflected. The front is trilobate, the median lobe is sharp, the lateral ones, corresponding to the intraorbital lobes, are blunt and finely denticulated on their inner edges. The denticulation fades out medially. The inner orbital margin is convex, the extraorbital tooth is acute. The upper orbital margin is cut by two rudimentary notches. A sharp infraorbital tooth is also present. The five anterolateral teeth are sharp, they decrease in size posteriorly. The surface of the carapace was probably smooth but it is somewhat eroded obscuring the finest details. The regions are almost invisible.

Subfamilia *C at o p t r i n a e* BORRADAILE, 1903

Genus *Carupa* DANA, 1851

?**Carupa** cf. *tenuipes* DANA, 1851

Pl. LXXIII, fig. 7.

Material seen: 3 ca. MV-6: CA.

Remarks: The specimens were compared with those preserved in the Mus. Natl. Histoire Naturelle, Paris. All features seen on the Visegrád specimens were very much like in the recent and fossil crabs. The lobulation of the front was the most convincing of the similarities.

Subfamilia *C a p h y r i n a e* ALCOCK, 1899

Genus *Lissocarcinus* ADAMS and WHITE, 1849

Remark: The name is used in a collective sense.

'**Lissocarcinus**' *szoeraenyiae* (MÜLLER, 1974)

Pl. LXXIII, figs. 4-6.

1974b. *Thia szoeraenyiae* n. sp. — MÜLLER: 278, 283, pl. 2, figs. 3-4.

1979a. *Lissocarcinus szoeraenyiae* (MÜLLER) — MÜLLER: 274, 281, 282, 289.

Material seen: holotype, ca. MDG-18-1 (=MD-1-1): NMÖ.

Others, 2 ca. MDG-18, 2 ca. MDN-3, ca. MOE-25: CA.

Description: Only the carapace is known. It is subpentagonal, flat and smooth. The front is prominent, faintly trilobate. The orbits are small. The anterolateral margin is adorned with five lobes separated from each other by shallow notches. The regions are ill-delimited, only the unified meso-urogastric and cardiac region is separated laterally from the others by shallow furrows. The anterolateral and frontal margins are sharp.

Remarks: The species was unfortunately assigned to the genus *Thia*. On my inquiry DR. L. B. HOLTHUIS suggested me to transfer the species among the *Portunids*. Indeed, compared with specimens or figures of *Lissocarcinus orbicularis* (the specimens collected by L. Bíró, on New Guinea, figures of STEPHENSON—CAMPBELL 1960, pl. 3, fig. 2.) only minute differences were observed. The fossil form has no epibranchial ridge, present on *L. orbicularis*. The anterolateral lobes are separated

from each other by deeper notches in *L. szoeraenyiae*. The extant *L.* species are, however, very small due to their commensal mode of living. In this case the size could be an important taxonomic character.

### Superfamilia Xanthoidea DANA, 1851

#### Familia Xanthidae DANA, 1851

**R e m a r k s :** The subdivision of this familia seems still unsatisfactory according to GLAESSNER's (1969, p. R515) opinion. Consequently the genera will be presented in an alphabetical order except the evidently natural group of *Pilumnus* and related forms, which will be described together.

### Genus *Actaea* DE HAAN, 1833

**R e m a r k s :** The old genus *Actaea* was recently split into numerous genera on solid basis. Our species probably belongs to *A.* sensu stricto, cf. GUINOT (1969). Most of *A.* species live in shallow waters, some of them are restricted to coral-reefs.

#### *Actaea turcocampestris* n. sp.

Pl. LXXIV, figs. 1-5.

**M a t e r i a l seen:** holotype, ca. (pl. LXXIV figs. 1, 2) MFR-1-1: CA.  
Paratypes, 3 ca., 2 ch. MTZ-10: CA.

**L o c u s t y p i c u s :** Fertőrákos, near Sopron.

**S t r a t u m t y p i c u m :** Badenian (probable Middle B.).

**D e r i v a t i o n o m i n i s :** From one of the localities, Törökmező (Lat.: Turcus Campus).

**D i a g n o s i s :** Elliptical shape, short anterolateral teeth.

**D e s c r i p t i o n :** The anterior parts of the carapace form an ellipsis. It is moderately convex. The front is narrow, almost straight, cut into two parts by a notch. The anterolateral margin is rounded, it is adorned with five small, but distinct teeth, the last of them is situated far behind the largest section of the carapace. The regions are well delimited by deep furrows. The epigastric region is cut into two parts by a longitudinal groove. The inner of these parts is half as wide as the outer one. The whole surface is covered with small tubercles. The chela is stout, strongly tuberculated. The upper proximal part of the dactylus is also covered with tubercles.

**R e m a r k s :** In its overall shape, because of its concave posterolateral margin, the species is close to *A. savignyi*, the type-species of the genus. The tubercles covering the surface of this recent form are much coarser.

**E c o l o g y :** The species was found on coral-reefs, in accord with the habits of most extant *A.* species.

### Genus *Carpilius* LEACH in DESMAREST, 1825

#### *Carpilius antiquus* GLAESSNER, 1928

Pl. LXXV, figs. 1-5.

1877. *Carpilius* sp. — BITTNER: 436.  
1928. *Carpilius antiquus* n. sp. — GLAESSNER: 191-193, text-figs. 5-6, pl. 3, fig. 13.  
1968. *Carpilius antiquus* GLAESSNER — GUINOT: 323.  
1976b. *Carpilius* cf. *antiquus* GLAESSNER — MÜLLER: 516.

**M a t e r i a l** seen: holotype, Baden-Rauchstallbrunngraben, No. 1860 V 8: NHMW.

Others, 8 ca., ch. MV-11, ca. MTZ-21: CA.

Right ch., Grosshöflein (as MNH), in an Austrian private collection.

**D e s c r i p t i o n :** The carapace is ovoid, very smooth, moderately convex. The frontal margin is undulate, composed of two round intraorbital and wide median lobes. The anterolateral margin is smooth. Its curvature increases toward the lateral angle. There is only a rudimentary lateral tooth.

**R e m a r k s :** GLAESSNER made a reconstruction of the holotype, based on additional specimens. The holotype is however, in all details identical the Hungarian ones, which have but the rudiment of a tooth on the lateral margin. This feature is in contrast to all extant *C.* species. The miocene species is in other details extremely similar to *C. convexus* (FORSKÅL).

### Genus *Chlorodiella* RATHBUN, 1897

**R e m a r k s :** The genus was reported from the Miocene of Java (*Chlorodius junghuhni* MARTIN, 1880). Its extant species live in the Indo-West-Pacific, and one in the Caribbean.

#### *Chlorodiella mediterranea* (LÖRENTHEY in LÖRENTHEY—BEURLEN, 1929)

Pl. LXXVI, figs. 1–6, pl. LXXVII, figs. 1–4.

1911. *Zozymus mediterraneus* nom. nud. — LÖRENTHEY: 529.  
1913. *Zozymus mediterraneus* nom. nud. — LÖRENTHEY: 326.  
1929. *Zozymus mediterraneus* n. sp. — LÖRENTHEY in LÖRENTHEY—BEURLEN: 34, 215–216, pl. 11, fig. 9. (false reconstruction!).  
non 1976a. “*Zosimus*” *mediterraneus* LÖRENTHEY — MÜLLER: 507, 510, pl. 3, figs. 1, 4. (n. ssp., see below).  
non 1976b. “*Zosimus*” *mediterraneus* LÖRENTHEY — MÜLLER: 516, 520. (n. ssp., see below).  
1979a. *Chlorodiella mediterranea* (LÖRENTHEY) — MÜLLER: 274, 282, 289.

**M a t e r i a l** seen: lectotype designated herein, Rákos M 2327/a: FI.

Paratypes, Rákos M 2327/b, c, d, e: FI.

Others, 250 ca. numerous ch. MRZ-6, 25 ca. MDZ-10, ca. MTZ-22, 4 ca. MZZ-5, 20 ca. MNH-5: CA. 5 ca., 2 ch. Gross-Höflein (in matrix of holotype of *Dromilites koberi*): NHMW.

**D e s c r i p t i o n :** The wide carapace is elliptical. The front is wide, cut into two parts by a notch. The frontal margin is double, with a horizontal furrow between the two edges. The orbits are round. The orbital region is slightly elevated. The convex anterolateral margin is adorned with four rudimentary teeth, only the last one is slightly larger. The surface is smooth covered with very small tubercles in partly decorticated specimens. The regions are not separated. There is a postfrontal and an epigastric ridge, and a slight elevation behind the outer part of the cervical groove. On some specimens this elevation is almost invisible.

The species is heterochelous, usually the right hand is bigger. The palm is elongate, its height gradually increases distally. Its surface is smooth. The fixed finger and the dactylus are also long, the finger-tips are spooned.

**R e m a r k s :** LÖRENTHEY compared his species to *Zozymus* (recte: *Zosimus*) *lacvis* DANA, a species which is identical with *Neoliomera insularis* WHITE (ODHNER 1925). The similarity is superficial, however, DR. L. B. HOLTHUIS kindly drew my attention to the similarities between LÖRENTHEY's species and *Chlorodiella*. Indeed, the wide and double front, the long chelae with spooned fingertips are common in both forms. The carapace is round at the Miocene species while more angular in the recent ones. The epibranchial ridges are also more angular in extant *Ch.* species.

**E c o l o g y :** The species was found invariably in reefal or coral-bearing (MZZ) layers.

#### *Chlorodiella mediterranea* (LÖRENTHEY) *tetenensis* n. ssp.

Pl. LXXVII, figs. 5–7.

- 1976a. “*Zosimus*” *mediterraneus* LÖRENTHEY — MÜLLER: 507, 510, pl. 3, figs. 1–4.  
1976b. “*Zosimus*” *mediterraneus* LÖRENTHEY — MÜLLER: 516, 520.

**M a t e r i a l** seen: holotype, ca. MRZ-19-1: CA.

Paratypes, 2 ca. MRZ-19, 11 ca. MKK-7, 30 ca. MV-13: CA.

**L o c u s t y p i c u s :** Rákos, railway-cut, Budapest.

**S t r a t u m t y p i c u m :** Upper Badenian.

**Derivation nominis**: After the name of the locality Tétény, its first known occurrence.

**Diagnosis**: Strong epibranchial, hepatic and epigastric ridges.

**Description**: The carapace bears a strong ornament, the ridges on the epibranchial, hepatic and epigastric regions are accentuated, but rounded in their cross-section.

**Remarks**: The new form is similar to *Ch. mediterranea*, but the ornament of the carapace is much stronger. Transitional forms are rare. Though the two forms were found together, they are clearly separated. Some localities (MKK, MV) yielded exclusively this subspecies, while at Rákos it is rare, 3 specimens for 250 of *Ch. mediterranea*. At MZZ LÖRENTHEY's form occurs exclusively.

**Ecology**: This subspecies substitutes LÖRENTHEY's form in localities where the petrographical conditions indicate agitated waters during the deposition.

### ***Chlorodiella loczyi* n. sp.**

Pl. LXXVIII, figs. 1-4.

**Material seen**: holotype, ca. MRZ-7-1: CA.

Paratypes, 14 ca. MRZ-7: CA.

**Locus typicus**: Rákos, railway-cut, Budapest.

**Stratum typicum**: Upper Badenian.

**Derivation nominis**: In honour of the great Hungarian geologist, LAJOS LÓCZY senior.

**Diagnosis**: Comparatively narrow front ornamented with pits.

**Description**: The carapace is ovoid, rather wide. The front is not very wide. The orbits are big. The main regions are well delimited especially the gastric and hepatic regions are bordered by deep furrows. The frontal region is adorned with pits arranged in horizontal rows. Five to nine pits compose each of the individual rows. The uneven furrow that gives the front a doubled appearance, also contains pits.

**Remarks**: The species differs from extant *Ch.* forms by its narrow front, in this respect it is close to *Ch. junghuhni* MARTIN (Miocene of Java) a species which is much more convex however. From the badenian species it differs by its narrow front and by the pits on the frontal region. The decoration of the surface and of the anterolateral margin are similar to that of *Ch. mediterranea tetenyensis*.

### ***Chlorodiella juglans* n. sp.**

Pl. LXXVIII, figs. 5, 6.

**Material seen**: holotype, ca. MDZ-11-1: CA.

Paratypes, 4 ca. MTZ-11: CA.

**Locus typicus**: Diósd, sand-pit.

**Stratum typicum**: Upper Badenian.

**Derivation nominis**: The name of the locality (Diósd) originates from the Hungarian name of walnut or walnut-tree, *Juglans* in Latin.

**Diagnosis**: Four strong but blunt anterolateral teeth, wide hepatic lobe.

**Description**: The carapace is suboctogonal, the fronto-orbital margin is extremely wide due to the big orbits. There are four strong but blunt teeth on the anterolateral margin. The gastric regions are well delimited. The postfrontal, epigastric and hepatic ridges are high, step-like but not sharp. The hepatic ridge is far from the cervical groove, consequently the hepatic region is wide.

**Remarks**: The species is even closer to extant *Ch.* species than the other Badenian forms are. The overall shape the anterolateral teeth and the ridges are similar to those of *Ch. nigra*. The front of the Miocene species is narrower and the number of its anterolateral teeth is inferior, however. From the Miocene species it differs by the presence of the anterolateral teeth and by its well-delimited gastric regions.

## **Genus *Daira* DE HAAN, 1833**

1871. *Phymatocarcinus* n. gen. — REUSS: 326.

**Daira speciosa (REUSS, 1871)**

Pl. LXXIX, figs. 1–6, pl. LXXX, figs. 1–2.

1871. *Phymatocarcinus speciosus* n. gen. n. sp. — REUSS: 326, figs. 1–4.  
 1877. *Phymatocarcinus speciosus* REUSS — BITTNER: 437, pl. 1.  
 1928. *Daira speciosa* (REUSS) — GLAESSNER: 191.  
 1929. *Daira speciosa* (REUSS) — LÖRENTHEY in LÖRENTHEY — BEURLEN: 197–198, 12, figs. 10, 11.  
 1949. *Daira speciosa* (REUSS) — SCHOUPPÉ: 139.  
 1969. *Daira speciosa* (REUSS) — YANAKEVICH: 25–26, pl. 1, figs. 1–3.  
 1976a. *Daira speciosa* (REUSS) — MÜLLER: 511.  
 1976b. *Daira speciosa* (REUSS) — MÜLLER: 516.  
 1977. *Daira speciosa* (REUSS) — YANAKEVICH: 79–80, pl. 10, figs. 5, 6.  
 1979a. ?*Eriphia* sp. C. — FÖRSTER: 98, text-fig. 12, pl. 1, fig. 2.  
 1979a. *Daira speciosa* (REUSS) — MÜLLER: 275.

Material not seen, lost (GLAESSNER 1928, p. 191): holotype, ca. Baden-Rauchstallbrunnengraben.

Material seen: others, 2 ca. MRZ-12, 10 ca. MKK-6, 6 ca., 3 ch. MV-12, 5 ca., 2 ch. MTZ-6, 2 ca., ch. MNH-8: CA.

**Description:** The carapace is elliptical, moderately convex. The front is adorned with four teeth. The orbits are small. The anterolateral margin is long, extending far behind the widest part of the carapace. It bears numerous teeth, the anterior ones are smaller, the posterior ones larger, but this feature is variable. The regions are not well visible since the ornament is dense, the whole surface is covered with round tubercles, which are larger on the anterior, smaller on the posterior parts. The test is extremely solid due to the peculiar structure described in details by GUINOT (1967, figs. 7, 8).

The chela is robust, tuberculated. The tubercles are much bigger on its upper surface than on the lower one. The inner surface is partly smooth.

**Remarks:** Morphologically it is almost impossible to distinguish the form from *D. perlata* (HERBST), (GUINOT 1967, fig. 3), or from its fossil specimens from the Pliocene of Fiji (RATHBUN 1945, pl. 59, figs. E, F, G, H). The species is common in the Messinien of the surroundings of Oran (pers. comm. of SAINT-MARTIN).

**Ecology:** The species is found invariably in reefal structures in the Paratethys and in the Mediterranean Miocene.

**Genus *Haydnella* n. gen.**

**Type-species:** *Haydnella steiningeri* n. sp., by monotypy.

**Derivation nominis:** From the great composer, JOSEPH HAYDN, who worked in Eisenstadt (Kismarton), near to the type-locality.

**Diagnosis:** A narrow xanthid carapace, straight front, round epibranchial lobe.

**Remarks:** In its overall, narrow shape it is somewhat similar to *Paraxanthias* and *Xanthias*, but these genera have *Xantho*-like, transversely elongated epibranchial subregion, and a more extended hepatic region. This latter is rather narrow at the new form. It reveals similarities with *Monodaeus* but the carapace is narrower in the Miocene form. "*Titanocarcinus*" *pulchellus* has some common features with it but, as far as it is possible to see on MILNE-EDWARDS' figure (the type is lost), the relationship fails in details. MILNE-EDWARDS' form might not belong to *Titanocarcinus* with its Cretaceous type-species.

***Haydnella steiningeri* n. sp.**

Pl. LXXX, figs. 3–5, pl. LXXXI, figs. 1–4.

partim 1928. *Titanocarcinus vulgaris* GLAESSNER: pl. 3, fig. 10.

1976b. *Titanocarcinus* aff. *sismondae* MILNE-EDW. — MÜLLER: 516, 520.

Material seen: holotype, ca. 1984/40/5: NHMW.

Paratypes, 6 ca., 2 ch. MNH-9, ca. MTZ-16, 10 ca. MV-15: CA.

Ch. Baden-Rauchstallbrunnengraben: NHMW.

**Locus typicus:** Gross-Höflein (Nagyhöflány), Burgenland, Austria, Quarry "Fenk" (Buchinger).

**Stratum typicum:** Badenian (Middle or lowermost Upper).

**Derivation nominis:** From PROF. FRITZ STEININGER, Vienna, whose help was essential in the study of this and other species and localities.

**Description:** The form of the carapace is subhexagonal, with rounded sides. The carapace is moderately convex latitudinally, more convex longitudinally. The front is straight, bilobated, the lobes are separated by a notch. The interior orbital angles are separated from the front by notches, too. The frontal margin is double as there is a horizontal fissure among two keel-like denticulated ridges. On the anterolateral margin there are five blunt teeth, including the extraorbital one which is even blunter than the others. The posterolateral margin is slightly convex. The regions are well delimited from each other. The epigastric regions are wide. The epibranchial region is peculiarly narrow, and the hepatic one is restricted to a small area.

The chelae are tuberculated, most tubercles are arranged in rows. The fingers are rather slender.

**Remarks:** As it was told the form is distinct from any extant or fossil species as far as it is possible to judge from the figures and from a limited number of comparison made in the Mus. Natl. Hist. Nat. Paris.

### Genus *Panopeus* H. MILNE-EDWARDS, 1834

#### *Panopeus wronai* n. sp.

Pl. LXXXI, figs. 5, 6, pl. LXXXII, figs. 1-4, pl. LXXXIII, figs. 1-4.

1979a. ?*Eriphia* sp. D. — FÖRSTER: 98-99, text-fig. 13, pl. 1, fig. 1.

**Material seen:** holotype, ca. 1984/41/1: NHMW.

Paratypes, 2 ca., ch. MNH-10, 5 ca. MTZ-9, ca. MDZ-2: CA.

**Material seen as cast:** paratype, ca. Grobie, Poland: Univ. Warsaw.

**Locus typicus:** Gross-Höflein, Burgenland, Austria.

**Stratum typicum:** Badenian.

**Derivation nominis:** DR. R. WRONA, University of Warsaw, found the first specimen in Grobie.

**Description:** The outline of the carapace is hexagonal, its upper surface is rather flat. The front is straight, cut by a median incision. The orbits are small. The upper orbital margin is denticulated. The anterolateral margin is ornamented with three teeth, the last of which is rudimentary and situated behind the widest part of the carapace. In place of the first two teeth, there is a keel-shaped edge. There are four pairs of transverse ridges on the frontal parts of the epigastric, protogastric, hepatic and epibranchial regions. These ridges consist of fine tubercles. There are two transverse grooves on the epibranchial region, which are better visible on internal moulds. The legs are preserved in two Gross-Höflein specimens, which is an exceptional case. The ambulatory legs are rather slender. The chelae are robust, the right one is much larger than the other. The palms are smooth. On the proximal part of the dactylus of the left chela there is an elongated crusher tooth.

**Remarks:** As FÖRSTER (1969a, p. 99) observed the species is close to *Panopeus*. The chelae, found together with other parts, are typically *Panopeus*-like, including the degree of heterochely, the smooth and robust palm and the form of the crusher tooth. The four pair of ridges on the carapace are also typical for *Panopeus*. The two pairs of grooves on the branchial part, however, are absent from extant forms as emphasized by FÖRSTER. These are, however, almost invisible on the outer surface and probably are of inferior systematic value in differentiation on species-level. All extant *P.* species have two teeth instead of a keel on the anterior end of their anterolateral margin. These tend to fuse, however in *P. herbstii*, which is similar to *P. wronai* in other respects, too.

### Genus *Pilodius* DANA, 1852 (= *Chlorodopsis* A. MILNE-EDWARDS, 1873)

The genus is used here as a collective taxon.

#### ‘*Pilodius*’ *vulgaris* (GLAESSNER, 1928)

Pl. LXXXIII, figs. 5, 6, pl. LXXXIV, figs. 1-4.

partim, only the holotype! 1928. *Titanocarcinus vulgaris* n. sp. — GLAESSNER: 185-189, pl. 3, fig. 9, non: pl. 3, figs. 10-12.

non 1953. *Titanocarcinus vulgaris* GLAESSNER — BACHMAYER: 254-257, pl. 4, figs. 1-9, pl. 5, figs. 1-2.

**Material seen:** holotype, ca. Baden-Rauchstallbrunnengraben, No 1927 I 2: NHMW.

Others, 11 ca., 5 ch. MZZ-7, ca. MNH-12: CA.

**Description:** Strongly areolated, elliptical carapace. The frontal margin is composed of many tubercles and tends to be double, much like *Chlorodiella*. There are five anterolateral teeth, the last ones are directed outwards. At Zebegény (MZZ) this is the most frequent *Xanthid*, and the most frequent chela from there is similar to the carapace in ornament and size. The carapace is decorated with tubercles which are coarse around the lateral teeth, on the hepatic and on the epibranchial parts, and near the front. The palm of the chela is similarly tuberculated as seen on the figures.

**Remarks:** As it was emphasized earlier (MÜLLER 1974a, p. 123), *Titanocarcinus vulgaris* is a compound species. Its holotype bears features lacking from other *Xanthids* so frequent in Badenian deposits (see below, *Xantho moldavicus*). These are the coarse granulation near the margins, the compound front and granulation of the chela. The species reveals similarities with *Pilodius* species in the ornament of the chela and in the overall shape of the carapace. Unfortunately, the tip of the finger, which could provide clues for systematic position of the species, has not been observed yet. (SERÈNE—VAN LUOM, 1958, 1959.)

### Genus *Trapezia* LATREILLE, 1828

#### *Trapezia glaessneri* MÜLLER, 1976

Pl. LXXXIV, figs. 5, 6, pl. LXXXV, figs. 1–4.

1976b. *Trapezia glaessneri* n. sp. — MÜLLER: 517, 518, 521, pl. 1, fig. 3, pl. 2, figs. 1–3.

**Material seen:** holotype, ca. MV-1-1: CA.

Paratypes, 3 ca. MV-1: CA.

Others, 8 ca., ch., merus and carpus MV-1, ca. MTZ-11: CA.

**Description:** The carapace is flat, smooth, suboctogonal. The front is undulate, it has three blunt lobes on each side. The notches between the lobes are shallow only the median one is slightly deeper. The orbits are comparatively deep, inner part of the orbital margin is gently elevated. The anterolateral margins are slightly curved, subparallel to each other. The posterolateral margin is also curved.

**Remarks:** A short merus of a right cheliped was found. It is a typical *T. merus*. Only its inner surface is visible. On its upper margin there are five teeth. The proximal one is small, the three subsequent ones are larger, while the distal tooth is twice as big as the previous one, and it is cut into two parts by a shallow notch.

**EcoLOGY:** The species was found in localities where *Stylophora subreticulata* REUSS, a seriatoporid coral is also present. It is known that *T.* species are commensal with seriatoporid corals (GARTH 1974), explaining their absence from the Caribbean.

**Stratigraphy:** The species was found only in Lower Badenian layers. The genus, however, was reported from the Upper Eocene, too (MÜLLER 1974b). The Eocene species is transitional between *Trapezia* and *Tetralia* and should be attributed to a new genus.

### Genus *Xantho* LEACH, 1804

#### *Xantho moldavicus* (YANAKEVICH, 1977)

Pl. LXXXV, figs. 5–8, Pl. LXXXVI, figs. 1–5, pl. LXXXVII, fig. 1.

- partim! 1928. *Titanocarcinus vulgaris* n. sp. — GLAESSNER: 185, 189, non pl. 3, fig. 9, nec fig. 10, ? fig. 11.  
non 1928. *Titanocarcinus vulgaris armatus* n. sp. — GLAESSNER: 189–190, pl. 3, fig. 12.  
1953. *Titanocarcinus vulgaris* GLAESSNER — BACHMAYER: 254–257, pl. 4, figs. 1–9, pl. 5, figs. 1–2.  
?1968. *Titanocarcinus aff. vulgaris* GLAESSNER — STANCU—ANDREESCU: 466.  
1969. *Medaeus* sp. — YANAKEVICH: 26–27, pl. 1, figs. 4–6.  
1974a. *Xantho* cf. *incisus* LEACH — MÜLLER: 123, pl. 3, figs. 1–2.  
1974b. *Xantho* cf. *incisus* n. sp.? — MÜLLER: 280.  
1976a. *Xantho* cf. *incisus* LEACH — MÜLLER: 510.

- 1976c. *Xantho* cf. *incisus* LEACH — MÜLLER: 152.  
 1977. *Medaeus moldavicus* n. sp. — YANAKEVICH: 80–81, pl. 10, fig. 4.  
 1979a. *Xantho* cf. *incisus* LEACH — MÜLLER: 274, pl. 20, figs. 1–5.  
 1979a. *Xantho* cf. *vulgaris* (GLAESSNER) — FÖRSTER: 100–101, text-fig. 14, 15, pl. 3, figs. 1–3, pl. 4, figs. 1–4.  
 1979b. *Xantho* cf. *vulgaris* (GLAESSNER) — FÖRSTER: 263–264, text-fig. 11, pl. 3, fig. 4. (plates exchanged!).

**M a t e r i a l** not seen, unavailable: holotype, ca. Shepteban, Moldavia, USSR, No 1084-Ya: coll. Gos. Ped. Inst. Tiraspol.

**M a t e r i a l** seen: numerous fingers, Korytnica, Lychów, Węglinek, Poland, leg. BAŁUK and KRACH, 2 ca. WEG-5, 7 ca., 5 ch. MOE-31, 25 ca., 4 ch. MGF-6, ca. MKF-3, ca. MM-3, 6 ca. MDG-22, ca. MRC-3, ca. MRS-6, 30 ca. MJL-15, 4 ca. MBH-10, 15 ca. MB-13, 4 ca. MUO-6, 4 ca. MBK-5, ch. MR-15, 9 ca. MKC-19, 4 ca. MOH-14, ca. MA-7, 13 ca. MRZ-11, 4 ca. MKK-3, 4 ca. MTZ-12, ca. MDZ-8: CA. 2 ca. Deutsch-Altenburg, leg. Oroszy: NHMW.

**D e s c r i p t i o n :** The carapace is hexagonal, broad. The front is straight or slightly bent forwards, rather narrow. The orbits are of medium size. The anterolateral margin bears five teeth, with forward directed tips. The posterolateral margin is straight. All regions are well delimited. The epibranchial region is transversely elongated, the hepatic is moderately extended. The surface is either smooth or finely denticulated on some parts. The small tubercles tend to fuse into short ridges. The palm is either smooth or denticulated proximally. The fingers are of moderate length, smooth and strongly spooned. On the upper outer surface of the dactylus there is a more or less developed furrow.

**R e m a r k s :** The problem of this most puzzling Crab of the Paratethys has not been solved yet. The fingers occurring together with the carapaces are invariably spooned, thus "*Titanocarcinus*" *vulgaris armatus* with its pointed finger-tips might not belong here. The earliest available name was that of YANAKEVICH. Indeed, many specimens (e.g. from MDZ, pl. LXXXVI, fig. 4) fully corresponds to YANAKEVICH' figures. Some others, as the Węglinek one, deviates in its stronger adornment. Intermediate forms occur, however, but not in the direction of the ornament of '*Pilodius*' *vulgaris*, which may be a good species. Considering the mutual similarity of so many extant *Xanthids* it is still possible that *X. moldavicus* is a compound taxon and not a single species.

## Group of *Pilumnus* LEACH, 1815

**R e m a r k s :** This is a coherent group, as discussed by BALSS (1933). It is easily recognisable by an extremely wide gastric field, a feature not mentioned by zoologists, but useful for palaeontologists. The existence of this group can be traced back to the Eocene (unpublished findings of the author).

## Genus *Pilumnus* LEACH, 1815

### *Pilumnus mediterraneus* (LÖRENTHEY, 1897)

Pl. LXXXVII, figs. 2–5, pl. LXXXVIII, figs. 1–5.

1897. *Pilodius mediterraneus* n. sp. — LÖRENTHEY: 160, 167, 169.  
 1898a. *Pilodius mediterraneus* LÖRENTHEY — LÖRENTHEY: 105, 113, 115.  
 1898b. *Pilodius mediterraneus* LÖRENTHEY — LÖRENTHEY: 126–129, pl. 8, figs. 5, 6.  
 1898c. *Pilodius mediterraneus* LÖRENTHEY — LÖRENTHEY: 99–101, pl. 8, figs. 5, 6.  
 partim? 1928. *Pilumnus* sp. — GLAESNER: 190.  
 1929. *Chlorodopsis mediterraneus* (LÖRENTHEY) — LÖRENTHEY in LÖRENTHEY—BEURLEN: 34, 225–227, pl. 12, figs. 13–17, 19.  
 1953. *Chlorodopsis mediterranea* (LÖRENTHEY) — BACHMAYER: 253, pl. 3, fig. 5.  
 ?1953. *Chlorodopsis mediterranea* (LÖRENTHEY) — BACHMAYER—TOLLMANN: 314.  
 1974a. "Pilodius" *mediterraneus* LÖRENTHEY — MÜLLER: 122, pl. 3, fig. 3.  
 1974b. *Pilumnus mediterraneus* (LÖRENTHEY) — MÜLLER: 280.  
 1976a. *Pilumnus mediterraneus* (LÖRENTHEY) — MÜLLER: 510.  
 1976c. *Pilumnus mediterraneus* (LÖRENTHEY) — MÜLLER: 152.  
 1979a. *Pilumnus mediterraneus* (LÖRENTHEY) — MÜLLER: 274, pl. 21, fig. 3.  
 1979a. *Eriphia* sp. — MÜLLER: pl. 21, figs. 1–2.  
 1979b. *Pilumnus* sp. — FÖRSTER: 260, 261, pl. 3, fig. 6, pl. 5, fig. 1, 3, text-figs. 8, 9.

**M a t e r i a l** seen: lectotype designated herein, ca. Rákos, M31/a: FI.

Paralectotype, ca. Rákos, M31/b: FI.

Others, ch. Rákos, M61/5634: NMÖ.

60 ca., 35 ch. MOE-18, 8 ca., 4 ch. MGF-6, 10 ca., 6 ch., MM-2, 5 ca., 2 ch. MDG-24, numerous specimens MJL-14, 25 ca. MBH-9, 48 ca. MB-14, ca. MUO-5, 5 ca., ch. MBK-6, ca., 3 ch. MR-14, ca., 2 ch. MR8-19, ca. MRF-11, 10 ca., 5 ch. MKC-18, 2 ca., ch. MRO-9, 6 ca., 2 ch. MEG-18, 10 ch. MOH-12, 2 ca. MFA-9, 2 ca. MRZ-8, ca. MTZ-7, ch. Máttraverebélly-Szentkút, Meszestető: CA.

**Description:** The outline of the carapace is elliptical, convex. The front is tetra-lobate, the two inner ones are wide, the outer ones rudimentary, separated from the inner orbital angle by notches. There are five anterolateral teeth, the second one is small, but invariably present. The regions are either well- or ill-defined on various specimens. The young ones are generally better areolated. Behind the orbits, at their inner side there is a protuberance. The anterior and anterolateral parts are tuberculated, but the tuberculæ are in some cases almost invisible. The chelæ are moderately stout, the palm is either smooth or tuberculated. The fingers are also moderately stout, the upper proximal part of the dactylus bears usually some tubercles. The left chela is generally more slender than the other and in most cases it is better adorned.

**Remarks:** The form is extremely close to recent Mediterranean species. Using the key of ZARIQUIEY ALVAREZ (1968, p. 390) the Miocene form is closest to *P. spinifer* regarding its chelæ, but even closer to *P. hirtellus* judging from the carapace. Probably *P. mediterraneus* was close to the common ancestor of the numerous, but variable mediterranean species, a part of which is regarded by some authors as a variety or subspecies (PESTA 1918).

**Ecology:** The species is probably the most common crab in the Badenian. It occurs in all types of biotopes studied, even in reefs, but it is the most abundant in very shallow, almost eulittoral environments.

**Stratigraphy:** *P. mediterraneus* occurs in all substages of the Badenian.

#### Genus *Actumnus* DANA, 1851

##### *Actumnus telegdii* (MÜLLER, 1974)

Pl. LXXXIX, figs. 1-5, pl. XC, figs. 1-5.

1974b. *Pilumnus telegdii* n. sp. — MÜLLER: 280, 281, 284, pl. 2, figs. 7-8.

1979a. *Actumnus telegdii* (MÜLLER) — MÜLLER: 274, 282, pl. 22, figs. 1-3.

**Material seen:** holotype, ca. MDG-23-1 (= MG-4-1): NMÖ.

Paratypes, 2 ca. MDG-23: CA.

Others, 2 ca. MDG-23, 28 ca., numerous ch. MOE-19, ca. MEG-17, 7 ch. MOH-13, 7 ca. MGY-20: CA.

**Description:** The carapace is very convex, its outline is elliptical. The front seems to be bilobated at first sight but there is a rudimentary lobe between the median one and the intraocular angle. On the anterolateral margin there are five teeth, but the second one is very small. The teeth consist of groups of tubercles. The main regions are well delimited, the gastric ones are very wide. On the anterior parts of the metagastric, frontal, hepatic and metabranchial regions there are groups of tubercles, but the regions near the posterior margin bear also such ornaments.

The chelæ and the fingers are stout. The palms are covered with tubercles, but the lower side of the right hand is almost smooth. Heterochely is distinct. In some cases the left chela is stronger.

**Remarks:** The species can not be confused with *Pilumnus mediterraneus* due to its adornment. Its fingers are much stouter than those of LÖRENTHEY's species. The species is close to *A. squamosus* and *A. setifer*, extant forms. The carapaces of *A. telegdii* and *A. setifer* are similar, but their chelæ are slightly different. (CAMPBELL—STEPHENSON 1970, p. 283.)

**Ecology:** Usually the species is rarer than *P. mediterraneus*, but in deposits of deeper water they are abundant.

##### *Actumnus* n? sp.

Pl. XC, figs. 6, 7.

**Material seen:** ca. MZZ-5: CA.

**Remarks:** The only known carapace is damaged in its frontal part. Otherwise it is similar to *A. telegdii*, but its decoration is the most marked near the posterolateral border, in contrast to the other form. Probably it belongs to a new species.

#### Genus *Glabropilumnus* BALSS, 1932

**Remarks:** The name is used as a collective one.

'*Glabropilumnus*' n. sp.

Pl. XCI, figs. 1-3.

Material seen: 3 ca. MTZ-8, ?ca. MRZ-9: CA.

Description: Unfortunately no specimen is complete, the front is invariably damaged. The preservation is moderate. The carapace is moderately convex, rather smooth, but on the hepatic and epigastric regions there are fine tubercles. The regions are ill-defined except the mesogastric one. The extraorbital angle bears no tooth. There are three teeth on the anterolateral margin, but there is a hiatus between the orbit and the first tooth.

Genus *Pilumnopeus* A. MILNE-EDWARDS, 1863

*Pilumnopeus paratethyensis* n. sp.

Pl. XCII, figs. 1-5.

Material seen: holotype, ca. MRZ-10-1: CA.

Paratypes, 16 ca. MRZ-10: CA.

?Others, ?2 ch. MRZ-10: CA.

Locus typicus: Rákos, railway-cut, Budapest.

Stratum typicum: Upper Badenian.

Derivation nominis: From the name of the sea in which the species lived: Paratethys.

Description: The carapace is convex in both directions, especially longitudinally. Its outline is elliptical. The front is bilobate, the lobes are directed downwards. The orbits are big, the inner part of the orbital region is delimited from the frontal lobes by a deep furrow. The anterolateral margin is adorned with a lobe-like, flat first tooth, and three other teeth. The outline of the postero-lateral margin is slightly convex. A step-like epibranchial ridge leaves the last anterolateral tooth, which ridge is broken twice. At the anterior end of the protogastric region, behind the front, there is also a step-like ridge. The mesogastric region is well delimited, while the others are not.

Remarks: The species is extremely similar to *Pilumnopeus vaquelinii*, Indo-West-Pacific form. The only constant difference is that the branchial ridge is much sharper in the recent form. Other *P.* species, as *P. indica* and *P. makiana* are less close to the Miocene form, as there are more ridges on their carapaces.

'*Pilumnopeus*' *tetenensis* n. sp.

Pl. XCI, figs. 4, 5.

Material seen: holotype, ca. MB-16-1: CA.

Paratypes, ca. MB-16, 2 ca. MRS-5: CA.

Locus typicus: Tétény-plateau, Budapest.

Stratum typicum: Upper Badenian.

Derivation nominis: From the locality, Tétény.

Diagnosis: Small, subglobular carapace, a deep notch behind the lateral angle.

Description: The small carapace is convex. Its front is wide, bilobate. The interior orbital angles are separated from these lobes by moderately deep furrows. The anterolateral margin forms a sharp keel on its anterior one-third, elsewhere it bears three subequal teeth. Most regions are ill-delimited except the most anterior part of the protogastric lobe. A faint outline of the very wide metagastric region is visible. There is a characteristic depression behind the lateral angle resulting in a comparatively narrow posterior part of the carapace. The surface is smooth.

Remarks: The shape of the carapace is peculiar with the depression on its posterior part. No similar feature was observed in other *Pilumnus*-related forms. The extremely wide gastric region is characteristic for this group.

Ecology: The species was found in layers containing oncoids, deposited from extremely shallow waters.

Familia Geryonidae COLOSI, 1924

Genus *Geryon* KRÖYER, 1837

**Geryon cf. latifrons VAN STRAELEN, 1936**

Pl. XCIII, fig. 1.

- ?1936. *Geryon latifrons* n. sp. — VAN STRAELEN: 477–478.  
?1955. *Geryon latifrons* VAN STRAELEN — VIA BOADA: 217–222.  
1979a. *Geryon cf. latifrons* VAN STRAELEN — MÜLLER: 275, 282.

M a t e r i a l seen: ca., drilling Z-2, 36.3 m, leg. KÓKAY: CA.

R e m a r k s : The specimen belongs undoubtedly to *Geryon*, but the frontal parts lack, where the diagnostic features of VAN STRAELEN's species are located, leaves the determination doubtful. The material of the sample (fine-grained sandy silt) and other fossils (*Amussium cristatum*) suggests a deeper sublittoral environment.

**Familia Goneplacidae MCLEAY, 1838**

Subfamilia *Goneopleaciinae* MCLEAY, 1838

Genus *Goneplax* LEACH, 1814

***Goneplax gulderi* BACHMAYER, 1953**

?Pl. XCIII, figs. 2, 3.

- 1953b. *Goneplax gulderi* n. sp. — BACHMAYER: 143–145, pl. 9, figs. 1–3.  
?1979b. *Goneplax cf. gulderi* BACHMAYER — MÜLLER: 6, pl. 1, figs. 2–3.

M a t e r i a l not seen: holotype, ca. Baden-Sooss, Austria: NHMW.

M a t e r i a l seen: other, ?ca. Oriakhovo, Bulgaria, leg. KOJUMDGIEVA: coll. Univ. Sofia.

R e m a r k s : As PESTA (1918, p. 437) pointed out, *Goneplax rhomboides* is extremely variable. Thus the remnants from Carpathian (unpublished findings of the author) and Badenian remnants require revision.

**Subfamilia *Carcinoplaciinae* H. MILNE-EDWARDS, 1852**

Genus *Pilumnoplax* STIMPSON, 1858

**“*Pilumnoplax*” *carnuntinus* BACHMAYER, 1953**

1953. *Pilumnoplax carnuntinus* n. sp. — BACHMAYER: 257–258, pl. 3, fig. 2.

M a t e r i a l seen: holotype, Deutsch-Altenburg, Austria, leg. OROSZY: NHMW.

D e s c r i p t i o n : The frontal region and a part of the gastric region is preserved. The mesogastric region is delimited by a deep furrow. The frontal region is prominent, it is cut by a longitudinal groove. The front consists of two straight parts, separated by a notch.

R e m a r k : The species most probably belongs to *Carcinoplaciinae* though its generic position can not be clarified on the basis of a single damaged specimen.

**Familia Grapsidae MCLEAY, 1838**

Subfamilia *Grapsinae* MCLEAY, 1838

Genus *Pachygrapsus* RANDALL, 1839

**Pachygrapsus hungaricus MÜLLER, 1974**

Pl. XCIII, fig. 4, pl. XCIV, figs. 1–3, 5–7.

1974a. *Pachygrapsus hungaricus* n. sp. — MÜLLER: 123–124, 126–127, pl. 4, figs. 1–3, pl. 5, fig. 1.

**M a t e r i a l seen:** holotype, ca. MB-5-1: NMÖ.

Paratypes, 8 ca. MB-5: CA.

Others, ca. MB-5, 12 ca. MJL-17, 17 ca., 2 ch. MRZ-4, 9 ca. MGF-7, 12 ca. MRS-7, 2 ca. MDZ-9, 2 ca. MBH-12, 3 ca. MBK-7: CA.

**D e s c r i p t i o n :** The carapace is almost square, slightly convex. Its front is very wide, virtually straight but faintly rounded at the inner orbital angles. The orbits are wide, open, externally delimited by teeth the axis of which diverges from the longitudinal axis of the carapace. The angle of this diversion decreases with increasing age of the individuals. There are two smaller teeth on the lateral margin. The posterior margin is wide, slightly shorter than the front. The chela is smooth, the fixed finger is spooned.

**R e m a r k s :** The species is closely allied to *P. marmoratus*, an extant form. There is a slight but well defined difference between them. The extraorbital tooth is more diverging in individuals of the same size in the Badenian form than in the extant one. For sake of comparison a photo of *P. marmoratus* is given (Pl. XCIV, fig. 4). The specimen is of same size than its Badenian counterpart on fig. 5. The Badenian form is slightly narrower at the posterior part of its carapace than the extant one.

**E c o l o g y :** The genus *P.* is well known for its eu- and supralittoral life on rocky shores. Beyond doubt its fossil representative lived also in similar habitats. It was found near to stromatolites or among oncoids, both usually intertidal features. *P.* occurred in reef-structures as well. At Rákos, there is sedimentological evidence for a temporal desiccation e.g. traces of roots, crashes. Rocky surfaces were present on the reefs and among oncoids.

The extremely poor preservational potential of eulittoral crabs makes this finding of special interest.

Subfamilia *V a r u n i n a e* H. MILNE-EDWARDS, 1852

Genus *Brachynotus* DE HAAN, 1833

**Brachynotus februariois MÜLLER, 1974**

Pl. XCV, figs. 1–5.

1974a. *Brachynotus februariois* n. sp. — MÜLLER: 123, 124, 127, pl. 5, figs. 2–4.

1979a. *Brachynotus februariois* MÜLLER — MÜLLER: 275.

**M a t e r i a l seen:** holotype, ca. MB-6-1: NMÖ.

Paratypes, 2 ca. MB-6, ca. MKF-5, ca. MM-4: CA.

Others, 16 ca. MB-6, 8 ca. MRS-8, 6 ca. MJL-16: CA.

**D e s c r i p t i o n :** The outline of the carapace is square. The front consists of two median lobes with sinusoid shape and of two intraocular teeth. The orbits are large, deep. The anterolateral margin bears three teeth, the last one is situated at the widest cross-section of the carapace. The posterolateral margins gently converge posteriorly. The surface is remarkably smooth. The median part of the cervical groove is well developed. No other grooves are visible. There is a faint epibranchial ridge originating from the last lateral tooth.

**R e m a r k s :** The species is closely allied to *B. sexdentatus* and to *B. foresti*. In recent forms the widest part of the carapace is situated at the 2nd tooth, but at the third one in *B. februariois*. The epibranchial ridge is far less sharp at the fossil form.

**E c o l o g y :** The species occurs together with *Pachygrapsus hungaricus* in accordance with the fact that extant *B.* species live in the highest infralittoral parts of the seas, and look for shelter under stones or in burrows of *Upogebia* (PESTA 1918, p. 450), but *Brachynotus februariois* was not found in reef-structures. Actually this species prefers bottoms of loose sediments while *Pachygrapsus* shelters in crevices of solid rocks.

Superfamilia *O c y p o d o i d e a* RAFINESQUE, 1815

Familia *Ocypodidae* RAFINESQUE, 1815

Subfamilia *O c y p o d i n a e* RAFINESQUE, 1815

Genus *Ocypode* WEBER, 1795

?**Ocypode** sp.

1979a. *Ocypode?* sp. — MÜLLER: 275, 282, pl. 23, figs. 1, 2.  
1980. *Ocypode?* sp. — VIA BOADA: 60, fig. 1.

M a t e r i a l seen: 2 ch. MGY-22: CA.

R e m a r k s : The outline of the chelae is similar to that of *Ocypode* species, but stridulating ridges are absent from the inner surface of palm. Some grapsid species, e.g. *Sesarma* forms have similar chelae, too.

Subfamilia *M a c r o p h t h a l m i n a e* DANA, 1852

Genus *Macrophthalmus* DESMAREST, 1823

**Macrophthalmus vindobonensis** GLAESSNER, 1924

Pl. XCV, fig. 6, pl. XCVI, figs. 1-2, 4-5, ?fig. 3.

- ?1879. *Macrophthalmus aquensis* n. sp. — A. MILNE-EDWARDS — BROCHI: 115.  
1924. *Macrophthalmus vindobonensis* n. sp. — GLAESSNER: 109-114, text-figs. 1-2.  
1928. *Macrophthalmus vindobonensis* GLAESSNER — GLAESSNER: 196-197.  
?1952. *Macrophthalmus aquensis* MILNE-EDWARDS — BROCHI — REMY: 27-39, text-fig. 1.  
1979a. *Macrophthalmus vindobonensis* GLAESSNER — FÖRSTER: 102, text-fig. 16, pl. 2, fig. 6.  
1980. *Macrophthalmus vindobonensis* GLAESSNER — VIA BOADA: 5-6, pl. 1, figs. 1-2.

M a t e r i a l seen: types, ca. Carpathian: NHMW.

Others, 3 ca. Niskowa, Poland: coll. Univ. Warsaw.

M a t e r i a l seen: others, 3 ca. MH34-1: CA.

?Syntypes of *Macrophthalmus aquensis*, 5 ca: Mus. Natl. Hist. Nat. Paris, a specimen from Teiritzberg, the type-locality, leg. STEININGER.

R e m a r k s : The Badenian forms do not differ from *M. aquensis* from France except for the fact that the surface of the best preserved type-specimen (external mould, casted with silicone rubber) is coarser granulated than that of the Badenian ones. This is, however, a variable feature, and also influenced by the state of preservation (REMY 1952). The type- and other specimens from Teiritzberg are usually somewhat narrower, than those from Hungary and France, also their lateral margins are somewhat shorter making the orbital margins more oblique than in our specimens. All these features are variable, even on the both sides of one and the same specimen (pl. XCVI, fig. 1). The (possibly only) European fossil *Macrophthalmus* species is remarkably similar to some extant forms, e.g. *M. barnesi* (SERÈNE 1971, p. 918), which also belongs to the subgenus *Venitus*.

E c o l o g y : Most *M. species* live in muddy eulittoral environments, some in Mangrove-swamps. Some species live however, in deep infralittoral regions, too. *M. (Venitus) latreillei* was reported from depth of 10 to 20 meters (SAKAI 1976, p. 616).

Superfamilia uncertain

Familia *Palicidae* BOUVIER, 1898

Genus *Palicus* PHILIPPI, 1838

**Palicus** sp.

Pl. XCVII, figs. 6, 7.

M a t e r i a l seen: 2 ca. MOE-32: CA.

R e m a r k s : The overall shape of the carapace and the ornament of the branchial regions is very much like that of *P. caronii*. The anterolateral margin is sharp, however, and cut by a notch into lobes, without teeth. The preservation of the specimen is rather poor at the margin, thus it is not impossible that the lobe-like feature might represent a fracture.

Genus *Crossotonotus* A. MILNE-EDWARDS, 1873

**Crossotonotus diosdensis n. sp.**

Pl. XCVII, figs. 1-5, Fig. 11.

**M a t e r i a l** seen: holotype, ca. MDZ-15-1: CA.

Paratype, ca. MDZ-15: CA.

**L o c u s t y p i c u s :** Diósd, south from Budapest.


**S t r a t u m t y p i c u m :** Upper Badenian.

**D e r i v a t i o n o m i n i s :** From the name of the locality.

**D i a g n o s i s :** More or less hexagonal carapace, lateral teeth flat and branching.

**D e s c r i p t i o n :** The carapace is strongly adorned, uneven, moderately convex. The front is damaged on both specimens. The orbits are big, cut by two notches. The anterolateral margin is adorned with four big teeth, two of which are branching. The cervical furrow is very deep. The epibranchial region is delimited from the more or less unified group of the cardiac, meta- and postero-branchial regions. All regions bear rather large regularly arranged tubercles.

**R e m a r k s :** The species is close to the extant *C.* forms in the peculiar arrangement of the regions, the tubercles, grooves. The anterolateral teeth are bigger, however, than those of any known form. No other fossil form of this genus is known. The most closely allied extant species is *Crossotonotus gardineri* (RATHBUN) which has lateral spines tending to be flat and even branching in the case of the third one. The teeth are much finer at the extant species than at *C. diosdensis*.



*Fig. 11. Crossotonotus diosdensis 3 × Diósd*

The Badenian decapod fauna is modern in its character. Only 12 of its 72 genera can be regarded as extinct thus about 85% of the genera still have living representatives. These numbers and ratios inevitably depend on certain subjective taxonomic decisions but support the above statement. Some archaic features of this fauna must not be overlooked, however. Callianassids are surprisingly diverse partly due to their good preservational potential. The probable homochely of the Badenian forms contrasts with the heterochely of most extant related forms. This situation requires further studies.

Galatheids and porcellanids are close to their recent relatives, in conformity with the high age of these groups. The same consideration applies to pagurids as well. In cases where direct comparison was possible, the fossil chelae were found not to differ substantially from those of their extant relatives (*Pagurus*, *Petrochirus* or *Dardanus arrosor*). *Albunea asymmetrica* is very similar to some extant species of its genus.

Uncommonly high is the number of Badenian dromioid species. Six different forms were detected although it was generally accepted that Neogene dromioids are rare. If forms bound to coral-reefs are omitted only two species remain, however, still more than the Mediterranean shallow-water forms. One, possibly homoloid species (?*Dorippe carpathica*) might be added to the list of *Dromiacea*.

Although calappid forms are not very numerous, their number still exceeds that found in the Mediterranean because genera recently confined to the Indo-West-Pacific are also present. The species do not differ substantially from their extant relatives, though *Mursia* and *Matuta* species represent primitive forms within their genera as indicated by their short lateral spines. Such presumably primitive species persist, however, in the Indo-West-Pacific. The phylogenetic significance of lateral spines is discussed by IMAIZUMI (1958). The leucosids are represented by eight species, five of which were determined to species level. The diversity of *Ebalia* species is high, similar to that observed in the Mediterranean. Two or three species belong to extinct genera or are of uncertain affinity.

A striking feature of the Badenian decapod fauna is the low number of majid species. Only five species are known, one of which was an inhabitant of reefs. There is a curious absence of see-grass-dwelling small forms, which, together with other small species, represent a major component of the Mediterranean crab-fauna. Three of the species belong to genera still present in the Mediterranean while the remaining two, including the reef-dwelling one, have extant Indo-West-Pacific relations. Of all crab species discussed the two parthenopids of the Badenian are most closely related to common Mediterranean forms.

The cancroids of the Badenian show several peculiar features. Forms belonging to the group of "Cancer" *szontaghii* seem not to have close extant relatives. The total number of cancroid species is rather high, especially in comparison to their number in the Mediterranean, even if this number inevitably includes also some erroneous identifications as well. The number of portunid species is similarly high, at least 17. Some of the species belong to modern genera, others belong to extinct ones. *Portunus*, *Liocarcinus*, *Thalamita* and *Charybdis* species are similar to their Mediterranean or Indo-West-Pacific counterparts. This family is important in modern faunas thus the rate observed in the Badenian record is moderate. As in modern seas, the number of xanthids is high, especially that of reef-dwelling ones.

The Badenian record represents only a small sample of the Neogene decapods of the World. Thus only few conclusions can be drawn about the evolution of this group on the basis of these data. It can be stated that evolution, at least in this area, was rather rapid for majids from the Miocene until today. Whether this fast change reflects immigrations or evolutionary changes in situ, cannot be decided yet. The *Dromiacea* display a regression in the area but this process depended on the disappearance of reefs which provided a habitat for most Miocene species of this group. Regression of

cancroids seems a real process of evolution. The group plays still an important role in American waters, however. The regression of xanthids, similarly to that of dromids, is connected with the extinction of coral reefs in European waters. If one compares their number with that of the Indo-West-Pacific forms, no regression is evident. The absence of some widespread recent groups from the Badenian record suggest a rapid evolution of this family. Other families are represented by low number of species thus do not permit speculations about evolution.

Three subunits are distinguished within the Badenian, an early (Moravian), a middle (Wieliczian) and a late (Kosovian) substages. The distinction between them is usually not easy if individual sections are considered. These substages are defined by stratotypes and by boundary stratotypes (PAPP et al. 1978), based mainly on foraminifers or on nannoflora. These fossils or at least the most characteristic forms are absent from limestones which provide the bulk of fossil decapods. Such difficulties notwithstanding, the crab-bearing rocks can be age-correlated to planctonic or benthic foraminifer-bearing strata through the use of geological, palaeogeographical methods or of molluscs which occur in both types of sedimentary rocks. The use of the former methods is facilitated by significant tectonic events occurring between the Early and Late Badenian times which changed the palaeogeographic pattern of a large part of the Central Paratethys. Of the two different areas, containing rich crab-faunas in the vicinity of Budapest one is of Upper, the other is of Lower Badenian in age.

Attribution of the limestones occurring within Budapest and in its closest vicinity (Rákos, Tétény-plateau and Bia, cf. chapter "Description of localities") to Upper Badenian is supported by its molluscs (BÁRTKÓ-KÓKAY 1966) and by the fact that they are covered by Sarmatian layers without disconformity. The radiometric age of rocks at Rákos (BALOGH-ÁRVÁNÉ-SÓS-PÉCSKAY 1980) is  $13.4 \pm 0.6$  million year B.P. which strongly supports their Upper Badenian age.

On the other hand, an area in the Pilis- and Börzsöny Mountains, at both sides of the Danube (Visegrád, Törökmező, Zebegény, p. 43) contains sedimentary rocks of Lower Badenian age, confirmed by their micro- and macrofossils (BÁLDI-BEKE et al. 1980, NAGYMAROSI 1980).

The decapod faunas of these two areas differ from each other in several aspects due to their different age. Ecologically similar conditions were present in both areas. The most marked difference between them is the mutually exclusive presence of some abundant species. The most common of these are in Lower Badenian sites:

- Dardanus arrosor*
- Kromtis koberi*
- Dynomene emiliae*
- Calappa praelata*
- Carpilius antiquus*

Exclusively in Upper Badenian rocks were found:

- Petrochirus priscus*
- Dardanus substriatiformis*
- Dromia eotvoesi*
- Calappa heberti*
- Matuta brocchii*

The differences between Upper and Lower Badenian faunas are the most obvious if one considers coral-reef dwelling forms. The well-known habitats of these remained quite similar ecologically. Table 2 presents these differences.

Though the list is not definitive, especially not for group 2, the differences in a number of species are sufficient to support the possibility of a stratigraphically determined change. According to PAPP and STEININGER (in PAPP et al. 1978, p. 199) marly layers situated in the vicinity of the Fenk-quarry in Gross-Höflein are of lowermost Upper Badenian age. These marly layers can be parallelized with some equally marly ones in the Fenk-quarry itself. As these are situated higher in the sequence as the decapod-yielding reef ("Oberer Korallenkalk"), this latter might belong to the uppermost Middle Badenian or to the lowermost Upper Badenian substages. Thus there are three biozones in Badenian layers which can be differentiated on the basis of reef-dwelling decapods and partly of corals.

Table 2

## Corals and decapods of groups of reef-origin decapod localities

Number of group (biozone)	1	2	3
<b>Corals</b>			
<i>Portites</i> sp.	+	+	+
<i>Tarbellastraea</i> sp.	+	+	+
<i>Acanthastraea</i> sp.	+	+	-
<i>Stylophora subreticulata</i>	+	+	-
Other hermatypic forms	+	-	-
<b>Decapods</b>			
<i>Petrolisthes haydnii</i>	+	+	-
<i>Petrolisthes magnus</i>	-	-	+
<i>Dardanus arrosor</i>	+	-	-
<i>Dardanus substrriatiformis</i>	-	+	+
<i>Dardanus hungaricus</i>	+	-	+
<i>Krommtilis koberi</i>	+	+	-
<i>Dynomene emiliae</i>	+	+	-
<i>Dromia neogenica</i>	+	-	+
<i>Calappa heberti</i>	-	-	+
<i>Charybdis mathiasi</i>	+	-	+
<i>Carupa</i> cf. <i>tenuipes</i>	+	-	-
<i>Rakosia carupoides</i>	-	+	+
<i>Panopeus wronai</i>	+	+	+
' <i>Pilodius</i> ' <i>vulgaris</i>	+	+	-
<i>Xantho moldavicus</i>	+	+	+
<i>Pilumnus mediterraneus</i>	+	?	+
<i>Pilumnopeus paralethynensis</i>	-	-	+
<i>Haydnella steiningeri</i>	+	+	-
<i>Chlorodiella juglans</i>	-	-	+
<i>Chlorodiella mediterranea</i>	+	+	+
<i>Chlorodiella mediterranea tetenyensis</i>	+	+	+
<i>Actaea turcocampestris</i>	+	?	-
<i>Carpilius antiquus</i>	+	?	-
<i>Daira speciosa</i>	+	+	+
<i>Trapezia glaessneri</i>	+	-	-

Group 1: Lower Badenian sites in the Börzsöny- and Pilis Mountains

Group 2: Gross-Höflein, "Oberer Korallenkalk"

Group 3: Upper Badenian localities in Budapest and Diósd

Table 3

## Tentative correlation of decapod zones to substages of Badenian

Decapod zones	Substages
4	
3	Upper Badenian
2	Middle Badenian
1	Lower Badenian

The presence of a fourth uppermost zone is evident in the Upper Badenian sequence in Budapest and Balatonakali. This zone is characterized by the absence of many species, extremely common in layers contemporary with the zone 3, i.e. *Callianassa munieri*, *C. chalmasii*, *Petrochirus priscus*, *Dromia eotvoesi*, *Calappa heberti*, *Thalamita fragilis* and all reef-bound species. Reefs are no longer present in this uppermost Badenian strata. The lower layers (indicated in section on Figs. 4, 5 as zone 3, i.e. MR8, MRO, MRL, MGY, MR45, MA etc.) are characterized by their rock rich in terrigenous and vulcanogenous material: gravel, sand, tefra, pumice. These layers are also characterized by intercalated lenses of reefs belonging to zone 3 (MRZ, MK, MDZ). They are covered by a series of layers virtually depleted of terrigenous material thus consisting of pure limestones (indicated on Figs. 4, 5 as zone 4, MRS, MOE, MB, MDG etc.).

The extinction of several decapods and the abrupt change in sedimentological character were contemporaneous and might be connected with climatic changes. Semi-arid conditions might cause fast erosion resulting in a high rate of terrigenous material in sediments. In the subsequent humid climate dense forests might cover the slopes and hinder erosion. A connected slight change in salinity and decreasing temperature of seawater might be responsible for the extinction of corals and some decapods.

The four zones tentatively characterized by presence or absence of species of decapods, could represent assemblage zones. Future studies will have to clarify whether these are general in the Central Paratethys or reflect only local patterns. Based on the above, decapod zones can be correlated to substages of Badenian as shown in Table 3.

Numerous decapod genera common in Badenian are recently confined to the Indo-West-Pacific realm. This fact was stressed by GLAESNER as early as 1928 (p. 205) and subsequently confirmed by others (FÖRSTER 1979a, p. 104, MÜLLER 1979c, p. 865). The Miocene decapods of the Indo-West-Pacific realm are poorly known making a comparison of Miocene faunas of that area and of the Paratethys impossible. The presence in the Badenian of genera which are now confined to the Indo-West-Pacific can easily be attributed to their post-Miocene extinction in the Mediterranean and in the eastern Atlantic subsequent to their disappearance from the Paratethys at the end of the Badenian. The extinction of coral-reef-bound species was inevitable in these waters with the disappearance of hermatypic corals. It is well known that the deterioration of the climate at the end of Tertiary was not so severe in South Asia than in Europe or in the Atlantic making the Indian and Pacific Oceans a natural refuge.

Genera, abundant in the Badenian, but recently confined to the Indo-West-Pacific are: *Matuta*, *Micippa*, *Macrophthalmus*, *Crossotonotus*, *Schizophrys* and *Actumnus* (some of them extended to a restricted area in the southeastern Atlantic, too). Some genera live elsewhere, too, but are predominantly Indo-West-Pacific, e.g. *Mursia*, *Trapezia*, *Daira*, *Chlorodiella*, *Carpilius*, *Dorippe*, *Pilumnopeus* and *Achaeus*. Genera determined with some uncertainty, e.g. *Pilodius*, *Glabropilumnus*, were not included in the above lists. Of the first list (six genera) all six occur in the Upper, while three or four in the Lower Badenian. From the second list (8 genera) six are Upper, and four Lower Badenian, while a genus is represented in older layers of Western Europe, too (*Mursia*). In contrast to earlier suggestions (MÜLLER 1979c), no or only slight temporal increase can be discerned in the proportion of Indo-West-Pacific-related forms, if we consider that the total number of known Upper Badenian genera is higher (53) than that of Lower Badenian ones (36).

Three species require special consideration. These, or very closely allied forms occur in the Pliocene of Fiji (RATHBUN 1945). Two of these (*Dardanus hungaricus* and *Daira speciosa*) occurred in zones 1 and 3 (Table 3) while *Dynomene emiliae* was found in zones 1 and 2 only. Thus the number of these species, presumably most closely related to Indo-West-Pacific forms, seems to decrease with time.

Consequently the forms with eastern relationships derived most probably from a common Tethyan fauna and subsequently vanished from European seas. Thus a connection of the Badenian sea with the Indian Ocean seems less probable now than some years ago. Nevertheless the existence of a sea-way permitting migration of certain euryhaline elements cannot be ruled out. The term euryhaline must be stressed, as such a way might have existed through the Eastern Paratethys to a sea somewhere in the Middle-East (Upper Fars series?). The salinity of the former sea was most probably inferior, and of the latter superior to that of the Oceans (RÖGL-STEININGER-C. MÜLLER 1978; STEININGER, RÖGL and C. MÜLLER in PAPP et al. 1978). Neither can another, though less probable, connection be ruled out along some unknown, partly molasse-like basins along the Hellenids, Taurids and Zagros, but our present knowledge of Miocene seas of these areas is highly controversial.

One point seems to be indisputable, however. The general character of the fauna remained constant for the whole Badenian thus the connection ensuring the oceanic (or nearly oceanic) salinity, i.e. the connection with the Mediterranean, also must have remained unchanged. At least four Badenian species are identical, while six others are almost identical with Mediterranean Miocene species.

A remarkable fact still awaits its interpretation. Certain highly abundant species are present exclusively in layers of the zone 3 (Table 3): *Petrochirus priscus*, *Dromia eotvoesi*, *Calappa heberti*, *Matuta brocchii*, *Micippa hungarica*, *Thalamita fragilis*. The abundant occurrence of these suggest that most of them are indeed absent from older strata. The majority of them vanished at the beginning of zone 4, others, as *Matuta*, are rarer in upper layers than in zone 3. One possible explanation is that these species were highly dependent on climate and/or on salinity. This is consistent with the supposed climatic change at the boundary of zones 3 and 4, causing regression or extinction of sen-

sitive, probably stenohaline species. Their probable immigration at the beginning of zone 3 requires the assumption of a source for these species. An Indo-West-Pacific source seems highly improbable as stenohaline species could hardly penetrate seas of abnormal salinity (cf. above). Perhaps these forms were present in southern parts of the Mediterranean during Early Badenian times but so far only *Petrochirus priscus* was recorded from this area (GLAESSNER 1928).

Various decapod sites of the Badenian yielded different types of associations which can be grouped even on a subjective basis. The clusters obtained in this way might reflect ecological conditions. To achieve a more objective clustering, an analytical investigation was carried out by dr. M. RAJCZY, Natural History Museum of Budapest. The similarity of individual associations (layers in our case) was computed by binary (absence vs. presence) and quantitative methods. Quantitative methods are used infrequently in macropaleontological investigations, especially for hand-picked samples, since the representative nature of such a sample can be questioned. Certain considerations (see below) and our rather convincing results support the reliability of quantitative evaluation of such collections, however. In hand-sampled collections for obvious reasons the sample will be distorted statistically. The rock is fragmented by the collector into pieces of a given average size. Specimens smaller than these pieces will remain hidden if located inside. Assuming that fragmentation is statistically independent of the position of fossils within the matrix, the probability that a fossil of given size will be observed:

$$p = 1 - \frac{(r-f)^3}{r^3} \quad (p_{\max} = 1)$$

where  $p$  is the probability of detection of a fossil of size  $f$ , and  $r$  is the average size of the pieces of fragmented rock. This formula was used to provide comparable distributive functions of sieved and of hand-picked collections. Although one cause of distortion was thus reduced, others, due to selective embedding, fossilization, incorrect determination etc., still remained. The data, corrected in this way, were used for computing similarities by binary (SØRENSEN 1948, SOKAL-MICHENER 1958) and by quantitative (HORN 1966, CZEKANOWSKI 1909, PINKHAM-PEARSON 1976, HUMMON 1974) functions. For clustering, the method of weighted averages was used. Calculations were based either on quantity of specimens of genera or of species. The results obtained by either approach were similar since few genera contained more than one species.

The main goal of the cluster-analysis was to define sets of individual sites and layers which can be studied subsequently together as groups. The most fidel and the most constant species of such groups can be discerned and used in further studies. Evidently such forms will provide more reliable basis for ecological considerations than those found in one or few sites only. The groups or clusters might be obtained from dendograms by cutting them on a given level of similarity. The choice of this level is arbitrary to some extent thus the method is inevitably subjective. The subjectivity will be reduced if different methods give similar results. The Badenian decapods yielded four groups based on functions of SØRENSEN, HUMMON, SOKAL-MICHENER and HORN and five more or less homogeneous clusters based on dendograms prepared according to the methods of PINKHAM-PEARSON and CZEKANOWSKI (Fig. 12, Table 4). Further levels of hierarchy were not evident. Localities of reef-origin formed a coherent cluster. The PINKHAM-PEARSON and SOKAL-MICHENER functions separated this group into two parts, however. The clustering by four different methods based on functions of SØRENSEN, CZEKANOWSKI, HORN and HUMMON gave almost identical results for most sites. The position of some layers, however, varied according the function used. These are: limestone at Kerepesi út (MKC), Węglinek (WEG) and one of the Diósd localities (MDN). The first seems transitional in many respects, while the two others are poorly collected. The results were consistent with the main petrologic character of rocks, i.e. presence or absence of oncoids or reefal structures. The groups were designated according to one of their constituents and an attempt will be made to evaluate the conditions under which the layers of these groups were formed (Table 4).

The most coherent cluster (MGY) includes layers MRO, MRF, MR8, MR45, MA, MOH, MEG, MFA and in certain dendograms MFK. Mátraverebély-Szentkút and Nyírád (Lower Badenian localities, MMS resp. MN) yielded similar associations, too with *Calappa praelata* replacing *C.*









Fig. 12. Dendograms showing similarities of species assemblages based on cluster-analysis. [Codes of assemblages (layers) are explained on pages 36—47]

Table 4

## Clusters of decapod localities

Name	Sites of layers	Abundant and constant forms	Occasional forms	Inferred conditions in habitat	Matrix
MGY	MGY, MRF, MRS, MRO, MR45, MA, MOH, MEG, MFA, (MFK)* (Métraverebélly-Szentháromság), (MNU), (MOS)	<i>Calappa</i> <i>Matuta</i> <i>Mursia</i> <i>Thalamita</i> <i>Dromia eobvoesi</i> <i>Portunus (Portunus)</i>	<i>Albunea</i> <i>Portunus (Monomia)</i> <i>Moja</i> <i>Dorippe</i>	Depth: 20–30 m ± oceanic salinity Loose substrate Warm-temperate climate	Sandstone, sandy limestone
MOE	MOE, MDG, MDN, (DAL), (WEG)	<i>Micippa</i> <i>Ebalia</i> <i>'Lissocarcinus'</i> <i>Liocarcinus</i> <i>Achelous</i> <i>Parthenope</i>	<i>Meja</i> <i>Cancer</i> <i>Dorippe</i> <i>Palicus</i> <i>Palaeomyra</i>	Depth > 20 m Slightly changing salinity Inhomogeneous, partly hard substrate Temperate climate	Fine or coarse calcarenite without oncoids
MB	MB, MRS, MGF, MBK, MBH, MJL, MM, (DAL)	<i>Picidina</i> <i>'Pis'a'</i> <i>Pachygrapsus</i> <i>Brachynotus</i> <i>Pilumnus</i>	Xantho	Depth 0–2 m Changing salinity Inhomogeneous, partly hard substrate, oncoids, crevices Temperate climate	Coarse calcarenite with oncoids or stromatolites
MRZ	MRZ, MDZ, MK, MV, MTZ, MNH, (MZZ)	<i>Chlorodiella</i> <i>Datra</i> <i>Charybdis (Goniozupradens)</i> <i>Pilumnopodus</i> <i>Petrolithes</i> <i>Galahea</i> <i>Dromia neogenica</i>	<i>Carpilius</i> <i>Panopus</i> <i>Cancer</i> <i>Pachygrapsus</i> <i>Crossollenatus</i>	Depth < 10 m ± oceanic salinity Patch-reefs Warm-temperate climate	Corals in rock-building quantity in various matrices (marl, sandstone, calcarenite)
SOOS	Soos, (Budapest drilling Z-2), (Walbersdorf), (Oriahovo)		Ranina Geryon Gonoplax "Cancer" szontaghii bonensis	Depth > 50 m ± oceanic salinity Soft, muddy substrate	Silt, clayey silt, rarely calcarenite

\* Designation within brackets: were assigned here only in certain dendograms, or assigned here subjectively.

*heberti*, but the low number of species found did not allow their inclusion in the numerical study. These associations are characterized by the abundance of *Calappa*, *Matuta*, *Mursia*, *Thalamita*, *Dromia eotvoesi* and *Portunus* (*Portunus*), while *Albunea* and *Portunus* (*Monomia*) were found in few sites only. These genera are rare or absent in other clusters thus they are fidel. The first five are constant as they are present in virtually all localities of this group.

The high diversity of 'Callianassa' forms (SAINT LAURENT-LE LOEFF 1979) can be attributed to a depth of 10 to 30 meter, as most extant *Callianassa* and *Callichirus* species prefer such conditions, though few species live intertidally, while some others descend below 100 meter. In some sites the presence of big species (*C. munieri*) indicates homogeneous sands on bottom. Most *Matuta* species live in shallow waters (SAKAI 1976), but *M. curtispina*, morphologically similar to *M. brocchii*, live on sands in depth of 35 to 80 meter. *Mursia* species prefer soft bottom in depth of 50 to 150 meter (SAKAI 1976), while *M. cristimana* was reported from shallow waters, too (17 m. BARNARD 1950). *Thalamita crenata* (similar to *Th. fragilis*) lives in extremely shallow water, on mud or soft sand. Other *Th.* species descend to 80 meter (SAKAI 1976). *Calappa* species live from 10 to 150 meter. Preference of depth depends on species (SAKAI 1976). *C. granulata*, similar to *C. praelata*, lives in depth of 30 to 150 meter (ZARIQUIEY ALVAREZ 1968). *Portunus* (*Portunus*) species live in shallow water, down to 20 or 30 meter. *Albunea carabus* lives on soft sandy bottom, in depth of 30 to 50 meter generally (ZARIQUIEY ALVAREZ 1968).

The actualistic interpretation of the mentioned data suggest a depth of 20 to 30 meter for most localities of this group. Where the sediments were homogeneous (MRF, MR8, MR45), big callianassids were abundant, in other places the smaller 'Callianassa' *pseudorakosensis* lived in masses (MFA, MFK). Abundance and diversity of calappid species suggest euhaline conditions and a warm-temperate climate. The occasional abundance of *Albunea* (MGY, MRF) may suggest water slightly deeper than 30 meter. The localities of this cluster share a set of species with other groups, mostly with MOE, as discussed below. The most important of these are *Maja*, with a preference for depth of 15 to 30 meter, or rarely of 10 to 200 meter (SAKAI 1976). *Maja squinado*, a European species, lives in depth of 2 to 73 meter (INGLE 1980). Extant *Dorippe* species live in depth of 15 to 50 meter (SAKAI 1976) or of 40 to 100 meter (ZARIQUIEY ALVAREZ 1968). Various *Liocarcinus* species live in different zones from the shoreline down to 220 meter (INGLE 1980).

Another cluster, MOE, includes also MDG, MDN, and in part of classifications, Deutsch-Altenburg and Węglerek as well. The sites of this group are characterized by abundance of *Micippa hungarica*, *Ebalia* species, 'Lissocarcinus', *Liocarcinus*, and in cases of better known localities, by the presence of *Achaeus*, *Actumnus*, *Dorippe*, *Maja*, *Parthenope*, *Cancer*, *Xantho*, *Palicus*. *Micippa thalia*, extant species, which is closely related to *M. hungarica*, lives in depth of 20 to 100 meter, on sand or broken shells (SAKAI 1976). Most *Achaeus* forms live between 10 to 50 meter, but some of them live in shallower habitats, too. *Ebalia* species live in various depths, they are reported from two to several hundreds of meters (INGLE 1980). *Palicus* species live mostly below 50 meter (SAKAI 1976, ZARIQUIEY ALVAREZ 1968). *Cancer* forms live in various depth (RATHBUN 1930), from the shore down to 1000 meter. *Parthenope* species are common between 10 to 40 meter (ZARIQUIEY ALVAREZ 1968). Extant *Actumnus* species live generally in deeper environments than *Pilumnus* forms, generally below 20 or 35 meter, but two forms occur intertidally as well (SAKAI 1976).

These data suggest a depth slightly greater than 20 meter similarly to the case of group MGY. One difference is the lack of big callianassids. This can be attributed to inhomogeneous sediments. Lack of *Calappa*, *Dromia*, *Thalamita*, *Portunus*, rarity of *Matuta*, *Mursia*, genera predominantly occurring under warm-temperate or tropical climates and in euhaline water, suggest climatic deterioration and slightly changing salinity of sea-water. As it was stated (p. 104), the MOE localities are characterized by lack of terrigenous sediments, probably related to a change of climate at the boundary of zones 3 resp. 4 (p. 105). Molluscs, studied by KÓKAY (KÓKAY-MIHÁLY-MÜLLER in press) suggest also a sea deeper than 20 or 30 meter.

Probably the most interesting cluster is MB as it represents an environment rarely preserved. Petrologically, its rocks contain oncoids, stromatolites or similar structures (LELKES-MÜLLER in press), suggesting extremely shallow environments. The group contains the following sites or layers: MRS, MGF, MB, MBK, MBH, MJL, MM and with certain reservations two additional ones, Węglerek and Deutsch-Altenburg. Faunistically, it is characterized by the abundance of *Pisidia kokayi*, 'Pisa' *oroszyi* and *Pachygrapsus*, which all are rare or absent elsewhere, by the exclusive presence of *Brachynotus februarius*, and by abundance of *Pilumnus mediterraneus* and *Xantho moldavicus*. *Pachygrapsus* species are typical eulittoral crabs living on rocky shores (PÉRÈS-PICARD 1964) which makes their occurrence particularly interesting. *Brachynotus* forms live mostly between 5 to 2 meter of depth (ZARIQUIEY ALVAREZ 1968). Most *Pisidia* species also live in such habitats but some occur on deeper bottoms as well. The last three forms prefer bottoms rich in shelters. *Pisa* species live in depth between one to 90 meter, prefer hard grounds (INGLE 1980). *Xantho* and *Pilumnus* forms are rather ubiquitous and also occur in extremely shallow water.

According to these data MB type layers were deposited in extremely shallow shore regions. Oncoids and stromatolites provided hard ground offering shelters while temporary regressions resulted in rocky intertidal conditions on shores covered with oncoids. Extant relatives of most forms are euryhaline organisms, as *Xantho*, *Pisidia*, *Pachygrapsus*, *Liocarcinus* and *Brachynotus* species live in the Black Sea (PESTA 1918), while some leucosids live either in the Marmara Sea (HOLTHUIS-GOTTLIEB 1958, p. 80), or crossed the hyperhaline barrier of the Suez Canal (POR 1978). This fact is consistent with changing salinities in high sublittoral and eulittoral environments.

The group MRZ representing coral-reef habitats is characterized by the nature of rock-containing hermatypic corals in rock-building quantity preserved either in living position or as removed blocks embedded in matrix. A series of genera or species is restricted to these sites: all species of *Chlorodiella*, *Daira speciosa*, *Carpilius*, *Charybdis* (*Goniosupradens*), *Dynomene*, *Dromia neogenica*, *Actaea*, 'Polidius', *Haydnella*, *Crossotonotus*, *Schizophrys*, *Petrolisthes*. Most of their extant relatives are reef-dwellers, as *Charybdis* (*Goniosupradens*), *Daira perlata*, *Chlorodiella*, most *Dynomene* species, *Carpilius*, *Trapezia* (SAKAI 1976, TAYLOR 1968). Others are not exclusively restricted to reefs but prefer hard bottom thus occasionally occur on reefs, too (*Schizophrys*, *Carupa*, *Daira americana*, *Crossotonotus*). Unexplained is the presence of *Panopeus* and *Cancer* in sites of reef-origin. *Panopeus wronai* occurs in various sites while *Cancer* only in Diósd (MDZ). Extant *Panopeus* species occur occasionally on reefs but prefer soft bottom while *Cancer* species were never found on reefs (RATHBUN 1930). *Calappa heberti*, though rarely, occurs also on reefs similarly to some extant *C.* species. Few specimens of *Matuta* were also found in rocks of reef-origin (MRZ). Characteristic is the consistent absence of some crabs abundant elsewhere, as *Parthenope*, leucosids, *Mursia* in agreement with the fact that these are absent or rare on extant reefs (TAYLOR 1968). In two of these sites (MDZ and MRZ) *Pisidia kokayi* and *Pachygrapsus hungaricus* occur indicating temporary emersions as proven also by sedimentological features (root-prints, desiccation cracks). The presence of coral-reefs points to euhaline water and to warm-temperate or tropical climate. The decrease of number of coral-genera (Table 3) with time suggests slight perturbations in salinity in upper parts of Badenian.

An association of a peculiar facies recently found in Žebegény (MZZ) (Fig. 6) was not yet included in the cluster-analysis. A fine-grained micritic matrix includes plentiful colonies of a small ahermatypic branching colonial coral-species similar to *Oculina*. Fields of ahermatypic corals may be found at different depths from one to several hundreds of meters, both in tropical or cold-temperate seas. Scarce presence of ?*Porites* indicates, however, a depth less than a few tens of meters. The decapods found are similar but less diverse than those found in reef-limestones. Interesting is the presence of two leucosid species, since leucosids are absent from Badenian reefs. Portunids and dromids are also absent from the sample taken comprising 30 specimens.

The existence of a fifth type of habitat is indicated from various localities, most of which did not yield more than one or two forms determinable to species level. Two localities yielded *Goneplax* species. In Oriakhovo it was found together with *Miocyclus bulgaricus*, in Sooss at Baden with "Cancer" szontaghii vindobonensis. A *Geryon* species was found in drilling Z-2. Budapest. *Goneplax* (40 to 760 meter, PESTA 1918), and *Geryon* (776 to 1500 meter, PESTA 1918) live in deeper sublittoral and bathyal environments. *Ranidina rosaliae* was found in two sites, in Walbersdorf (Borbolya) it occurred together with "Cancer" cf. *bittneri*. Decapods of deep sublittoral environments with soft bottom are thus known from sporadic findings.

The results presented above are in accord with those obtained by analysis of invertebrates other than decapods and by sedimentological methods (Table 4). Preliminary results of such studies are presented in various publications (MÜLLER 1974a, 1974b, 1976a, 1979a; KÓKAY-MIHÁLY-MÜLLER in press; LELKES-MÜLLER in press).

Shannon diversity indices were calculated for the samples analysed above, also by M. RAJCZY. The extreme values, expressed as  $H' \log_2$  for groups of sites are as follows:

- Group MGY:  $H' = 2.18-3.66$
- Group MOE:  $H' = 3.25-4.00$
- Group MRZ:  $H' = 2.10-3.63$
- Group MB:  $H' = 1.24-2.83$

The diversity of some intertidal and shallow subtidal crab-assotiations was studied by ABELE (1974). Though his samples were taken from warm-temperate and tropical seas of higher temperatures than those suggested for Badenian seas, his highest value,  $H' \log_2 = 4.346$  does not exceed significantly our highest values. Fossil samples might also contain specimens from more than one habitat which could increase diversity. Whether this effect was significant in Badenian localities cannot be a priori decided. Nevertheless a post-mortem transport of decapod remnants is less likely than that of most fossil invertebrates.

An interesting pattern of diversity was revealed along a section on the Tétény-plateau. For localities of the group MB (Fig. 2) diversity clearly increased from layer MM, presumably situated

closest to the shore, toward south, i. e. the open sea, from a value of  $H' \log_2 = 1.24$  through  $H' = 2.93$  (MBH) to  $H' = 3.25$ , value of site MDG. Since these layers were deposited more or less simultaneously, it is likely that the observed increase of  $H'$  reflects increasing depth in a section normal to the shore. This result is somewhat in disagreement with ABELE (1974) who found some of the highest values in rocky intertidal habitats, while such a habitat (MM) yielded our lowest value. ABELE did investigate only one subtidal community, however, namely a *Pocillopora* association of high diversity,  $H' \log_2 = 4.346$ . If the results concerning Badenian associations are correct (at least at a level limited by the restricted size of samples), not the values for coral-reef (MRZ) associations are the highest ones (2.10–3.63), but those for sublittoral communities which lived on a complex bottom with shells, calcareous sand, oyster-beds (MOE,  $H' \log_2 = 4.00$ ). ABELE concluded (1974) that diversity correlated best with complexity of the substrate. In this respect our results are in agreement with ABELE, as a sublittoral environment like MOE may compete even with a coral-reef being similarly complex.

## ADDENDUM

After closing this manuscript I had the opportunity to study some material in collections in Vienna and make additional collections at Austrian localities. This has led to some taxonomical changes.

Study of the type-series of *Cancer carniolicus* BITTNER, 1884 (kept in the Naturhistorisches Museum, lectotype designated herein: pl. 1 fig. 8a in BITTNER, 1884, seen by the courtesy of dr. BACHMAYER) and that of some new specimens of *Cancer szontaghii* LÖRENTHEY, 1897 from its type-locality, demonstrated convincingly their identity. *Cancer bittneri* TOULA, 1904 (holotype kept also in the NHMW), though its carapace is slightly smoother and wider than that of BITTNER's and LÖRENTHEY's specimens, belongs also here. In JANSSEN—MÜLLER (in press) MÜLLER described a new monotypical genus, *Tasadia*, for *Cancer szontaghii* LÖRENTHEY (here mentioned as nomen nudum). Thus the name *Tasadia carniolica* (BITTNER, 1884) should be used for all of the mentioned forms after the appearance of JANSSEN—MÜLLER (in press).

Seven new specimens of *Atelecyclus szontaghii vindobonensis* BACHMAYER—KÜPPER, 1952 were collected at the type-locality (Baden-Sooss, clay-pit, leg. P. PERVESLER and me). This form, though having many features in common with *Tasadia carniolica* (outline of the carapace, presence of tuberculated elevations on the main regions), differs from it by the much smaller relative size of these elevations and by the much smoother inter-elevation surface, as it was mentioned by its author. Thus the subspecies should be maintained under the name *Tasadia carniolica vindobonensis*.

Poorly preserved specimens from Walbersdorf (coming from a layer of Lower Badenian age, RÖGL, pers. comm.), kept at the Geologische Bundesanstalt and in the NHMW (mentioned by BITTNER, 1893 p. 32 as *Cancer* cfr. *illyricus* and by GLAESSNER, 1928 p. 178 as *Cancer bittneri*) belong either to *Tasadia carniolica* (BITTNER) or are transitional between the nominal subspecies and ssp. *vindobonensis*.

The type-series (10 specimens) of *Ranidina rosaliae* BITTNER, 1893 is deposited in the collection of the Geologische Bundesanstalt. Lectotype, designated herein, pl. 2 fig. 2 (non fig. 2a) in BITTNER (1893).

## REFERENCES

- ABELE L. G. 1974: Species diversity of Decapod Crustaceans in marine habitats. — Ecology. 55 (1): 156—161.
- BACHMAYER F. 1950: Neue Dekapoden aus dem Österreichischen Tertiär. — Ann. Naturhist. Mus. Wien. 57: 133—140. pl. 1.
- BACHMAYER F. 1953a: Die Dekapodenfauna des tortonischen Leithakalkes von Deutsch-Altenburg (Niederösterreich). — Mitt. Geol. Ges. Wien. 44: 237—262. pl. 1—5.
- BACHMAYER F. 1953b: *Goneplax gulderi*, eine neue Crustaceen-Species aus dem tortonischen Tegel des Wiener Beckens. — Paläont. Zeitschr. 27 (3—4): 143—145. pl. 9.
- BACHMAYER F. 1953c: Zwei neue Crustaceen-Arten aus dem Helvet-Schlier von Ottwang. — Ann. Naturhist. Mus. Wien. 59: 137—140. pl. 5.
- BACHMAYER F. 1954: Zwei bemerkenswerte Crustaceen-Funde aus dem Jungtertiär des Wiener Beckens. — Österr. Akad. Wiss. Sitzungsber. Math.-Naturwiss. Kl. Abt. I. 163 (1—2): 63—70. pl. 1.
- BACHMAYER F. 1962: Die Calappiden (Crustacea, Decapoda) aus dem tortonischen Ablagerungen des Wiener Beckens. — Ann. Naturhist. Mus. Wien. 65: 39—46. pl. 1—3.
- BACHMAYER F.—KÜPPER K. 1952: Eine bemerkenswerte Krabbe aus dem Badener Tegel (Torton) von Sooss im Wiener Becken. — Verh. Geol. Bundesanst. Wien. (4): 200—205. text-fig. 1, 2.
- BACHMAYER F.—TOLLMANN A. 1953: Die Crustaceen-Fauna aus dem tortonischen Leithakalk (Steinbrüche der Firma Fenk) bei Gross-Höflein im Burgenland. — Kober-Festschrift: 309—314. pl. 1. Wien.
- BÁLDI T.—KÓKAY J. 1970: A kismarosi tufit faunája és a börzsönyi andezitvulkánosság. — Földt. Közl. 100 (4): 274—284.
- BÁLDINÉ BEKE M.—BOHNNÉ HAVAS M.—KORECZNIÉ LAKY I.—NAGYNÉ GELLAI Á.—NAGY LÁSZLÓNÉ 1980: Újabb őslénytani és rétegtani eredmények a Börzsöny-hegység és távolabbi környékének oligocénjéből és miocénjéből (Recent paleontological and stratigraphical results on the Oligocene and Miocene of the Börzsöny Mountain and its surroundings). — Őslénytani Viták. 26: 61—102.
- BALOGH KADOSA—ÁRVÁNÉ SÓS E.—PÉCSKAY Z. 1980: Hazai kőzetek K/Ar kora. — Atommag Kutató Intézet, kézirat, Debrecen.
- BALSS H. 1933: Beiträge zur Kenntniss der Gattung *Pilumnus* (Crustacea Decapoda) und verwandte Gattungen. — Capita Zoologica. 4 (3): 1—47. pl. 1—7.
- BARNARD K. H. 1950: Descriptive catalogue of South African Decapod Crustaceans. — Ann. South Afr. Mus. 38: 1—837. text-figs. 1—154.
- BARTKÓ L.—KÓKAY J. 1966: Lajtamészkkő előfordulás a Kerepesi úton (Leithakalksteinvorkommen in Budapest Kerepeser Strasse). — Földt. Közl. 96 (3): 301—305.
- BEURLEN K. 1939: Neue Decapoden-Krebse aus dem ungarischen Tertiär. — Pal. Zeitschr. 21 (2): 135—160. pl. 7. text-figs. 1—5.
- BEURLEN K. 1958: Contribuição a Paleontologia do Estado do Para, Crustáceos decápodes da Formação Pirabas. — Bol. Mus. Paraense Emílio Goeldi, Nova Serie, Geologia. 5: 2—48. pl. 1—4.
- BITTNER A. 1877: Über *Phymatocarcinus speciosus* Reuss. — Sitzungsber. Math. Nat. Cl. Akad. Wiss. Wien. I. Abt. 75: 435—447. pl. 1.
- BITTNER A. 1884: Beiträge zur Kenntniss tertärer Brachyuren-Faunen. — Denkschr. Math. Nat. Cl. Akad. Wiss. Wien. I. Abt. 48: 15—30. pl. 1, 2.
- BITTNER A. 1893: Decapoden des pannonischen Tertiärs. — Sitzungsber. Math. Nat. Cl. Akad. Wiss. Wien. I. Abt. 102: 10—37. pl. 1, 2.
- BONFIGLIO L.—DONADEO G. 1982: Cancer sismondai Meyer nel Pliocene di Torre dell'Orso (Puglia). — Atti Soc. Ital. Sci. Nat. Mus. Civic. Stor. Nat. Milano. 123 (2—3): 255—296. pl. 36—44, text-figs. 1—27.
- BOUVIER E. L. 1940: Décapodes marcheurs. In: Faune de France. — 37: 1—399. pl. 1—14, text-figs. 1—222.
- BROCCHI P. 1883: Note sur les Crustacés Fossiles des Terrains tertiaires de la Hongrie. — Ann. Sci. Geol. Paris. 14: 1—8. pl. 4, 5.
- CAMPBELL B. M.—STEPHENSON W. 1970: The sublittoral Brachyura (Crustacea Decapoda) of Moreton Bay. — Mem. Quensld. Mus. 15 (4): 235—301. pl. 22, text-figs. 1—49.
- CHADIDA M. R.—PAPP A. 1977: Verbreitung von Oberoligozän und Untermiozän im Zentral-Iran und dessen faunistisch-paläogeographische Beziehung zur Paratethys. — N. Jb. Geol. Paleont. (7): 402—406.
- COUFFON O. 1908: Sur quelques Crustacés des Faluns de Touraine et d'Anjou, suivi d'un essai de Prodrome des Crustacés podophtalmaires miocènes. — Feuille des Jeunes Naturalistes. 4 (457): 1—40. pl. 1, 2. Paris.
- COUFFON O. 1910: *Maia Orbigniana* Millet. In: Palaeont. Universalis: 130.

- CZEKANOWSKI J. 1909: Zur Differenzial Diagnose der Neanderthalgruppe. — Korresp. Bl. dt. Ges. Anthropol. 40: 44—47.
- DOFLEIN F. 1904: Brachyura. — In: Wissenschaftliche Ergebnisse der deutschen Tiefsee-Expedition, Valdivia. 6: I—XIV, 1—314, text-figs. 1—314, pl. 1—57.
- DULLO W. CH. 1983: Die miozäne Leithakalk der zentralen Paratethys. — Facies. 8: 1—112. pl. 1—12, text-figs. 1—22.
- FÖRSTER R. 1979a: Decapod Crustaceans from the Middle Miocene (Badenian) deposits of Southern Poland. — Acta Geol. Pol. 29 (1): 89—106. pl. 1—4.
- FÖRSTER R. 1979b: Decapod Crustaceans from the Korytnica basin (Middle Miocene; Holy Cross Mountains, Central Poland). — Acta Geol. Pol. 29 (3): 253—268. pl. 1—5.
- FRANZENAU A. 1881: Adatok a rákosi (Budapest) felső mediterrán-emelet Foraminifera-faunájához [Beitrag zur Foraminiferen Fauna der Rákoser (Budapest) Ober-Mediterran Stufe]. — Földt. Kőzl. 11: 31—107.
- GALOPIM DA CARVALHO A. M. 1959: Malacostráceos das formações glauconiosas do Miocénico superior do litoral a norte do Cabo Espichel. — Bol. Mus. Lab. Miner. Faculd. Ciencias. Lisboa. 8 (1): 77—82. pl. 1, 2.
- GARTH J. S. 1974: On the occurrence in the Eastern Tropical Pacific of Indo-West-Pacific Decapod Crustaceans commensal with reef-building corals. — In: Proc. II. Intern. Coral Reef Symp. 1: 397—404. Brisbane.
- GEORGIADES-DIKEOULIA E. 1977: Présence de Cancer sismondai H. v. Meyer dans les couches Pliocènes d'Heraklion (Crète). — Ann. Geol. Pays Hell. 28: 418—426.
- GLAESSNER M. 1924: Über eine neue miozäne Krabbe und die Brachyurenfauna des Wiener Beckens. — Verh. Geol. Bundesanst. Wien. (6): 109—118. text-figs. 1, 2.
- GLAESSNER M. 1928: Die Dekapodenfauna des Österreichischen Jungtertiärs. — Jahrb. Geol. Bundesanst. Wien. 78: 161—219. pl. 3, 4.
- GLAESSNER M. 1929: Crustacea Decapoda. — Fossilium Catalogus. 41: 1—464. Berlin.
- GLAESSNER M. 1930: Dekapodenstudien. — Neues Jahrb. Geol. Paleont. 63 (Beilage-Band), Abt. B: 137—176. pl. 6—10.
- GLAESSNER M. F. 1969: Decapoda. — Treatise on Invertebrate Paleontology. R. Arthropoda. 4: 309—533.
- GRIPP K. 1964a: Erdgeschichte Schleswig-Holsteins. — Neumünster.
- GRIPP K. 1964b: Krebse aus Miozän-Geschieben. — Lauenburgische Heimat. Ratzeburg. Neue Folge. 45: 55—57.
- GRIPP K. 1967: Dekapode Krebse tertiären Alters aus Schleswig-Holstein. — Meyniana. 17: 1—3. pl. 1, text-fig. 1.
- GRIPP K. 1968: Neue Funde aus dem Miozän und Pliozän von Amrum und Sylt. — Meyniana. 18: 1—8. pl. 1—3.
- GRIPP K. 1969: Fossilien aus Norddeutschland. — Meyniana. 19: 79—89. pl. 1—3.
- GUINOT D. 1967: Recherches préliminaires sur les groupements naturels chez les Crustacés Décapodes Brachyoures. III: A propos des affinités des genres Dairodes Stebbing et Daira de Haan. — Bull. Mus. Natl. Hist. Nat. Paris. 2<sup>e</sup> 39 (3): 540—563.
- GUINOT D. 1968: Recherches préliminaires sur les groupements naturels chez les Crustacés Décapodes Brachyoures. VI: Les Carpiliinae. — Bull. Mus. Natl. Hist. Nat. Paris. 2<sup>e</sup> 40 (2): 320—334.
- GUINOT D. 1969: Sur divers Xanthidae, notamment sur Actaea de Haan et Paractaea gen. nov. (Crustacea Decapoda Brachyura). — Cahiers du Pacifique. 13: 223—269. text-figs. 1—36.
- GUINOT D. 1977: Propositions pour une nouvelle classification des Crustacés Décapodes Brachyoures. — C. R. hebd. Séances Ac. Sci. Paris. 285 (D/10): 1049—1053.
- HAIG J. 1981: Three new species of Petrolisthes (Decapoda Anomura Porcellanidae) from the Indo-West Pacific. — Journ. Crust. Biol. 1 (2): 265—271. text-figs. 1—3.
- HÁMOR G. 1985: A Nógrád—Cserhát kutatási terület földtani viszonyai. — Geol. Hung. ser. Geol. 22 (in press).
- HOJNOS R. 1923: Adatok a Kis-Zagyva környékének földtani ismeretéhez. — 1—8. Budapest.
- HOJNOS R. 1933: Über fossilen Krebsbruchteile der tuffitische Sande des Cserháts. — In: A Mátra gazdasággeológiai jelentősége. Budapest.
- HOLTHUIS L. B. 1949: Fossil Decapod Crustacea from Miocene and younger deposits of the Netherlands. — Meded. Geol. Stichting, Nieuwe Ser. 3: 55—68. pl. 1, 2.
- HOLTHUIS L. B. 1959: The Crustacea Decapoda of Suriname (Dutch Guiana). — Zool. Verhandel. 44: 1—296. pl. 1—16. Leiden.
- HOLTHUIS L. B.—GOTTLIEB E. 1958: An annotated list of the Decapod Crustacea of the Mediterranean Coast of Israel. — Bull. Research Council Israel. 7B (1—2): 1—126.
- HORN H. S. 1966: Measurement of "overlap" in comparative ecological studies. — American Naturalist. 100: 419—424.
- HÖPFNER G. 1974: Krabben Mursia Desmarest und Panopeus H. Milne-Edwards. — Der Geschiebesammler. 9 (1): 31—32. text-figs. 1, 2.
- HUMMON W. D. 1974: A similarity index based on shared species diversity, used to assess temporal and spatial relationships among intertidal marine Gastrichia. — Oecologia. 17: 203—220.
- IMAIZUMI R. 1958: Super-Specialization of Crab. — Congr. Geol. Int. XX. Mexico. Sobretiro Secc. VII: 275—286. pl. 1. text-figs. 1—18.
- INGLE R. W. 1980: British Crabs. — 1—222, pl. 1—34. Oxford.
- JANSSEN A. W.—MÜLLER P. (in press): Miocene Decapoda and Mollusca from Ramsel, municipality of Herselt (Belgium, province of Antwerp), with descriptions of a new crab genus and a new coleoid cephalopod species. — Scripta Geol. Leiden.
- JANSSEN R. 1972: Beiträge zur Kenntnis der Bryozoa, Vermes, Crustacea und Echinodermata aus dem nordwestdeutschem Mittel- und Obermiozän. — Veröff. Überseemus. Bremen. A. 4 (11): 71—108.
- KELLY S. R. A.—MC LACHLAN A. 1980: The use of silicone rubbers in the preparation of casts from natural fossil moulds. — Geol. Mag. 117 (5): 447—454. pl. 1.

- KOJUMDGIEVA E. 1975: Paléoécologie des communautés des Mollusques du Miocène en Bulgarie du Nord-Ouest. II. Communautés des Mollusques du Badénien (Miocène moyen) en Bulgarie du Nord-Ouest. — Geol. Balc. 6 (2): 63—94.
- KÓKAY J.—MÍHÁLY S.—MÜLLER P. (in press.) Bádeni korú rétegek a budapesti Örs vezér tere könyékén (Badenian exposures near Örs vezér tere, Budapest). — Földt. Közl.
- KRACH W. 1981: Badeńskie utwory rafowe na Roztoczu Lubelskim Oraz Ślimaki. — Geol. Prace, 121.
- LELKE Gy.—MÜLLER P. (in press.) Foraminifera-alga onkoidok a budapesti miocénből (Foraminiferal-algal oncoids from the Miocene of Budapest). — Földt. Közl.
- LÖRENTHEY I. 1897: Adatok Magyarország harmadkorú rák faunájához. — Math. és Term. Tud. Ért. 15: 149—169.
- LÖRENTHEY I. (E.)\* 1898a: Beiträge zur Decapodenfauna des ungarischen Tertiärs. — Math. Naturwiss. Berichte aus Ungarn. 14: 92—115.
- LÖRENTHEY I. 1898b: Adatok Magyarország harmadkorú rák-faunájához. — Math. és Term. Tud. Közl. 27 (2): 103—271. pl. 1—9.
- LÖRENTHEY I. (E.) 1898c: Beiträge zur Decapodenfauna des Ungarischen Tertiärs. — Természettrajzi Füz. 21: 1—133. pl. 1—9.
- LÖRENTHEY I. 1901a: „Andorina” és „Darányia” két új ráknem Magyarországból. — Math. és Term. Tud. Közl. 27 (5): 831—838. pl. 1.
- LÖRENTHEY I. (E.) 1901b: Andorina und Darányia, zwei neue Brachyurengattungen aus Ungarn. — Math. Wiss. Berichte aus Ungarn. 17: 328—336. pl. 1.
- LÖRENTHEY I. 1904a: Palaeontológiai tanulmányok a harmadkorú rákok köréből. — Math. és Term. Tud. Ért. 22: 160—167.
- LÖRENTHEY I. (E.) 1904b: Paläontologische Studien über tertiäre Decapoden. — Math. Naturwiss. Berichte aus Ungarn. 22: 29—36.
- LÖRENTHEY I. 1907: Adatok Sardínia harmadidőszaki rák-faunájához. — Math. és Term. Tud. Közl. 19 (2): 243—296. pl. 1, 2.
- LÖRENTHEY I. (E.) 1909: Beiträge zur tertiärer Decapodenfauna Sardiniens. — Math. Naturwiss. Berichte aus Ungarn. 24: 202—259. pl. 1, 2.
- LÖRENTHEY I. 1911: Újabb adatok Budapest környéke harmadidőszaki üledékeinek geológiájához. — Math. és Term. Tud. Ért. 29: 515—535.
- LÖRENTHEY I. (E.) 1913: Neuere Beiträge zur Stratigraphie der Tertiärbildungen in der Umgebung von Budapest. — Math. Naturwiss. Berichte aus Ungarn. 27: 282—394.
- LÖRENTHEY I. (E.)—BEURLEN K. 1929: Die fossilen Dekapoden der Länder der Ungarischen Krone. — Geol. Hung. ser. Pal. 3: 1—420. pl. 1—12, text-figs. 1—49.
- MARTIN K. 1880: Die Tertiärschichten auf Java. — Leiden.
- MILLET DE LA TURTAUDIÈRE P. 1854: Paléontologie de Maine-et-Loire. — Angers.
- MILLET DE LA TURTAUDIÈRE P. 1866: Paléontographie de Maine-et-Loire ou description des fossiles nouveaux du terrain tertiaire marin ou terrain miocène supérieur du département de Maine-et-Loire. — Angers.
- MÜLLER P. 1974a: Decapoda (Crustacea) fauna a budapesti miocénből 1. [Les faunes de Crustacés Décapodes des calcaires miocènes de Budapest.] — Földt. Közl. 102 (2): 119—132. pl. 1—5, text-fig. 1.
- MÜLLER P. 1974b: Decapoda (Crustacea) fauna a budapesti miocénből 2. [Faune de Décapodes (Crustacés) du Miocène de Budapest.] — Földt. Közl. 102 (3): 275—287. pl. 1—3.
- MÜLLER P. 1976a: Decapoda (Crustacea) fauna a budapesti miocénből 3. [Faune de Décapodes (Crustacés) du Miocène de Budapest.] — Földt. Közl. 105 (4): 506—512. pl. 1—3.
- MÜLLER P. 1976b: Trapezia (Crustacea Decapoda) a magyar eocénból és miocénből [Trapezia (Crustacea Decapoda) dans l'Éocène et le Miocène de Hongrie]. — Földt. Közl. 105 (4): 516—523. pl. 1, 2.
- MÜLLER P. 1976c: Decapoda (Crustacea) fauna a budapesti miocénből 4. [Faune de Décapodes (Crustacés) dans le Miocène de Budapest.] — Földt. Közl. 106 (2): 149—160. pl. 1—5.
- MÜLLER P. 1979a: Decapoda (Crustacea) fauna a budapesti miocénből 5. [Faune de Décapodes (Crustacés) dans le Miocène de Budapest]. — Földt. Közl. 108 (3): 272—312. pl. 1—23.
- MÜLLER P. 1979b: Décapodes du Badénien et Sarmatiens de Bulgarie. — Palaeontology. Strat. and Lith. 10: 3—7. pl. 1—3. Sofia.
- MÜLLER P. 1979c: The Indo-West-Pacific character of the Badenian Decapod Crustaceans of the Paratethys. — Ann. Géol. Pays Hellen., Tome hors série, 2: 865—869.
- NAGYMAROSI A. 1980: A magyarországi bádenien korrelációja nannoplankton alapján (Correlation of the Badenian in Hungary on the basis of the Nannoplankton). — Földt. Közl. 110 (2): 206—245. pl. 1—7.
- NOSZKY J. sen. 1940: A Cserhát-hegység földtani viszonyai (Die Geologie des Cserhát-Berges). — Magy. Tájak Földt. Leír. 3. Budapest.
- ODHNER T. 1925: Monographierte Gattungen der Krabbenfamilie Xanthidae I. — Göteborgs Kungl. Vetens. och Vitterhets Samhälles Handlingar, Fjärde Följden. 29 (1): 1—92. pl. 1—5, text-figs. 1—7.
- ORTMANN A. E. 1897: Carcinologische Studien. — Zool. Jahrb. Abth. Syst. Geogr. Biol. der Thiere. 10 (3): 258—272. pl. 1—17.
- PAPP A.—CICHA I.—SENEŠ J.—STEININGER F. 1978: M<sub>4</sub> Badenien. — Chronostratigraphie und Neostratotypen, Miozän der Zentralen Paratethys. 6. Bratislava.
- PAPP A.—ZAPFE H.—BACHMAYER F.—TAUBER A. F. 1947: Lebensspuren mariner Krebse. — Sitzungsber. Akad. Wiss. Wien. I. 155 (8—10): 281—317. text-figs. 1—11.

\* see page 32.

- PERÈS J. M.—PICARD J. 1964: Nouveau Manuel de Bionomie Benthique de la Mer Méditerranée. — Recueil des Travaux de la Station Marine d'Endoume. 31: 1—47.
- PESTA O. 1918: Die Decapodenfauna der Adria. — 500 p. text-figs. 1—152. Leipzig—Wien.
- PHILIPPE M.—SECRETAN S. 1971: Crustacés Décapodes du Burdigalien des Courances (Vaucluse). — Ann. de Paléont. Invert. 57 (2): 3—20. pl. A—C, text-figs. 1, 2.
- PINKHAM C. F. A.—PEARSON J. O. 1976: Application of a new coefficient of similarity to pollution surveys. — J. Water Poll. Contr. Fed. 48: 717—723.
- POR F. D. 1978: Lessepsian Migration. The influx of Red Sea Biota into the Mediterranean by way of the Suez Canal. — 1—228. pl. 1—10. Springer Verlag. Berlin—Heidelberg—New York.
- RÖGL F.—STEININGER F.—MÜLLER C. 1978: Middle Miocene Salinity Crisis and the Paleogeography of the Paratethys (Middle and Eastern Europe). — Initial Rep. D. S. D. P. 42 (1): 985—990. Washington.
- RATHBUN M. J. 1911: Marine Brachyura. In: The Percy Sladen Trust Expedition to the Indian Ocean in 1905. — Trans. Linn. Soc. 2nd ser. Zool. 14 (2): 191—261. pl. 15—20. London.
- RATHBUN M. J. 1925: The Spider Crabs of America. — Smithson. Inst. Bull. 129: 1—613. pl. 1—283.
- RATHBUN M. J. 1930: The Cancroid Crabs of America. — Smithson. Inst. Bull. 152: 1—609, pl. 1—230.
- RATHBUN M. J. 1937: The Oxystomatous and allied Crabs of America. — Smithson. Inst. Bull. 166: 1—278. pl. 1—86.
- RATHBUN M. J. 1945: Decapod Crustacea. In: Ladd—Hoffmeister: Geology of Lau. — Bernice P. Bishop Museum Bull. 181: 373—391. pl. 54—62.
- RÉMY J. M. 1952: *Macrophthalmus aquensis* A. Milne-Edwards et Brocchi 1897, son stade Mégalope. Le genre *Macrophthalmus* Latreille 1829. — C. R. Somm. et Bull. Soc. Geol. France, Notes et Mémoires, 6-eme ser. 2: 27—39. text-fig. 1.
- REUSS A. 1859: Zur Kenntniss fossiler Krabben. — 1—90. pl. 1—24. Wien.
- REUSS A. E. 1867: Die fossile Fauna der Steinzalzablagerung von Wieliczka in Galizien. — Sitzungsber. Math. Naturwiss. Cl. Akad. Wiss. Wien. Abt. I. 40: 173—176. pl. 8.
- REUSS A. E. 1872: Die fossilen Korallen des österreichisch-ungarischen Miozäns. — Denkschr. Akad. Wiss. Wien. Math. Naturwiss. Cl. 31: 197—270, pl. 1—21.
- REUSS R. 1871: *Phymatocarcinus speciosus*, eine neue fossile Krabbe aus dem Leithakalk des Wiener Beckens. — Sitzungsber. Akad. Wiss. Wien. Abt. 2. 63: 325—330. pl. 1.
- SAINT LAURENT DE M. 1980a: Sur la classification et la phylogénie des Crustacés Décapodes Brachyoures, I Podotremata Guinot 1977 et Eubrachyura sect. nov. — C. R. Acad. Sci. 290 (D/19): 1265—1268. Paris.
- SAINT LAURENT DE M. 1980b: Sur la classification et la phylogénie des Crustacés Décapodes Brachyoures, Hetrotremita et Thoracotremata Guinot, 1977. — C. R. Acad. Sci. 290 (D/20): 1317—1320. Paris.
- SAINT LAURENT DE M.—LE LOEFF P. 1979: Crustacés Décapodes Thalassinidea. I Upogebiidae et Callianassidae. In: Campagnes de la Calypso au large des côtes Atlantiques Africaines (1956 et 1959), 22. — Ann. Inst. Océanogr. Nouv. Sér. 55: 29—101. text-figs. 1—28.
- SAKAI T. 1976: Crabs of Japan and the adjacent Seas. Kodansha.: 1—773. pl. 1—251, text-figs. 1—379. Tokio.
- SCHAFAZIK F. 1903: Budapest harmadik főgyűjtőcsatornájának földtani szelvénye (The geologic section of the trench of the third main drainpipe of Budapest). — Földt. Közl. 33 (1—4): 45—53. pl. 1.
- SCHAFAZIK F.—VENDL A. 1929: Geológiai kirándulások Budapest környékén. — 1—343. Budapest.
- SCHÄFER W. 1951: Fossilizations-Bedingungen Brachyuren Krebse. — Abh. Senck. Nat. Forsch. Ges. 485: 225—230.
- SCHÄFER W. 1954: Form und Function der Brachyuren-Schere. — Abh. Senck. Nat. Forsch. Ges. 489: 1—65.
- SCHAFFER F. X. 1927: Der Begriff der „miozänen Mediterraenstufen“ ist zu streichen. — Verh. Geol. Bundesanst.: 86—88. Wien.
- SCHOLZ G. 1970: A visegrádi Fekete-hegy tortonai korall-faunája (The “tortonian” coralfuna of Fekete-hill at Visegrád). — Földt. Közl. 100: 192—206. pl. 1—5.
- SCHOUPPÉ A. 1949: Zwei Decapoden aus dem Torton von Retznei. — Mitt. Naturwiss. Vereines Steiermark. 77—78: 139—141. text-figs. 1—2. Graz.
- SERÈNE R. 1965: A review of Philippine Albuneidae, with description of two new species. — The Philippine Journ. Sci. 94 (1): 87—116. pl. 1—6.
- SERÈNE R. 1969: Observations on species of the group *Trapezia rufopunctata*, with a provisional key for all species of *Trapezia*. — Journ. Mar. Biol. Ass. India. 11 (1—2): 126—148. text-figs. 1—33. Madras.
- SERÈNE R.—NGUYEN VAN LUOM 1958: Chlorodopsis (Brachyura) du Viêt-Nam. — pp.: 1—147. pl. 1—4. Saigon.
- SERÈNE R.—NGUYEN VAN LUOM 1959: Note additionnelle sur les espèces de Chlorodopsis (Brachyours). — Ann. Fac. Sci.: 301—340. pl. 1—3.
- SHOUP J. B. 1968: Shell opening by crabs of the genus Calappa. — Science. 160 (3830): 887—888.
- SØRENSEN T. 1948: A method for establishing groups of equal amplitude in plant sociology based on similarity of species content and its application to analyses of the vegetation on Danish commons. — Biol. Skr. 5 (4): 1—34.
- SOKAL R. R.—MICHENER C. D. 1958: A statistical method for evaluating systematic relationships. — Univ. Kansas Sci. Bull. 38: 1409—1438.
- STANCIU I.—ANDREESCU E. 1968: Fauna tortoniana din regiunea Rugi-Delinesti (Bazinul Caransebesului). — Studii și Cercetări de Geol. Geofiz. Geogr. ser. Geol. 2: 455—471. pl. 1—7.
- STEPHENSON W. 1972: An annotated check-list and key to the Indo-west Pacific Swimming Crabs (Crustacea: Decapoda: Portunidae). — Bull. Roy. Soc. New Zealand. 10: 1—64.
- STEPHENSON W.—CAMPBELL B. 1960: The Australian Portunids (Crustacea: Portunidae), IV Remaining Genera. — Austr. Journ. of Marine and Freshwater Research. 11 (1): 73—122. pl. 1—6.

- STEPHENSON W. — HUDSON J. J. 1957: The Australian Portunids (Crustacea: Portunidae), I The genus Thalamita. — Austr. Journ. of Marine and Freshwater Research. 8 (3): 312—368.
- STOJASPAL FR. 1975: (Anhang) Katalog der Typen und Abbildungsoriginale der Geologischen Bundesanstalt. 1. Teil. Wirbellose des Kainozoikums (exclusive Microfossilien). — Verh. Geol. Bundesanst. (1): A159—A192. Wien.
- STOLL N. R. et al. (red.) 1961: International Code of Zoological Nomenclature Adopted by the XV. International Congress of Zoology. — London.
- STRAUSZ L. 1924: A biai miocén (The Miocene of Bia). — Földt. Közl. 53: 53—58.
- STRAUSZ L. 1927: Eine neue Fundstätte der Obermediterrans in Budapest und eine neue Pecten-Form. — Centralbl. Min. Geol. Pal. B.: 340—343, text-figs. 1—7.
- STRAUSZ L. 1966: Die Miozän-Mediterranen Gastropoden Ungarns. — 534 p., pl. 1—79. Akad. Kiadó, Budapest.
- SZABÓ J. 1879: Budapest geológiai tekintetben. — : 1—116. pl. 1—3, 1 map. Budapest.
- TAYLOR J. D. 1968: Coral reef and associated invertebrate communities (mainly Molluscan) around Mahé, Seychelles. — Phil. Trans. Roy. Soc. ser. B (793) 254: 129—206. London.
- TOULA F. 1904: Über eine neue Krabbe (Cancer Bittneri n. sp.) aus dem miozänen Sandsteine von Kalksburg bei Wien. — Jahrb. Geol. Reichsanst. 54: 161—168. text-figs. 1—5. Wien.
- VADÁSZ M. E. 1906: Budapest — Rákos felsőmediterrán korú faunája (Über die obermediterranen Fauna von Budapest — Rákos). — Földt. Közl. 36: 256—351.
- VAN STRAELEN V. 1937: Sur quelques Crables du Sahélien moyen des environs d'Oran (Algérie). — C. R. Somm. des Séances. Soc. Géol. France, 1936: 473—480. pl. 33.
- VASS D. et al. 1978: Chronology of volcanic events in Eastern Slovakia on the ground of isotopical-paleomagnetical researches. — Geol. Prace 71: 77—89.
- VEIGA FERREIRA O. DA 1954: Malacostráceos do Miocenico marinho de Portugal. — Comm. Serv. Geol. Port. 35: 5—23. pl. 1—6.
- VEIGA FERREIRA O. DA 1958: Descoberto de „Calappa heberti” no Tortoniano do Penedo. — Comin. Serv. Geol. Port. 42: 203—207. pl. 1, text-figs. 1—3.
- VEIGA FERREIRA O. DA 1961: Nova especie de Callianassa no Miocenico do Bacia do Tejo. — Comm. Serv. Geol. Port. 45: 479—482. pl. 1.
- VEIGA FERREIRA O. DA 1965: Nova contribuição para o conhecimento dos Malacostráceos do Miocénico Marinho de Portugal. — Comm. Serv. Geol. Port. 48: 5—17. pl. 1, 2.
- VIA L. 1948: Braquiuros fósiles de Barcelona y sus alrededores. — An. Serv. Técn. Agricult. 7: 143—152. pl. 1. Barcelona.
- VIA BOADA L. 1955: Un Braquiuro fósil (*Geryon latifrons*) en el Plaisencience. — Estud. Geol. España, 26: 219—222. pl. 34.
- VIA BOADA L. 1980: Ocypodoidea (Crustacés Décapodes) du Cénozoïque Méditerranéen, origine et évolution de cette super-famille. — Ann. Pal. Invert. 66: 51—66. pl. 1.
- VIA BOADA L. 1981: Les Crustacés Décapodes du Cénomanien de Navarra (Espagne): premiers résultats de l'étude des Galatheidae. — Geobios. 14 (2): 247—251. text-figs. 1—5.
- WEIMER R. J.—HOYT J. H. 1964: Burrows of *Callianassa major* Say, geologic indicators of littoral and shallow neritic environment. — Journ. Pal. 38 (4): 761—767.
- WOODWARD H. 1873: 6th report of the committee app. for the purpose of cont. researches in foss. Crust. — Rep. 42nd Meeting Brit. Assoc. Adv. Sci. Brighton.
- (YANAKEVICH A. N.) Янакевич А. Н. 1969: К характеристике крабов из биогермных известняков миоцена северо-западной Молдавии. — Палеонт. сб. 6 (I) Львов.
- (YANAKEVICH A. N.) Янакевич А. Н. 1977: Среднемиоценовые рифы Молдавии. — Кишинев.
- ZARIQUIEY ALVAREZ R. 1968: Crustaceos Decápodos Ibericos. — Investig. Peñq. 32 (I—XV): 1—510. figs. 1—164.

LIST OF SPECIES

	Page	Plate
<b>Ordo DECAPODA</b> .....	49	
<b>Infraordo ANOMURA</b> .....	49	
<b>Familia Laomediidae</b> .....	49	
<i>Jaxea kuemeli</i> BACHMAYER .....	49	—
<b>Familia Callianassidae</b> .....	49	
‘ <i>Callianassa</i> ’ <i>munieri</i> BROCCHI .....	50	I, II
‘ <i>Callianassa</i> ’ <i>chalmasi</i> BROCCHI .....	50	II
‘ <i>Callianassa</i> ’ <i>rakosiensis</i> LÖRENTHEY .....	51	III
<i>Callianassa brocchii</i> LÖRENTHEY .....	51	III
‘ <i>Callianassa</i> ’ <i>pseudorakosiensis</i> LÖRENTHEY .....	52	III, IV, V
‘ <i>Callianassa</i> ’ <i>kerepesiensis</i> MÜLLER .....	52	VI
‘ <i>Callianassa</i> ’ cf. <i>kerepesiensis</i> MÜLLER .....	52	VII
‘ <i>Callianassa</i> ’ cf. <i>jahringensis</i> GLAESSNER .....	53	—
‘ <i>Callianassa</i> ’ <i>sismondai</i> A. MILNE-EDWARDS .....	53	—
‘ <i>Callianassa</i> ’ <i>norica</i> GLAESSNER .....	53	—
‘ <i>Callianassa</i> ’ sp. ind. .....	53	V
? <i>Callianassa szobensis</i> n. sp. .....	53	VII
<b>Familia Upogebiidae</b> .....	54	
<i>Upogebia</i> sp. (?div.) .....	54	VIII
<b>Familia Paguridae</b> .....	54	
<i>Pagurus rakosensis</i> MÜLLER .....	54	X
<i>Pagurus concavus</i> MÜLLER .....	55	IX
‘ <i>Pagurus</i> ’ <i>albus</i> MÜLLER .....	55	XI
‘ <i>Anapagurus</i> ’ <i>miocenicus</i> MÜLLER .....	55	XI
‘ <i>Anapagurus</i> ’ <i>marginatus</i> MÜLLER .....	56	XII
? <i>Pagurus turcus</i> n. sp. .....	56	XII, XIII
‘ <i>Pylopagurus</i> ’ <i>leganyii</i> n. sp. .....	56	XIV
<i>Pagurinae</i> sp. 1. .....	57	XV
<i>Pagurinae</i> sp. 2. .....	57	XV
<i>Diogenes matrensis</i> n. sp. .....	57	XVI
<i>Diogenes</i> cf. <i>pugilator</i> (ROUX) .....	57	XVI
<i>Dardanus arrosor</i> (HERBST) .....	58	XVII
<i>Dardanus substriatiformis</i> (LÖRENTHEY) .....	58	XVII
<i>Dardanus hungaricus</i> (LÖRENTHEY) .....	58	XVIII
<i>Paguristes cserhatensis</i> n. sp. .....	59	XIX
<i>Petrochirus priscus</i> (BROCCHI) .....	59	XIX, XX, XXI
<b>Familia Galatheidae</b> .....	60	
<i>Galathea weinfurteri</i> BACHMAYER .....	60	XXI, XXII
<b>Familia Porcellanidae</b> .....	60	
<i>Petrolisthes magnus</i> n. sp. .....	60	XXIII, XXIV, XXV
<i>Petrolisthes haydni</i> n. sp. .....	61	XXVI
<i>Pisidia kokayi</i> (MÜLLER) .....	61	XXVII
<i>Pisidia</i> cf. <i>kokayi</i> (MÜLLER) .....	62	XXVII, XXVIII

	Page	Plate
Familia Albuneidae .....	62	
<i>Albunea asymmetrica</i> (MÜLLER) .....	62	XXVIII
Infraordo BRACHYURA .....	62	
Familia Dromiidae .....	62	
<i>Dromia neogenica</i> MÜLLER .....	63	XXIX
<i>Dromia eotvoesi</i> (MÜLLER) .....	63	XXIX, XXX
<i>Kromitis koberi</i> (BACHMAYER) .....	64	XXXI
<i>Kerepesia viari</i> MÜLLER .....	64	XXXI, XXXII
<i>Dromiidarum</i> n? gen. n. sp. ....	65	XXXII
Familia Dynomenidae .....	65	
<i>Dynomene emiliae</i> MÜLLER .....	65	XXXIII
Familia Dorippidae .....	65	
<i>Dorippe margaretha</i> LÖRENTHEY .....	66	XXXIV
<i>?Dorippe carpathica</i> (FÖRSTER) .....	66	—
Familia Calappidae .....	66	
<i>Calappa praelata</i> LÖRENTHEY .....	66	XXXV, XXXVI
<i>Calappa heberti</i> BROCHI .....	67	XXXVII, XXXVIII
<i>Calappilia matzkei</i> (BACHMAYER) .....	68	—
<i>Mursia lienharti</i> (BACHMAYER) .....	68	XXXIX, XL
<i>Matuta brocchii</i> GLAESSNER .....	69	XLI, XLII
Familia Leucosiidae .....	69	
<i>Ebalia vanstraeleni</i> BACHMAYER .....	69	XLIII
<i>Ebalia oersi</i> MÜLLER .....	69	XLIII
<i>Ebalia</i> sp. 1. ....	70	XLIV
<i>Ebalia</i> sp. 2. ....	70	XLIV
<i>Ebalia</i> sp. 3. ....	70	XLIV
<i>'Ebalia'</i> hungarica MÜLLER .....	70	XLIV, XLV
<i>Andorina elegans</i> LÖRENTHEY .....	70	XLVI
<i>'Palaeomyra'</i> globulosa (MÜLLER) .....	71	XLVII
Familia Raninidae .....	71	
<i>Ranidina rosaliae</i> BITTNER .....	71	XLVI
Familia Majidae .....	71	
<i>Maja biaensis</i> LÖRENTHEY .....	71	XLVIII, XLIX
<i>Schizophrys visegradensis</i> n. sp. ....	72	XLIX, L
<i>Achaeus magnus</i> MÜLLER .....	72	L, LI
<i>'Pisa'</i> oroszyi (BACHMAYER) .....	73	LI, LII
<i>Micippa hungarica</i> (LÖRENTHEY) .....	73	LII, LIII
Familia Parthenopidae .....	74	
<i>Parthenope szaboi</i> MÜLLER .....	74	LIII, LIV
<i>Parthenope tetenyensis</i> n. sp. ....	74	LV, LVI
Familia Corystidae .....	75	
<i>Corystites latifrons</i> (LÖRENTHEY) .....	75	—
Familia Canceridae .....	75	
<i>Cancer sismondai</i> (MEYER) .....	75	—
<i>Cancer styriacus</i> BITTNER .....	76	LVII, LVIII
<i>Cancer illyricus</i> BITTNER .....	76	LVIII
<i>Microdium nodulosum</i> REUSS .....	76	—
<i>Miocyclus bulgaricus</i> MÜLLER .....	77	LVIII
<i>"Cancer"</i> szontaghii LÖRENTHEY .....	77	LIX, LX
<i>"Cancer"</i> szontaghii vindobonensis (BACHMAYER — KÜPPER) .....	78	—
<i>"Cancer"</i> carniolicus BITTNER .....	78	—
<i>"Cancer"</i> bittneri TOULA .....	78	—

	Page	Plate
<b>Familia Pirimelidae</b> . . . . .	78	
<i>Pirimela lorentheysi</i> (MÜLLER) . . . . .	78	LX
<i>Trachypirimela grippi</i> (MÜLLER) . . . . .	79	LXI
<b>Familia Portunidae</b> . . . . .	79	
<i>Portunus monspeliensis</i> (A. MILNE-EDWARDS) . . . . .	79	LXII
<i>Portunus neogenicus</i> MÜLLER . . . . .	80	LXII
<i>Portunus (Monomia) miocaenicus</i> n. sp. . . . .	80	LXII
<i>Charybdis (Goniosupradens) mathiasi</i> n. sp. . . . .	81	LXIII, LXIV, LXVII
<i>Thalamita fragilis</i> MÜLLER . . . . .	81	LXV, LXVI
<i>Thalamita</i> n? sp. . . . .	82	LXVII
<i>Rakosia carupoides</i> n. sp. . . . .	82	LXVIII
<i>Necronectes schafferi</i> GLAESSNER . . . . .	82	LXVI, LXIX
<i>Liocarcinus rakosensis</i> (LÖRENTHEY) . . . . .	83	LXIX, LXX
<i>Liocarcinus oroszyi</i> (BACHMAYER) . . . . .	83	LXXI
<i>Liocarcinus kuehni</i> (BACHMAYER) . . . . .	84	LXXI
<i>Miopipus pygmeus</i> (BROCCHI) . . . . .	84	LXXII
<i>Portumnus tricarinatus</i> LÖRENTHEY . . . . .	85	—
‘ <i>Xaiva</i> ’ <i>bachmayeri</i> n. sp. . . . .	85	LXXII
<i>Mioxoiva psammophila</i> MÜLLER . . . . .	86	LXXIII
? <i>Carupa</i> cf. <i>tenuipes</i> DANA . . . . .	86	LXXIII
‘ <i>Lissocarcinus</i> ’ <i>szoeraenyiae</i> (MÜLLER) . . . . .	86	LXXIII
<b>Familia Xanthidae</b> . . . . .	87	
<i>Actaea turcocampestris</i> n. sp. . . . .	87	LXXIV
<i>Carpilius antiquus</i> GLAESSNER . . . . .	87	LXXV
<i>Chlorodiella mediterranea</i> (LÖRENTHEY) . . . . .	88	LXXVI, LXXVII
<i>Chlorodiella mediterranea tetenyensis</i> n. ssp. . . . .	88	LXXVII
<i>Chlorodiella loczyi</i> n. sp. . . . .	89	LXXVIII
<i>Chlorodiella juglans</i> n. sp. . . . .	89	LXXVIII
<i>Daira speciosa</i> (REUSS) . . . . .	90	LXXIX, LXXX
<i>Haydnella steiningeri</i> n. sp. . . . .	90	LXXX, LXXXI
<i>Panopeus wronai</i> n. sp. . . . .	91	LXXXI, LXXXII, LXXXIII
‘ <i>Pilodius</i> ’ <i>vulgaris</i> (GLAESSNER) . . . . .	91	LXXXIII, LXXXIV
<i>Trapezia glaessneri</i> MÜLLER . . . . .	92	LXXXIV, LXXXV
<i>Xantho moldavicus</i> (YANAKEVICH) . . . . .	92	LXXXV, LXXXVI, LXXXVII
<i>Pilumnus mediterraneus</i> (LÖRENTHEY) . . . . .	93	LXXXVII, LXXXVIII
<i>Actumnus telegdii</i> (MÜLLER) . . . . .	94	LXXXIX, XC
<i>Actumnus</i> ?n. sp. . . . .	94	XC
‘ <i>Glabropilumnus</i> ’ n. sp. . . . .	95	XCI
<i>Pilumnopeus paratethyensis</i> n. sp. . . . .	95	XCII
‘ <i>Pilumnopeus</i> ’ <i>tetenensis</i> n. sp. . . . .	95	XCI
<b>Familia Geryonidae</b> . . . . .	95	
<i>Geryon</i> cf. <i>latifrons</i> VAN STRAELEN . . . . .	96	XCIII
<b>Familia Gonoplacidae</b> . . . . .	96	
<i>Goneplax gulderi</i> BACHMAYER . . . . .	96	XCIII
“ <i>Pilumnoplax</i> ” <i>carnuntinus</i> BACHMAYER . . . . .	96	—

	Page	Plate
<b>Familia Grapsidae</b> . . . . .	96	
<i>Pachygrapsus hungaricus</i> MÜLLER . . . . .	97	XCIII, XCIV
<i>Brachynotus februarius</i> MÜLLER . . . . .	97	XCV
<b>Familia Ocypodidae</b> . . . . .	97	
? <i>Ocypode</i> sp. . . . .	98	—
<i>Macrophthalmus vindobonensis</i> GLAESSNER . . . . .	98	XCV, XCVI
<b>Familia Palicidae</b> . . . . .	98	
<i>Palicus</i> sp. . . . .	98	XCVII
<i>Crossotonotus diosdensis</i> n. sp. . . . .	99	XCVII




## **PLATES**

Scanning electron micrographs were taken by Mrs. B. TAKÁCS. All other photographs were made by the author.

Abbreviations of collections: FI: Geological Institute, Budapest; NMÖ: Palaeontological Department of Natural History Museum Budapest; CA: author's collection; NHMW: Naturhistorisches Museum Wien. Codes of layers are explained in chapter "Description of localities" (p. 36).

**PLATE I**

- 1, 4. '*Callianassa*' *munieri* BROCCHI  
Rákos, MR-5-1, right propodus 2.5×
- 2—3. '*Callianassa*' *munieri* BROCCHI  
Rákos, MR-5-4, right dactylus 4.7×
5. '*Callianassa*' *munieri* BROCCHI  
Rákos, FI, leg. SCHAFARZIK, unidentified layer, three articles of left cheliped 2.5×
6. '*Callianassa*' *munieri* BROCCHI  
Rákos, MR-5-2, left carpus 4×
7. '*Callianassa*' *munieri* BROCCHI  
Rákos, unidentified layer, FI, leg. STOCZEK, merus of left cheliped 4.9×



**PLATE II**


1. '*Callianassa*' *munieri* BROCCHI  
Rákos, MR-5-5, right dactylus 4.6×
2. '*Callianassa*' *munieri* BROCCHI  
Rákos, MR-5-6, left dactylus 3.2×
3. '*Callianassa*' *chalmasii* BROCCHI  
Gyakorló út, MGY-9-3, left propodus 5.6×
4. '*Callianassa*' *chalmasii* BROCCHI  
Gyakorló út, MGY-9-2, left propodus 5.5×
- 5—7. '*Callianassa*' *chalmasii* BROCCHI  
Gyakorló út, MGY-9-1, right chela 4.6×




1




2




3



4



5



6



7

**PLATE III**

- 1—2. '*Callianassa*' *rakosiensis* LÖRENTHEY  
Gyakorló út, MGY-6-1, right propodus 6.2×
3. *Callianassa brocchii* LÖRENTHEY  
Gyakorló út, MGY-24-1, right propodus 7.3×
- 4—5. '*Callianassa*' *pseudorakosensis* LÖRENTHEY Lectotype  
Rákos, unidentified layer, leg. LÖRENTHEY, M 139, FI, right chela 8.5×






PLATE IV

- 1—2. '*Callianassa*' *pseudorakosensis* LŐRENTHEY Lectotype  
Rákos, unidentified layer, leg. LŐRENTHEY, M 139. FI 8.5×
3. '*Callianassa*' *pseudorakosensis* LŐRENTHEY  
Gyakorló út, MGY-7-2, left propodus 6.8×
- 4—6. '*Callianassa*' *pseudorakosensis* LŐRENTHEY  
Gyakorló út, MGY-7-3, right propodus 7.5×



1




2



4



3



6



5

**PLATE V**

- 1—3. '*Callianassa*' *pseudorakosensis* LÖRENTHEY  
Szentkút, leg. STREDA?, M 61/2633 NMÖ, left dactylus 9×
4. '*Callianassa*' *pseudorakosensis* LÖRENTHEY  
Szentkút, MSZ-1-1, right merus of cheliped 11.5×
- 5—6. *Callianassa* sp. (*Semiranina oroszyi* BACHMAYER)  
Tétény-plateau, MDG-1-1, anterior part of carapace 10×
- 7—8. '*Callianassa*' sp.  
Órs vezér tere, MOE-34-1 and 34-2 9×




**PLATE VI**


- 1—2. '*Callianassa*' *kerepesiensis* MÜLLER Holotype  
Kerepesi út, MKC-2-1, right propodus 13×
3. '*Callianassa*' *kerepesiensis* MÜLLER  
Szentkút, leg. STREDA?, M 62/2640 NMÖ, right propodus, cutting edge 9×
4. '*Callianassa*' *kerepesiensis* MÜLLER  
Örs vezér tere, MOE-11-1, left propodus inner side 10×
5. '*Callianassa*' *kerepesiensis* MÜLLER  
Örs vezér tere, MOE-11-2, left propodus outer side 10×
6. '*Callianassa*' *kerepesiensis* MÜLLER  
Szentkút, leg STREDA?, M 83/153 NMÖ, right dactylus 17×




1




2




3



4




5



6

**PLATE VII**

- 1—2. '*Callianassa*' cf. *kerepesiensis* MÜLLER  
Örs vezér tere, MOE-12-3, right propodus 14.5×
- 3—4. ?*Callianassa szobensis* n. sp. Holotype  
Szob, Damásdi-patak, MSZ-1-1, left propodus 4.4×
- 5—6. ?*Callianassa* cf. *szobensis* n. sp.  
Szob, Damásdi-patak, MSZ-2-1, dactylus? 8.5×



1




2




3



5




4




6

**PLATE VIII**


1. *Upogebia* sp. Holotype of *U. scabra* MÜLLER  
Tétény-plateau, MDG-3-1 (= MG-1-1), NMÖ, anterior part of carapace 14×
- 2—3. ?*Upogebia* sp.  
Örs vezér tere, MOE-10-1, left dactylus 16.7×
- 4—6. ?*Upogebia* sp.  
Gyakorló út, MGY-10-1, right dactylus 12.7×



2




3



4




5



6


**PLATE IX**

- 1—2. *Pagurus concavus* MÜLLER  
Örs vezér tere, MOE-1-10, right propodus 7.3×
3. *Pagurus concavus* MÜLLER  
Örs vezér tere, MOE-1-10, cutting edge 21×
4. *Pagurus concavus* MÜLLER  
Örs vezér tere, MOE-1-11, left propodus 11×
5. *Pagurus concavus* MÜLLER  
Örs vezér tere, MOE-1-11, right dactylus 15×
6. *Pagurus concavus* MÜLLER  
Örs vezér tere, MOE-1-11, cutting edge 18×




**PLATE X**

- 1—2. *Pagurus rakosensis* MÜLLER Holotype  
Rákos, MR-1-1, right dactylus 7.4×
3. *Pagurus rakosensis* MÜLLER  
Örs vezér tere, MOE-6-1, right propodus 11×
4. *Pagurus rakosensis* MÜLLER  
Örs vezér tere, MOE-6-1, cutting edge 30×
5. *Pagurus rakosensis* MÜLLER  
Örs vezér tere, MOE-6-2, left propodus 16×




1




2



4




3



5


**PLATE XI**

- 1—2. '*Pagurus*' *albus* MÜLLER  
Örs vezér tere, MOE-2-10, right propodus 7.2×
3. '*Pagurus*' *albus* MÜLLER  
Örs vezér tere, MOE-2-10, cutting edge 20×
- 4—6. '*Anapagurus*' *miocenicus* MÜLLER Holotype  
MGY-1-1, right propodus 13.6×
7. '*Anapagurus*' *miocenicus* MÜLLER Paratype  
MGY-1-2, right propodus cutting edge SEM 120×
- 8—9. '*Anapagurus*' *miocenicus* MÜLLER Paratype  
MGY-1-2, right propodus SEM 25×




**PLATE XII**


- 1—3. '*Anapagurus*' *marginatus* MÜLLER  
Örs vezér tere, MOE-3-10, right propodi 12×
- 4—5. '*Anapagurus*' *marginatus* MÜLLER  
Örs vezér tere, MOE-3-11, -12, cutting edges 18×
6. ?*Pagurus turcus* n. sp. Paratype  
Törökmező, MTZ-17-2, right propodus 13.4×




1




4




2



3




5



6


**PLATE XIII**

- 1--3. ?*Pagurus turcus* n. sp. Holotype  
Törökmező, MTZ-17-1, right propodus 13×
4. ?*Pagurus turcus* n. sp. Paratype  
Törökmező, MTZ-17-3, right propodus 7.5×



**PLATE XIV**

- 1—3. '*Pylopagurus*' *leganyii* n. sp. Holotype  
Sámsonháza, Csúcs-hegy, M 62/3253 NMÖ, leg. F. LEGÁNYI, right propodus 7.6×
4. ?*Pylopagurus* sp.  
Törökmező, MTZ-20-1, right cheliped, mould 8.9×
5. ?'*Pylopagurus*' *leganyii* n. sp.  
Szentkút, M 62/3153 NMÖ, fragment of right propodus 4.5×




1




2



3



4








5

5

**PLATE XV**


1—3. *Pagurinae* n.? sp. 1.  
Rákos, MRF-12-1, right propodus 7.2×

4—5. *Pagurinae* n.? sp. 2.  
Kerepesi út, MKC-11-1, right propodus 7.5×




**PLATE XVI**


- 1—3. *Diogenes matrensis* n. sp. Holotype  
Mátraverebély-Szentkút, M 61/2633 NMÖ, left propodus 9.3×  
4—5. *Diogenes* cf. *pugilator* (ROUX)  
Budapest, drilling Váci út 120, 17 m, left propodus 33×




1



2



3




4



5

**PLATE XVII**

- 1--3. *Dardanus substriatiformis* (LÖRENTHEY)  
Kerepesi út, MKC-7-1, left propodus 6×
4. *Dardanus substriatiformis* (LÖRENTHEY)  
Kerepesi út, MKC-7-1, cutting edge 14×
5. *Dardanus arrosor* (HERBST)  
Törökmező, MTH-1-1, right propodus 5.6×




1




2



3




4



5

**PLATE XVIII**


- 1—4. *Dardanus hungaricus* (LÖRENTHEY)  
Tétény-plateau, MK-3-1, right propodus 6.5×
5. *Dardanus hungaricus* (LÖRENTHEY)  
Tétény-plateau, MK-3-1, cutting edge 13×



1



2



3




5



4


**PLATE XIX**

- 1—4. *Paguristes cserhatensis* n. sp. Holotype  
Sámonháza, Halastópuszta “Basis sands”, leg. HARMAT?, M 61/3907 NMÖ, left propodus 29×  
5. *Petrochirus priscus* (BROCCHI)  
Rákos, unidentified layer, M 23 FI, leg. LŐRENTHEY?, right chela 1.6×




**PLATE XX**

- 1—2. *Petrochirus priscus* (BROCCHI)  
Rákos, unidentified layer, M 23 FI, leg. LŐRENTHEY ?, right chela 2.3×
3. *Petrochirus priscus* (BROCCHI)  
Rákos, MR-2-1, right dactylus 6.4×
4. *Petrochirus priscus* (BROCCHI)  
Rákos, unidentified layer M 2364, FI, right chela 7.2×
5. *Petrochirus priscus* (BROCCHI)  
Rákos, MRF-3-1, left dactylus 5×




1




2



3



4




5

**PLATE XXI**

1. *Petrochirus priscus* (BROCCHI)  
Rákos, MRF-3-2, left fixed finger 7.6×
2. *Petrochirus priscus* (BROCCHI)  
Rákos, unidentified layer, M 2365, FI, leg. STOCZEK, right fixed finger 10.7×
3. *Petrochirus priscus* (BROCCHI)  
Rákos, MRF-3-3, right fixed finger 10.7×
4. *Galathea weinfurteri* BACHMAYER  
Rákos, MRZ-1-5, left propodus 10×
5. *Galathea weinfurteri* BACHMAYER  
Örs vezér tere, MOE-33-2, right propodus 13.5×




1




2



3




5



4

**PLATE XXII**


1. *Galathea weinfurteri* BACHMAYER  
Rákos, MRZ-1-1, carapace 6.8×
2. *Galathea weinfurteri* BACHMAYER  
Rákos, MRZ-1-2, carapace 8.6×
3. *Galathea weinfurteri* BACHMAYER  
Rákos, MRZ-1-3, rostrum 15×
4. *Galathea weinfurteri* BACHMAYER  
Örs vezér tere, MOE-33-1, carapace 13×
5. *Galathea weinfurteri* BACHMAYER  
Rákos, MRZ-1-4, carapace, *Bopyridae*-infected 7.4×




1



2



5




3



4

PLATE XXIII


- 1—2. *Petrolisthes magnus* n. sp. Holotype  
Rákos, MRZ-2-1, carapace 4.5×
3. *Petrolisthes magnus* n. sp. Paratype  
Rákos, MRZ-2-2, carapace 5×
4. *Petrolisthes magnus* n. sp. Paratype  
Rákos, MRZ-2-5, carapace, *Bopyridae*-infected 8×
5. *Petrolisthes boscii* (AUDOIN)  
Arabie, leg. BONNIE et PEREZ, det. NOBILI 1905, gift of Mus. Natl. Hist. Nat. Paris,  
carapace 7×



1



3



2





4



5

**PLATE XXIV**

- 1—2. *Petrolisthes magnus* n. sp. Paratype  
Rákos, MRZ-2-9, left chela 4.7×
- 3—4. *Petrolisthes magnus* n. sp. Paratype  
Rákos, MRZ-2-8, carpus of left cheliped 8.8×



**PLATE XXV**

1—3. *Petrolisthes boscii* (AUDOIN)


Arabie, leg. BONNIE et PEREZ, det. NOBILI, gift of Mus. Natl. Hist. Nat. Paris, left chela and carpus 7×

4. *Petrolisthes magnus* n. sp. Paratype

Rákos, MRZ-2-6, carpus of right cheliped 8.5×

5. *Petrolisthes magnus* n. sp. Paratype


Rákos, MRZ-2-10, carpus of left cheliped 8.8×




1




3



2




4



5

**PLATE XXVI**

1. *Petrolisthes haydni* n. sp. Holotype Gross-Höflein, NHMW, carapace, 22×
2. *Petrolisthes haydni* n. sp. Paratype Gross-Höflein, MNH-1-4, carapace 12×
3. *Petrolisthes haydni* n. sp. Paratype Gross-Höflein, MNH-1-3, carapace 10×
- 4—5. *Petrolisthes haydni* n. sp. Paratype Zebegény, MZZ-1-1, right chela 5.5×
6. *Petrolisthes haydni* n. sp. Paratype Gross-Höflein, MNH-1-8, carpus of right cheliped 10.5×
7. *Petrolisthes haydni* n. sp. Paratype Gross-Höflein, MHN-1-3, front 23×
8. *Petrolisthes haydni* n. sp. Paratype Gross-Höflein, KMNH-1-9, right propodus 8×



2

3

6

5

8

7

**PLATE XXVII**

1. *Pisidia kokayi* (MÜLLER) Holotype  
Tétény-plateau, MB-2-1, carapace 15.5×
2. *Pisidia kokayi* (MÜLLER)  
Rákos, MRZ-18-1, carapace 14.8×
- 3–4. *Pisidia kokayi* (MÜLLER)  
Zebegény, MZZ-2-1, carapace 21×
5. *Pisidia kokayi* (MÜLLER)  
Rákos, MRZ-18-2, carapace, *Bopyridae*-infected 15×
6. *Pisidia* cf. *kokayi* (MÜLLER)  
Törökmező, MTZ-1-1, carapace 16.5×








PLATE XXVIII


1. *Pisidia* cf. *kokayi* (MÜLLER)  
Törökmező, MTZ-1-2, carapace, rubber cast of external mould 14×
2. *Pisidia* cf. *kokayi* (MÜLLER)  
Törökmező, MTZ-1-2, carapace 14×
3. ?*Pisidia* cf. *kokayi* (MÜLLER)  
Törökmező, MTZ-1-3, internal mould of left propodus 12.2×
4. *Albunea asymmetrica* (MÜLLER) Holotype + Paratype  
MGY-2-1 and MGY-2-2, mosaic picture of carapace (holotype: left upper part 3.6×, right lower part 4×)
5. *Albunea asymmetrica* (MÜLLER) Paratype  
MGY-2-12, carpus 4.6×
6. *Albunea asymmetrica* (MÜLLER) Paratype  
MGY-2-11, chela 4.3×



1




2



5



3



6



4


**PLATE XXIX**

- 1—2. *Dromia neogenica* MÜLLER  
Diósd, MDZ-3-1, rubber-cast of internal mould, carapace 1.8×
3. *Dromia neogenica* MÜLLER Holotype  
Kerepesi út, MKC-4-1, internal mould, carapace 2.5×
- 4—5. *Dromia neogenica* MÜLLER  
Rákos, MRZ-16-2, right dactylus 5.8×
6. *Dromia neogenica* MÜLLER  
Rákos, MRZ-16-1, right propodus 2.7×
7. *Dromia eotvoesi* (MÜLLER)  
Rákos, MRO-3-1, left dactylus 5.5×
8. *Dromia eotvoesi* (MÜLLER)  
Gyakorló út, MGY-12-17 12.7×



**PLATE XXX**

- 1—2. *Dromia eotvoesi* (MÜLLER)  
Gyakorló út, MGY-12-1, carapace 4.75×
3. *Dromia eotvoesi* (MÜLLER) Holotype  
Tétény-plateau, MFA-1-1, carapace 5.7×
- 4—5. *Dromia eotvoesi* (MÜLLER)  
Gyakorló út, MGY-12-17, left propodus 12.7×
- 6—7. *Dromia eotvoesi* (MÜLLER)  
Rákos, MRO-3-1, left chela 4.8×



**PLATE XXXI**


1. *Kromtitis koberi* (BACHMAYER — TOLLMANN) Holotype  
Gross-Höflein, NHMW, carapace 3.5×
2. *Kromtitis koberi* (BACHMAYER — TOLLMANN)  
Visegrád, MV-5-2, rubber cast, carapace 3.3×
- 3—4. *Kromtitis koberi* (BACHMAYER — TOLLMANN)  
Visegrád, MV-5-3, carapace 7.7×
5. *Kerepesia viai* MÜLLER Holotype  
Kerepesi út, MKC-3-1, carapace 4.8×
- 6—7. *Kerepesia viai* MÜLLER  
Örs vezér tere, MOE-30-3, carapace, left part of the mosaic is mirror image 9.5×





1




4




3



5




6




7

**PLATE XXXII**


1. *Kerepesia viai* MÜLLER  
Örs vezér tere, MOE-30-3, carapace, right part of the mosaic is mirror image 9.5×
2. *Kerepesia viai* MÜLLER  
Kerepesi út, MKC-3-7, carapace 3.7×
- 3—4. ?*Kerepesia viai* MÜLLER  
Örs vezér tere, MOE-30-2, left chela 8.1×
- 5—6. *Dromiidarum* n? gen. n. sp.  
Törökmező, MTZ-23-1, decorticated carapace and rubber cast 5×
7. *Dromiidarum* n? gen. n. sp.  
Törökmező, MTZ-23-2, rubber cast of frontal part 9×




1




2




3



4



5




6



7

**PLATE XXXIII**

- 1—2. *Dynomene emiliae* MÜLLER Holotype  
Iasen, leg. E. KOJUMDGIEVA, coll. Univ. Sofia, carapace 3.9×
- 3—4. *Dynomene emiliae* MÜLLER  
Gross-Höflein, MNH-4-1, carapace 2.8×
5. *Dynomene emiliae* MÜLLER  
Gross-Höflein, MNH-4-2, carapace 5.4×
6. *Dynomene emiliae* MÜLLER  
Visegrád, MV-4-2, carapace 5.4×



1



3



2



5




4



6

**PLATE XXXIV**

- 1—3. *Dorippe margaretha* LÖRENTHEY Holotype  
Rákos, unidentified layer, M 25 Fl, leg. LÖRENTHEY, internal mould of a carapace  $3.5 \times$   
4—5. *Dorippe margaretha* LÖRENTHEY  
Tétény-plateau, MDG-14-1,  $3.8 \times$




1

2



3




4



5


**PLATE XXXV**

- 1 -- 2. *Calappa praelata* LŐRENTHEY Holotype  
Mátraverebély-Szentkút, M 27 FI, leg. NOSZKY, carapace 2.6×
- 3 -- 5. *Calappa cf. praelata* LŐRENTHEY  
Oslip, sand-pit Bauer, MOS-1-1, leg. STEININGER and MÜLLER, right dactylus 4.5×
6. *Calappa cf. praelata* LŐRENTHEY  
Oslip, sand-pit Bauer, MOS-2-1, leg. STEININGER and MÜLLER, right dactylus with proximal lobe 4.5×
7. *Calappa praelata* LŐRENTHEY  
Mátraverebély-Szentkút, M 63/310 NMÖ, leg. STREDA, right dactylus 13×




**PLATE XXXVI**

1. *Calappa granulata* (LINNAEUS)  
Sicily, leg. SOLT, right carpus and propodus 2.6×
2. *Calappa granulata* (LINNAEUS)  
Sicily, leg. SOLT, carapace 2.2×
- 3—5. *Calappa granulata* (LINNAEUS)  
Sicily, leg. SOLT, right dactylus 5×
6. *Calappa cf. praelata* LÖRENTHEY  
Szob-Kerék-hegy, MSK-1-1, right propodus 3×



**PLATE XXXVII**

- 1—2. *Calappa heberti* BROCCHI  
Rákos, unidentified layer, M 26, FI, carapace 3.7×
3. *Calappa heberti* BROCCHI  
Rákos, unidentified layer, M 26, FI, carapace 1.2×
4. *Calappa heberti* BROCCHI  
Rákos, MR8-12-3, carapace 12×
- 5—7. *Calappa heberti* BROCCHI  
Erdősi út, MER-5-2, right dactylus 4×



1



2



3



6



4




5



7


**PLATE XXXVIII**

1. *Calappa heberti* BROCCHI  
Rákos, MRF-5-1, carapace with fragments of chelae 0.95×
2. *Calappa heberti* BROCCHI  
Rákos, unidentified layer, M 2379 FI, fixed finger of left propodus 1.6×
- 3—4. *Calappa heberti* BROCCHI  
Rákos, unidentified layer, M 26 FI, right chela with carpus 2.7× and 1.8×
5. *Calappa heberti* BROCCHI  
Gyakorló út, MGY-14-6, merus 3×
6. *Calappa heberti* BROCCHI  
Rákos, unidentified layer, M 2379 FI, left chela with carpus 1.6×




**PLATE XXXIX**


- 1—2. *Mursia lienharti* (BACHMAYER)  
Gyakorló út, MGY-17-1, leg. SOLT, carapace 5.6×
3. *Mursia lienharti* (BACHMAYER)  
Kerepesi út MKC-1-5, carapace 2.8×
4. *Mursia lienharti* (BACHMAYER)  
Rákos, MR8-20-5, left dactylus 8.5×
5. *Mursia lienharti* (BACHMAYER)  
Rákos, MR8-20-1, right propodus 8.5×
6. *Mursia lienharti* (BACHMAYER)  
Rákos, MR8-20-2, right propodus 4×




1



2



3



4




6




5

**PLATE XL**


- 1—3. *Mursia lienharti* (BACHMAYER)  
Örs vezér tere, MOE-33-1, right dactylus 7.5×
4. *Mursia armata* DE HAAN  
Indonesia, leg. and det. KREMPF, Mus. Nat. Hist. Nat. Paris, right dactylus 3.5×
5. *Mursia lienharti* (BACHMAYER)  
Rákos, MR8-12-3, fragment of right propodus with fixed finger 8×
6. *Mursia lienharti* (BACHMAYER)  
Rákos, MR8-12-4, left propodus 8×



1




5



2



4




3




6

**PLATE XLI**

1. *Matuta brocchii* GLAESSNER  
Gyakorló út, MGY-13-1, carapace 2.8×
- 2—3. *Matuta brocchii* GLAESSNER  
Gyakorló út, MGY-13-2, carapace 5.6×
- 4—5. *Matuta brocchii* GLAESSNER  
Rákos, MR45-2-1, carapace 3.7×
6. *Matuta brocchii* GLAESSNER  
Gyakorló út, MGY-13-10, stridulating ridges on left propodus 9×




1




4



3






5



6


**PLATE XLII**

- 1—3. *Matuta brocchii* GLAESSNER  
Gyakorló út, MGY-13-11, left carpus and propodus 4.8×  
4—5. *Matuta brocchii* GLAESSNER  
Gyakorló út, MGY-13-10, right dactylus 9×




**PLATE XLIII**

- 1—2. *Ebalia vanstraeleni* BACHMAYER Holotype  
Deutsch-Altenburg, NHMW, carapace (internal mould) 12×
- 3—4. *Ebalia oersi* MÜLLER Holotype  
Örs vezér tere, MOE-4-1, carapace 13.8×, 10×
- 5—7. *Ebalia oersi* MÜLLER  
Örs vezér tere MOE-4-10, carapace 11×
8. ?*Ebalia oersi* MÜLLER  
Örs vezér tere, MOE-4-11, female abdomen 11×




1

2




3




6



5



7



8

**PLATE XLIV**

1. *Ebalia* sp. 1.  
Törökmező, MTM-1-1, external mould of a carapace 14×
- 2—3. *Ebalia* sp. 2.  
Zebegény, MZZ-2-1, internal mould of a carapace 22×
4. *Ebalia* sp. 3.  
Zebegény, MZZ-3-1, internal mould of a carapace 22×
5. '*Ebalia*' *hungarica* MÜLLER Holotype  
Tétény-plateau, MB-3-1, carapace 10×



1




2



3




4




5

**PLATE XLV**


- 1—3. '*Ebalia*' *hungarica* MÜLLER  
Rákos, MR8-18-1, carapace 8.1×
- 4—5. '*Ebalia*' *hungarica* MÜLLER  
Tétény-plateau, MBH-5-1, carapace 6×




1



4



2



5



3

**PLATE XLVI**

- 1—3. *Andorina elegans* LÖRENTHEY Holotype  
Budapest, Telepy utca, M 30 FI, leg. MOESZ, carapace 7.7×
4. *Andorina elegans* LÖRENTHEY Holotype  
Budapest, Telepy utca, M 30 FI, leg. MOESZ, detail of surface 18×
5. ? '*Palaeomyra*' *globulosa* (MÜLLER)  
Budapest, Örs vezér tere, MOE-14-26, female pleon 8.8×
6. *Ranidina rosaliae* BITTNER  
Gorna Mitropolia, Bulgaria, coll. Univ. Sofia, leg. KOJUMDGIEVA, carapace 2.5×




2




1




3



5



6



4


**PLATE XLVII**

- 1—2. '*Palaeomyra*' *globulosa* (MÜLLER) Holotype  
Tétény-plateau, MA-1-1, carapace 9×
- 3—5. '*Palaeomyra*' *globulosa* (MÜLLER)  
Örs vezér tere, MOE-14-1, carapace 4.7×
6. '*Palaeomyra*' *globulosa* (MÜLLER)  
Örs vezér tere, MOE-14-2, carapace 6.2×
7. '*Palaeomyra*' *globulosa* (MÜLLER)  
Örs vezér tere, MOE-14-25, right carpus and propodus 8×




**PLATE XLVIII**

1. *Maja biaensis* LÖRENTHEY  
Bia, unidentified layer, MN-1-1, carapace 0.95×
2. *Maja biaensis* LÖRENTHEY  
Örs vezér tere, MOH-9-1, rostrum of carapace 1.1×
- 3—4. *Maja biaensis* LÖRENTHEY  
Gyakorló út, MGY-23-1, carapace 1.1×
- 5—6. *Maja biaensis* LÖRENTHEY  
Örs vezér tere, MOE-20-1, carapace 1.1×




**PLATE XLIX**


1. *Maja biaensis* LŐRENTHEY  
Bia, unidentified layer, FI, plaster cast of lost holotype, carapace  $0.95\times$
2. *Maja biaensis* LŐRENTHEY  
Gyakorló út, MGY-23-3, left propodus  $4\times$
3. *Maja biaensis* LŐRENTHEY  
Örs vezér tere, MOE-20-4, left dactylus  $3.2\times$
- 4—5. *Schizophrys visegradensis* n. sp. Holotype  
Törökmező, MTZ-20-1, carapace  $4.4\times$



1



2



3




4



5

**PLATE L**

1. *Schizophrys visegradensis* n. sp. Holotype  
Törökmező, MTZ-20-1, rubber cast of the external mould 4.4×
2. *Schizophrys visegradensis* n. sp. Paratype  
Törökmező, MTZ-20-2, rubber cast, carapace 4.4×
3. *Schizophrys visegradensis* n. sp. Paratype  
Törökmező, MTZ-20-3, rubber cast, carapace 4.4×
4. *Achaeus magnus* MÜLLER Holotype  
Örs vezér tere, MOE-5-1, carapace 3.8×
- 5—6. *Achaeus magnus* MÜLLER Paratype  
Örs vezér tere, MOE-5-4, right propodus 10×



2


1



3

5


4



6


**PLATE LI**

- 1—2. *Achaeus magnus* MÜLLER Paratype  
Örs vezér tere MOE-5-3, carapace 3.8×
- 3—4. 'Pisa' *oroszyi* (BACHMAYER)  
Tétény-plateau, MB-12-1, carapace 5.6×
- 5—6. 'Pisa' *oroszyi* (BACHMAYER)  
Tétény-plateau, MB-12-3, right carpus and propodus 5.6×



**PLATE LII**


- 1 — 2. '*Pisa' oroszyi* (BACHMAYER)  
Tétény-plateau, MB-12-2, carapace 4.7×
3. '*Pisa' oroszyi* (BACHMAYER)  
Tétény-plateau, MB-12-4, carapace 4.2×
4. *Micippa hungarica* (LŐRENTHEY) Holotype  
Bia, unidentified layer, M 2330 FI, carapace 3.2×
- 5 — 6. *Micippa hungarica* (LŐRENTHEY) var.?  
Rákos, unidentified layer, M 60/5964 NMÖ, leg. FRANZENAU, carapace 4.2×



2



1



4





5



6


**PLATE LIII**

1. *Micippa hungarica* (LŐRENTHEY)  
Örs vezér tere, MOE-17-1, carapace 3.1×
2. *Micippa hungarica* (LŐRENTHEY)  
Örs vezér tere, MOE-17-2, carapace 4.4×
- 3—4. *Micippa hungarica* (LŐRENTHEY)  
Örs vezér tere, MOE-17-3, carapace 5.2×
5. *Parthenope szaboi* MÜLLER Holotype  
Tétény-plateau, MDG-9-1, NMÖ, carapace 7×




1


2



3



4



5

**PLATE LIV**

1—2. *Parthenope szaboi* MÜLLER

Örs vezér tere, MOE-15-1, carapace 6.2×

3. *Parthenope szaboi* MÜLLER

Örs vezér tere, MOE-15-2, carapace 4×

4—5. *Parthenope szaboi* MÜLLER

Örs vezér tere, MOE-15-11, left (nipper) dactylus 10×

6. *Parthenope szaboi* MÜLLER


Rákos, MR8-13-1, right propodus 8×

7. *Parthenope szaboi* MÜLLER

Örs vezér tere, MOE-15-12, left (nipper) propodus and carpus 5.5×



8—9. *Parthenope szaboi* MÜLLER

Örs vezér tere, MOE-15-13, left propodus 7.3×



**PLATE LV**

- 1—2. *Parthenope tetenyensis* n. sp. Holotype  
Örs vezér tere, MOE-16-1, carapace 4.2×
- 3—4. *Parthenope tetenyensis* n. sp. Paratype  
Örs vezér tere, MOE-16-2, carapace 3×
- 5—6. *Parthenope tetenyensis* n. sp. Paratype  
Örs vezér tere, MOE-16-3, carapace 10×
7. *Parthenope tetenyensis* n. sp. Paratype  
Örs vezér tere, MOE-16-4, carapace 6.8×



1




4



3



5




7



6

**PLATE LVI**


1. *Parthenope tetenyensis* n. sp. Paratype  
Örs vezér tere, MOE-16-16, right (crusher) chela 8×
2. *Parthenope tetenyensis* n. sp. Paratype  
Osip (Burgenland), sand-pit Bauer, MOS-3-1, leg. STEININGER and MÜLLER, left (crusher!) dactylus 9×
3. *Parthenope tetenyensis* n. sp. Paratype  
Örs vezér tere, MOE-16-17, left (nipper) propodus 6×
4. *Parthenope tetenyensis* n. sp. Paratype  
Örs vezér tere, MOE-16-18, left (nipper) dactylus 7×
- 5—7. *Parthenope tetenyensis* n. sp. Paratype  
Örs vezér tere, MOE-16-19, right (crusher) dactylus 7×




1




3



4



5



7



6

**PLATE LVII**


1. *Cancer styriacus* BITTNER  
Örs vezér tere, MOE-27-1, carapace 2.8×
2. *Cancer styriacus* BITTNER  
Diósd, MDZ-4-2, rubber cast of external mould, right lateral margin of carapace 6×
- 3—4. *Cancer styriacus* BITTNER  
Diósd, MDZ-4-1, carapace 2.5×




1



2





3



4

**PLATE LVIII**

1. *Cancer styriacus* BITTNER  
Örs vezér tere, MOE-27-3, left propodus 5.1×
2. *Cancer styriacus* BITTNER  
Örs vezér tere, MOE-27-4, right propodus, upper view 6×
3. *Cancer illyricus* BITTNER (? = *C. sismondai* MEYER) Holotype  
Zagor, Yugoslavia, coll. Univ. Graz, plaster cast of carapace 1.7×
4. *Miocyclus bulgaricus* MÜLLER Paratype  
Oriakhovo, Bulgaria, coll. Univ. Sofia, leg. KOJUMDGIEVA, No. Nt 533/b, carapace 2×
5. *Miocyclus bulgaricus* MÜLLER Holotype  
Oriakhovo, Bulgaria, coll. Univ. Sofia, leg. KOJUMDGIEVA, No. Nt 533/a, carapace 2×
6. *Miocyclus bulgaricus* MÜLLER Holotype  
Oriakhovo, Bulgaria, coll. Univ. Sofia, leg. KOJUMDGIEVA, No. Nt 533/a, part of right antero-lateral margin 7×



**PLATE LIX**

- 1—3. “*Cancer*” *szontaghii* LÖRENTHEY Holotype  
Tasádfő, Transsylvania, Rumania. M 35 FI. leg. SzONTÁGH, complete specimen 1.7×
4. “*Cancer*” *szontaghii* LÖRENTHEY Holotype  
Tasádfő, Transsylvania, Rumania, M 35 FI. leg. SzONTÁGH, right chela 2×
5. “*Cancer*” *szontaghii* LÖRENTHEY  
Tasádfő, M 2451 FI. leg. SZONTÁGH, frontal part of carapace



1




2



3




4



5

**PLATE LX**

1. "Cancer" *szontaghii* LÖRENTHEY  
same specimen as on pl. LIX. fig. 1., basis-ischium of right 3rd maxilliped 9×
2. "Cancer" *szontaghii* LÖRENTHEY  
same specimen as on pl. LIX. fig. 5., ventral side 2.5×
- 3–5. *Pirimela lorentheyi* (MÜLLER) Holotype  
Tétény-plateau, MB-4-1, carapace 3.8×



1




2



3





4



5

**PLATE LXI**

- 1—2. *Trachypirimela grippi* (MÜLLER)  
Holotype of *T. radula*  
Tétény-plateau, MDG-15-2 (= MA-1-1), NMÖ, carapace 12×
3. *Trachypirimela grippi* (MÜLLER)  
Örs vezér tere, MOE-28-1, carapace, lateral view 12×
4. *Trachypirimela grippi* (MÜLLER)  
Örs vezér tere, MOE-28-1, carapace, dorsal view 15×
5. *Trachypirimela grippi* (MÜLLER) Holotype  
Tétény-plateau, MDG-15-1 (= MG-3-1), carapace 9×
6. *Trachypirimela grippi* (MÜLLER)  
Tétény-plateau, MJL-12-1, carapace 8×



1

2

3



4




6



5

**PLATE LXII**


- 1—2. *Portunus monspeliensis* (A. MILNE-EDWARDS)  
Kishajmás-Szatina, railway cut, MKS-1-1, leg. SOLT, carapace, ventral side 1.5×
- 3—4. *Portunus neogenicus* MÜLLER Holotype  
Gyakorló út, MGY-3-1, carapace 3.8×
5. *Portunus (Monomia) miocaenicus* n. sp. Holotype  
Rákos, MRO-10-1, carapace 3.8×




1



2




3



5

**PLATE LXIII**


- 1, 4. *Charybdis (Goniosupradens) mathiasi* n. sp. Holotype  
Rákos, MRZ-5-1, carapace 1.7 $\times$
2. *Charybdis (Goniosupradens) mathiasi* n. sp. Paratype  
Rákos, MRZ-5-2, carapace 1.1 $\times$
3. *Charybdis (Goniosupradens) mathiasi* n. sp. Paratype  
Rákos, MRZ-5-3, rubber cast, carapace 1.4 $\times$




1



2




3




4

**PLATE LXIV**

1. *Charybdis (Goniosupradens) mathiasi* n. sp.  
same specimen as on pl. LXIII, fig. 3., postfrontal part  $5\times$
2. *Charybdis (Goniosupradens) mathiasi* n. sp. Paratype  
Rákos, MRZ-5-11, right chela  $3.6\times$
- 3-5. *Charybdis (Goniosupradens) mathiasi* n. sp. Paratype  
Rákos, MRZ-5-12, left chela with carpus  $1.95\times$



1




2



3



4



5


**PLATE LXV**

1. *Thalamita fragilis* MÜLLER Holotype  
Gyakorló út, MGY-4-1, fragment of carapace 5.2×
2. *Thalamita fragilis* MÜLLER  
Rákos, MRF-10-1, carapace 2.9×
3. ?*Thalamita fragilis* MÜLLER  
Gyakorló út, MGY-4-4, left part of a carapace 6.5×
4. *Thalamita fragilis* MÜLLER  
Rákos, MRO-8-1, carapace 3×
5. *Thalamita fragilis* MÜLLER (subspecies?)  
Rákos, MR8-17-1, carapace 4.9×



1

2



3


4



5


**PLATE LXVI**

- 1—2. *Thalamita fragilis* MÜLLER  
Rákos, MRF-10-5, right chela with carpus 4.5×
- 3—6. *Thalamita fragilis* MÜLLER  
Örs vezér tere, MOH-11-1, -2, right fixed fingers 5.1×
7. *Thalamita fragilis* MÜLLER  
Gyakorló út, MGY-4-6, right dactylus 6.1×
8. *Necronectes schafferi* GLAESSNER  
Diósd, MDZ-10-1, carapace, median regions 1.05×
9. *Necronectes schafferi* GLAESSNER  
Diósd, MDZ-10-1, carapace, frontal part 1.9×



**PLATE LXVII**

- 1—2. *Thalamita* n? sp.  
Rákos, MRZ-13-1, carapace 2.6×
- 3—5. *Charybdis (Goniosupradens) mathiasi* n. sp.  
Rákos, MRZ-5-12, right propodus with carpus 1.7×



1



2



3

4

5


**PLATE LXVIII**

- 1—2. *Rakosia carupoides* n. gen. n. sp. Holotype  
Gross-Höflein, NHMW, rubber cast of external mould, carapace  $4.5\times$
3. *Rakosia carupoides* n. gen. n. sp. Paratype  
Rákos, MRZ-14-1, carapace  $4.1\times$
4. *Rakosia carupoides* n. gen. n. sp. Paratype  
Rákos, MRZ-14-16, left propodus with fragment of dactylus  $5.4\times$
5. *Rakosia carupoides* n. gen. n. sp. Paratype  
Rákos, MRZ-14-17, left propodus with dactylus  $5.4\times$
- 6—7. *Rakosia carupoides* n. gen. n. sp. Paratype  
Rákos, MRZ-14-18, right chela with dactylus and carpus  $3.6\times$



1

3




2



4



6




7



5


## PLATE LXIX

1. ?*Necronectes schafferi* GLAESSNER  
Rákos, MRZ-19-1, right chela 1.8×
2. *Liocarcinus rakosensis* (LÖRENTHEY) Lectotype  
Rákos, unidentified layer, M 2328/a FI, leg. LÖRENTHEY, carapace 4.5×
- 3—4. *Liocarcinus rakosensis* (LÖRENTHEY)  
Örs vezér tere, MOE-21-1, carapace with straight frontal margin 3.8×
5. *Liocarcinus rakosensis* (LÖRENTHEY)  
Örs vezér tere, MOE-21-2, carapace with typical front 5.9×
6. *Liocarcinus rakosensis* (LÖRENTHEY)  
Örs vezér tere, MOE-21-3, carapace 5.9×






**PLATE LXX**

1. *Liocarcinus rakosensis* (LÖRENTHEY)  
Örs vezér tere, MOE-21-46, right propodus with carpus 4.6×
2. *Liocarcinus rakosensis* (LÖRENTHEY)  
Örs vezér tere, MOH-10-1, left palm 7.5×
3. *Liocarcinus rakosensis* (LÖRENTHEY)  
Örs vezér tere, MOE-21-47, left (crusher) dactylus 6.9×
4. *Liocarcinus rakosensis* (LÖRENTHEY)  
Örs vezér tere, MOE-21-48, left (nipper) propodus 7.7×
5. *Liocarcinus rakosensis* (LÖRENTHEY)  
Örs vezér tere, MOE-21-49, left (nipper) dactylus 8.5×
6. *Liocarcinus rakosensis* (LÖRENTHEY)  
Örs vezér tere, MOE-21-50, left (nipper) fixed finger 8.5×
- 7—8. *Liocarcinus rakosensis* (LÖRENTHEY)  
Örs vezér tere, MOE-21-51, right dactylus 8.5×



**PLATE LXXI**


- 1—2. *Liocarcinus oroszyi* (BACHMAYER) Holotype  
Deutsch-Altenburg, NHMW, leg. K. OROSZY, carapace 6.1×
- 3—4. *Liocarcinus kuchni* (BACHMAYER) Holotype  
Deutsch-Altenburg, NHMW, carapace 5.5×



4

**PLATE LXXII**

- 1—3. *Miopipus pygmeus* (BROCCHI) n. gen.  
Keresztúri út, MEG-16-1, carapace, 10.1×
4. *Miopipus pygmeus* (BROCCHI)  
Keresztúri út, MEG-16-2, 10.1×
- 5—7. 'Xaiva' *bachmayeri* n. sp.  
Örs vezér tere, MOE-26-1, carapace 4.8×
8. ?'Xaiva' *bachmayeri* n. sp.  
Örs vezér tere, MOE-26-3, left propodus 7.5×




1




2



3




5



4



8




7




6

**PLATE LXXIII**


- 1—3. *Mioxaiva psammophila* MÜLLER Holotype Gyakorló út, MGY-5-1, carapace, the right side of the mosaic picture is a mirror image 11×
4. ‘*Lissocarcinus*’ *szoeraenyiae* (MÜLLER) Diósd, MDN-3-1, carapace, internal mould 5.2×
- 5—6. ‘*Lissocarcinus*’ *szoeraenyiae* (MÜLLER) Holotype Tétény-plateau, MDG-18-1 (=MD-1-1), NMÖ, carapace 3.4×
7. ?*Carupa* cf. *tenuipes* DANA Visegrád, MV-6-1, carapace 2.7×




1




2




4




3



5



6




7

**PLATE LXXIV**


- 1—2. *Actaea turcocampestris* n. sp. Holotype  
Fertőrákos, drilling FRK-21, 268 m, MFR-1-1, carapace 11×
- 3—4. *Actaea turcocampestris* n. sp. Paratype  
Törökmező, MTZ-10-4, right chela 5.8×
5. *Actaea turcocampestris* n. sp. Paratype  
Törökmező, MTZ-10-1, rubber cast, carapace 8.6×




1



3



4




2



5

**PLATE LXXV**

1. *Carpilius antiquus* GLAESSNER Holotype  
Baden-Rauchstallbrunnengraben, No 1860/8, NHMW, internal mould, carapace 2.2×
- 2—3. *Carpilius antiquus* GLAESSNER  
Törökmező, MÁTZ-21-1, internal mould, carapace 2.3×
4. *Carpilius antiquus* GLAESSNER  
Visegrád, MV-11-2, front of carapace 6.8×
5. *Carpilius antiquus* GLAESSNER  
Visegrád, MV-11-9, right propodus 2×




**PLATE LXXVI**

1. *Chlorodiella mediterranea* (LÖRENTHEY) Lectotype  
Rákos, M 2327/a FI, same layer as MRZ (coral-layer), carapace 5.5×
2. *Chlorodiella mediterranea* (LÖRENTHEY)  
Rákos, MRZ-6-2, carapace 6.8×
- 3—4. *Chlorodiella mediterranea* (LÖRENTHEY)  
Rákos, MRZ-6-1, carapace 4.2×
5. *Chlorodiella mediterranea* (LÖRENTHEY)  
Rákos, MRZ-6-101, right propodus with carpus 5.3×
6. *Chlorodiella mediterranea* (LÖRENTHEY)  
Rákos, MRZ-6-102, left chela with carpus 5.3×




**PLATE LXXVII**

1. *Chlorodiella mediterranea* (LÖRENTHEY)  
same specimen as on pl. LXXVI. fig. 5.
2. *Chlorodiella mediterranea* (LÖRENTHEY)  
same specimen as on pl. LXXVI. fig. 6.
- 3—4. *Chlorodiella mediterranea* (LÖRENTHEY)  
Rákos, MRZ-6-103, right dactylus 6.3×
5. *Chlorodiella mediterranea* (LÖRENTHEY) *tetenensis* n. ssp. Holotype  
Rákos, MRZ-19-1, carapace 8.5×
- 6—7. *Chlorodiella mediterranea* (LÖRENTHEY) *tetenensis* n. ssp. Paratype  
Rákos, MRZ-19-2, carapace 4×



1



2



3



5



4




6



7

**PLATE LXXVIII**


- 1—3. *Chlorodiella loczyi* n. sp. Holotype  
Rákos, MRZ-7-1, carapace 13.6×
4. *Chlorodiella loczyi* n. sp. Paratype  
Rákos, MRZ-7-3, carapace 12×
5. *Chlorodiella juglans* n. sp. Holotype  
Diósd, MDZ-11-1, carapace 6.3×
6. *Chlorodiella juglans* n. sp. Holotype  
Diósd, MDZ-11-1, cast of external mould 6.3×




1



2



3



4




5

**PLATE LXXIX**


1—2. *Daira speciosa* (REUSS)  
Rákos, MRZ-12-1, carapace 4.2×

3. *Daira speciosa* (REUSS)  
Törökmező, MTZ-6-2, cast of carapace 2.9×


4—6. *Daira speciosa* (REUSS)  
Törökmező, MTZ-6-4, right palm and carpus, internal mould and cast of external mould 3×



1



4




2



3




5



6

**PLATE LXXX**

1. *Daira speciosa* (REUSS)  
Törökmező, MTZ-6-1, internal mould of carapace 2.5×
2. *Daira speciosa* (REUSS)  
Rákos, MRZ-12-2, surface of carapace 10×
- 3—5. *Haydnella steiningeri* n. gen. n. sp. Paratype  
Gross-Höflein, MNH-9-3, right chela with carpus 4.9×




2




1



3



4



5

**PLATE LXXXI**

- 1--3. *Haydnella steiningeri* n. gen. n. sp. Holotype  
Gross-Höflein, NHMW, carapace 7.9×
4. *Haydnella steiningeri* n. gen. n. sp. Paratype  
Gross-Höflein, MNH-9-2, cast of external mould, carapace 5.9×
- 5--6. *Panopeus wronai* n. sp. Paratype  
Gross-Höflein, MNH-10-6, fingers 8.6×




1




3



2



5




6



4

**PLATE LXXXII**

- 1—2. *Panopeus wronai* n. sp. Holotype  
Gross-Höflein, NHMW, cast of external mould 1.8×  
3—4. *Panopeus wronai* n. sp. Paratype  
Gross-Höflein, MNH-10-2, cast, carapace 2.9×



4

3

**PLATE LXXXIII**


1. *Panopeus wronai* n. sp.  
same specimen as on pl. LXXXI. figs. 5., 6, rubber cast of palm, original fingers 5×
- 2-4. *Panopeus wronai* n. sp. Paratype  
Gross-Höflein, MNH-10-2, internal mould of carapace, cast of chela 2.9×
5. 'Pilodius' *vulgaris* (GLAESSNER) Holotype  
Baden-Rauchstallbrunngaben, No 1927 I 2, NHMW, carapace 4.8×
6. 'Pilodius' *vulgaris* (GLAESSNER)  
Zebegény, MZZ-7-1, carapace 11.5×



1



3




2



4




5




6

**PLATE LXXXIV**


- 1 — 2. '*Pilodius*' *vulgaris* (GLAESSNER)  
Zebegény, MZZ-7-8, right chela with carpus 9.5×
3. '*Pilodius*' *vulgaris* (GLAESSNER)  
Zebegény, MZZ-7-9, right palm 7.4×
4. '*Pilodius*' *vulgaris* (GLAESSNER)  
Zebegény, MZZ-7-2, frontal part of carapace 7.4×
5. *Trapezia glaessneri* MÜLLER  
Visegrád, MV-1-12, merus of right cheliped 10.5×
6. *Trapezia glaessneri* MÜLLER  
Visegrád, MV-1-5, fragment of left propodus 6×




1




2



3



4




5



6

**PLATE LXXXV**

1. *Trapezia glaessneri* MÜLLER Holotype  
Visegrád, MV-1-1, part about the right orbit 10.5×
2. *Trapezia glaessneri* MÜLLER Holotype  
Visegrád, MV-1-1, carapace 5×
3. *Trapezia glaessneri* MÜLLER Paratype  
Visegrád, MV-1-2, carapace 7.8×
4. *Trapezia glaessneri* MÜLLER Paratype  
Visegrád, MV-1-3, carapace 4.5×
- 5—7. *Xantho moldavicus* (YANAKEVICH)  
Węglinek, Poland, MWE-3-3, right (crusher) dactylus 7.5×
8. *Xantho moldavicus* (YANAKEVICH)  
Węglinek, Poland, MWE-3-4, leg. KRACH, right (crusher) fixed finger 7.5×



3

4

2

5


6

7

8

**PLATE LXXXVI**

- 1 - 2. *Xantho moldavicus* (YANAKEVICH)  
Węglinek, Poland, MWE-3-1, carapace, leg. KRACH 5.9×
3. *Xantho moldavicus* (YANAKEVICH)  
Örs vezér tere, MOE-31-8, right chela with propodus 9.5×
4. *Xantho moldavicus* (YANAKEVICH)  
Diósd, MDZ-8-1, rubber cast, carapace 4.9×
5. *Xantho moldavicus* (YANAKEVICH)  
Węglinek, Poland, MWE-3-5, right (nipper) dactylus 7.5×




1




2



4



3




5

**PLATE LXXXVII**


1. *Xantho moldavicus* (YANAKEVICH)  
Kerepesi út, MKC-19-1, carapace 13.8×
2. *Pilumnus mediterraneus* (LÖRENTHEY)  
Örs vezér tere, MOE-18-1, carapace 13.5×
3. *Pilumnus mediterraneus* (LÖRENTHEY) Lectotype  
Rákos, unidentified layer, M 31/a FI, leg. LÖRENTHEY carapace 4.95×
4. *Pilumnus mediterraneus* (LÖRENTHEY)  
Örs vezér tere, MOE-18-57, right chela 7×
5. *Pilumnus mediterraneus* (LÖRENTHEY)  
Örs vezér tere, MOE-18-58, right dactylus 7.8×




1



2



3



4



5

**PLATE LXXXVIII**

1. *Pilumnus mediterraneus* (LÖRENTHEY)

Örs vezér tere, MOE-18-2, carapace 3.2×

- 2—3. *Pilumnus mediterraneus* (LÖRENTHEY)


Örs vezér tere, MOE-18-3, carapace 5×

4. *Pilumnus mediterraneus* (LÖRENTHEY)

Örs vezér tere, MOE-18-59, left (nipper) chela with carpus 7.8×

5. *Pilumnus mediterraneus* (LÖRENTHEY)

Örs vezér tere, MOE-18-60, right chela 8.4×




1




2



3






4



5

**PLATE LXXXIX**

1. *Actumnus telegdii* (MÜLLER) Holotype  
Tétény-plateau, MDG-23-1 (MG-4-1), carapace 12×
- 2—3. *Actumnus telegdii* (MÜLLER)  
Örs vezér tere, MOE-19-1, carapace 4.1×
4. *Actumnus telegdii* (MÜLLER)  
Örs vezér tere, MOE-19-29, left (nipper) chela with carpus 7×
5. *Actumnus telegdii* (MÜLLER)  
Örs vezér tere, MOE-19-30, left (crusher!) chela 9.4×



3

5

**PLATE XC**

1—2. *Actumnus telegdii* (MÜLLER)

Örs vezér tere, MOE-19-2, carapace 6×

3. *Actumnus telegdii* (MÜLLER)


Örs vezér tere, MOE-19-31, right (crusher) chela with carpus 9.2×

4—5. *Actumnus telegdii* (MÜLLER)

Örs vezér tere, MOE-19-32, left (crusher) dactylus 7.2×

6—7. *Actumnus* n? sp.

Zebegény, MZZ-5-1, carapace 13.5×



3

2

5


4

7


6

**PLATE XCI**

- 1—3. '*Glabropilumnus*' n. sp.  
Törökmező, MTZ-8-1, carapace, fig. 3. rubber cast 6.7×
- 4—5. '*Pilumnopeus*' *tetenyensis* n. sp. Holotype  
Tétény-plateau, MB-16-1, carapace 12.7×




1




3



2




5



4

**PLATE XCII**


1. *Pilumnopeus paratethyensis* n. sp. Holotype  
Rákos, MRZ-10-1, carapace 16.2×
2. *Pilumnopeus paratethyensis* n. sp. Paratype  
Rákos, MRZ-10-2, front of carapace 16.2×
- 3—4. *Pilumnopeus paratethyensis* n. sp. Paratype  
Rákos, MRZ-10-3, carapace 10.3×
5. *Pilumnopeus paratethyensis* n. sp. Paratype  
Rákos, MRZ-10-4, carapace 11×
6. ?*Pilumnopeus paratethyensis* n. sp.  
Rákos, MRZ-10-18, right palm with carpus 11×



1




3



2



4



5



6

**PLATE XCIII**

1. *Geryon* cf. *latifrons* VAN STRAELEN


Thököly út, drilling Z-2, 36.3 m, MTT-1-1, carapace  $3.2\times$

- 2—3. *Goneplax* cf. *gulderi* BACHMAYER


Oriakhovo, Bulgaria, coll. Univ. Sofia, leg. KOJUMDGIEVA, carapace  $5\times$

4. *Pachygrapsus hungaricus* MÜLLER


Rákos, MRZ-4-18, left propodus  $6.7\times$



1



2




3




4

PLATE XCIV


- 1—2. *Pachygrapsus hungaricus* MÜLLER Holotype  
Tétény-plateau, MB-5-1, carapace 10×
3. *Pachygrapsus hungaricus* MÜLLER Paratype  
Tétény-plateau, MB-5-2, carapace 6.2×
4. *Pachygrapsus marmoratus* (FABRICIUS)  
Adriatic Sea, leg. MÜLLER, carapace 4.8×
5. *Pachygrapsus hungaricus* MÜLLER  
Rákos, MRZ-4-1, carapace 4.8×
6. *Pachygrapsus hungaricus* MÜLLER  
Rákos, MRZ-4-2, carapace 9.5×
7. *Pachygrapsus hungaricus* MÜLLER Paratype  
Tétény-plateau, MB-5-3, carapace 6.2×




1



2




3



4



5




6




7

**PLATE XCV**


- 1—2. *Brachynotus februarius* MÜLLER Holotype  
Tétény-plateau, MB-6-1, NMÖ, carapace 11×
3. *Brachynotus februarius* MÜLLER Paratype  
Tétény-plateau, MB-6-2, carapace 6.8×
4. *Brachynotus februarius* MÜLLER  
Tétény-plateau, MB-6-4, carapace 8×
5. *Brachynotus februarius* MÜLLER Paratype  
Tétény-plateau, MB-6-3, carapace 13×
6. *Macrophthalmus vindobonensis* GLAESSNER  
Mező Imre út drilling H-34, 32.6 m, MH34-1-4, right fixed finger 13.6×



1



3




2



4



5



6

**PLATE XCVI**

1. *Macrophthalmus vindobonensis* GLAESSNER  
Mező Imre út drilling H-34, 33 m, MH34-1-1, carapace 4.9×
2. *Macrophthalmus vindobonensis* GLAESSNER  
Mező Imre út drilling H-34, 33 m, MH34-1-1, posterolateral part 10×
3. *Macrophthalmus aquensis* A. MILNE-EDWARDS-BROCCHI  
Aix, Syntype, Mus. Natl. Hist. Nat. Paris, rubber cast, carapace 5.1×
4. *Macrophthalmus vindobonensis* GLAESSNER  
Mező Imre út drilling H-34, 25.8 m, MH34-1-2, carapace 4.9×
5. *Macrophthalmus vindobonensis* GLAESSNER  
Mező Imre út drilling H-34, 32.6 m, MH34-1-5, left dactylus 13.6×



1




2



3



4



5

**PLATE XCVII**

- 1., 3. *Crossotonotus diosdensis* n. sp. Holotype  
Diósd, MDZ-15-1, carapace 1.5×
2. *Crossotonotus diosdensis* n. sp. Holotype  
Diósd, MDZ-15-1, rubber cast of external mould 1.5×
4. *Crossotonotus diosdensis* n. sp. Paratype  
Diósd, MDZ-15-2, rubber cast of external mould 6×
5. *Crossotonotus diosdensis* n. sp. Paratype  
Diósd, MDZ-15-2, carapace 6×
6. *Palicus* sp.  
Örs vezér tere, MOE-32-2, right side, mirror image, carapace 12×
7. *Palicus* sp.  
Örs vezér tere, MOE-32-1, rubber cast of inner surface of poorly preserved carapace 15×

