



Dr. Péter Ozsvárt is geologist and paleontologist. He is a member of the Research Group for Paleontology at the Hungarian Academy of Sciences and Hungarian Natural History Museum in Budapest. His doctoral thesis is on paleoecology of Eocene benthic foraminifera and their paleoceanographic applications from the Hungarian Paleogene Basin. Currently his research is focused on taxonomy, biostratigraphy and paleoecology of Triassic and Jurassic radiolarians from Hungary, Greece and Turkey.

*Middle and Late Eocene benthic foraminiferal fauna from the Hungarian Paleogene Basin: systematics and paleoecology* presents a synthesis about the well-preserved and in most cases incredibly diverse Eocene benthic foraminiferal faunas from the Hungarian Paleogene Basin. This report provides a brief review of geological setting and biostratigraphy of the Paleogene Basin and provides comprehensive coverage of the systematic analysis, paleoecology and paleoenvironmental interpretations of the Eocene benthic foraminiferal fauna from Hungary.

  
Hantken Press  
Budapest, 2007




Middle and Late Eocene benthic foraminiferal fauna, Hungarian Paleogene Basin

Geologica Pannonica Special Publication 2

# Middle and Late Eocene benthic foraminiferal fauna from the Hungarian Paleogene Basin: systematics and paleoecology

Péter Ozsvárt



  
Hantken Press  
Budapest, 2007

Geologica Pannonica Spec. Publ. 2

PÉTER OZSVÁRT

Middle and Late Eocene  
benthic foraminiferal fauna of the  
Hungarian Paleogene Basin:  
systematics and paleoecology

## Geologica Pannonica Special Publications

1. HORVÁTH, F. & GALÁ CZ, A. (eds) (2006): The Carpathian-Pannonian Region – A Review of Mesozoic-Cenozoic Stratigraphy and Tectonics. Vols 1–2. 650 p., 260 figs, 3 coloured maps
2. OZSVÁ RT, P (2007): Middle and Late Eocene Benthic foraminiferal fauna of the Hungarian Paleogene Basin: systematics and paleoecology. 127 p., 20 figs, 2 t., 11 plates

### Forthcoming

TARI, G (ed.): Extensional collapse of the Alpine orogene and hydrocarbon prospects in the basement and basin fill of the western Pannonian Basin  
The Pannonian and Transylvanian Basins  
Hydrocarbon geology of the Carpathian-Pannonian region

Geologica Pannonica Special Publication 2

Middle and Late Eocene benthic foraminiferal fauna  
of the Hungarian Paleogene Basin:  
systematics and paleoecology

PÉTER OZSVÁRT



Hantken Press  
Budapest  
2007

PÉTER OZSVÁRT  
Hungarian Academy of Sciences–Hungarian Natural History Museum  
Research Group for Paleontology  
P.O.Box 137, H-1431 Budapest, Hungary  
Email: ozsi@nhmus.hu

This volume is available from

The Hantken Press  
P O Box 23  
H-1241 Budapest  
Hungary

E-mail: hantkenpress@gmail.com  
<http://hantkenpress.fw.hu>

Sponsored by  
Book and Journals Committee  
of the Hungarian Academy of Sciences

Suggested reference to this volume:

OZSVÁRT P. (2007): Middle and Late Eocene benthic foraminiferal fauna from the Hungarian Paleogene Basin: systematics and paleoecology. – *Geologica Pannonica Special Publication* 2, 1–127, 20 figs, 2 tables, 11 plates, Hantken Press, Budapest

ISBN 978-963-87436-4-0  
ISSN HU 1788-4004

© PÉTER OZSVÁRT 2007  
© Hantken Kiadó, Budapest, 2007

## Contents

7	Introduction	42	
7	Previous work	42	
8	Geological setting	42	
8	Localities	44	
9	Biostratigraphy	44	
10	Methods	45	
12	Sections	45	
12	<i>SW Bakony - Zala Paleogene Subbasin</i>	46	
16	<i>NE Bakony – Vértes Paleogene Subbasin</i>	46	
22	<i>Gerecse Paleogene Subbasin</i>	46	
25	<i>North Hungarian Paleogene Subbasin</i>	47	
27	Paleoecology	47	
27	<i>Benthic foraminiferal assemblages</i>	47	
27	<i>Water temperature and salinity</i>	50	
28	Acknowledgements	52	
29	Systematic paleontology	52	
29	Family Astrorhizidae	52	
29	Genus <i>Astrorhiza</i>	53	
29	Family Bathysiphonidae	53	
29	Genus <i>Bathysiphon</i>	55	
30	Family Rhabdamminidae	55	
30	Genus <i>Rhabdammina</i>	56	
31	Family Hormosinidae	58	
31	Genus <i>Reophax</i>	60	
32	Family Haplophragmoididae	60	
32	Genus <i>Haplophragmoides</i>	61	
32	Family Lituolidae	61	
32	Genus <i>Ammomarginulina</i>	62	
32	Family Spiroplectamminidae	62	
32	Genus <i>Bolivinopsis</i>	63	
33	Genus <i>Spiroplectammina</i>	63	
33	Genus <i>Vulvulina</i>	63	
34	Family Prolixoplectidae	64	
34	Genus <i>Plectina</i>	64	
35	Family Verneuilinidae	65	
35	Genus <i>Verneuilina</i>	65	
35	Family Eggerellidae	66	
35	Genus <i>Dorothia</i>	66	
35	Genus <i>Martinottiella</i>	66	
36	Family Textulariidae	67	
36	Genus <i>Textularia</i>	68	
37	Family Pseudogaudryinidae	68	
37	Genus <i>Clavulinoides</i>	71	
39	Family Valvulamminidae	71	
39	Genus <i>Clavulina</i>	72	
40	Genus <i>Cylindroclavulina</i>	72	
40	Genus <i>Tritaxilina</i>	72	
41	Family Cornuspiridae	73	
41	Genus <i>Cornuspira</i>	73	
41	Family Spiroloculinidae	74	
41	Genus <i>Adelosina</i>	75	
			Genus <i>Spiroloculina</i>
			Family Hauerinidae
			Genus <i>Quinqueloculina</i>
			Genus <i>Pyrgo</i>
			Genus <i>Articulina</i>
			Family Miliolidae
			Genus <i>Miliolina</i>
			Genus <i>Massilina</i>
			Family Peneroplidae
			Genus <i>Spirolina</i>
			Family Nodosariidae
			Genus <i>Chrysalogonium</i>
			Genus <i>Dentalina</i>
			Genus <i>Nodosaria</i>
			Genus <i>Pseudonodosaria</i>
			Genus <i>Gonatosphaera</i>
			Genus <i>Fronicularia</i>
			Family Vaginulinidae
			Genus <i>Lenticulina</i>
			Genus <i>Marginulinopsis</i>
			Genus <i>Saracenaria</i>
			Genus <i>Marginulina</i>
			Genus <i>Vaginulinopsis</i>
			Genus <i>Planularia</i>
			Genus <i>Vaginulina</i>
			Family Lagenidae
			Genus <i>Lagena</i>
			Family Polymorphinidae
			Genus <i>Globulina</i>
			Genus <i>Ramulina</i>
			Family Ellipsolagenidae
			Genus <i>Fissurina</i>
			Family Glandulinidae
			Genus <i>Glandulina</i>
			Family Bolivinidae
			Genus <i>Bolivina</i>
			Family Cassidulinidae
			Genus <i>Globocassidulina</i>
			Family Buliminidae
			Genus <i>Bulimina</i>
			Family Uvigerinidae
			Genus <i>Uvigerina</i>
			Family Reussellidae
			Genus <i>Reussella</i>
			Family Fursenkoinidae
			Genus <i>Fursenkoina</i>
			Family Pleurostomellidae
			Genus <i>Ellipsoglandulina</i>
			Genus <i>Nodosarella</i>
			Genus <i>Pleurostomella</i>
			Family Stilostomellidae

75	Genus <i>Orthomorphina</i>	87	Genus <i>Pullenia</i>
75	Genus <i>Stilostomella</i>	88	Family Almaenidae
78	Family Bagginidae	88	Genus <i>Almaena</i>
78	Genus <i>Cancris</i>	88	Genus <i>Queraltina</i>
78	Genus <i>Valvulineria</i>	88	Family Chilostomellidae
78	Family Eponididae	88	Genus <i>Chilostomella</i>
78	Genus <i>Eponides</i>	89	Family Heterolepidae
79	Family Discorbidae	89	Genus <i>Anomalinoidea</i>
79	Genus <i>Discorbis</i>	90	Genus <i>Heterolepa</i>
80	Family Parrelloididae	91	Family Gavelinellidae
80	Genus <i>Cibicidoides</i>	91	Genus <i>Gyroidinoidea</i>
81	Family Planulinidae	92	Family Gavelinellinae
81	Genus <i>Planulina</i>	92	Genus <i>Gavelinella</i>
81	Family Cibicididae	92	Genus <i>Hanzawaia</i>
81	Genus <i>Cibicides</i>	93	Family Linderinidae
85	Genus <i>Dyocibicides</i>	93	Genus <i>Eoannularia</i>
85	Family Asterigerinidae	93	Family Rotaliidae
85	Genus <i>Asterigerina</i>	93	Genus <i>Pararotalia</i>
85	Family Epistomariidae	93	Genus <i>Rotalia</i>
85	Genus <i>Nuttallides</i>	94	Genus <i>Ammonia</i>
85	Family Nonionidae	94	Family Elphidiidae
86	Genus <i>Nonion</i>	94	Genus <i>Elphidium</i>
86	Genus <i>Nonionella</i>	104	References
86	Genus <i>Melonis</i>	116	Plates

## Introduction

The well-preserved and in most cases extremely diverse Eocene benthic foraminiferal faunas of the Hungarian Paleogene Basin (HPB) have been studied for more than a century. This report is the third in a series of studies dealing with systematic analysis, paleoecology and paleoenvironmental interpretations of Middle and Late Eocene benthic foraminifera from the HPB by the author. The first part of the series was the paleoenvironmental analysis in a Middle Eocene transgressive sequence by benthic foraminiferal fauna from Csordakút Basin, Gerecse Mountains, Hungary (OZSVÁRT, 1999). The second is on describing new benthic foraminiferal species from the HPB (OZSVÁRT, 2003a). The principal objective of this third study, which is the main part of author's PhD Thesis (OZSVÁRT, 2003b), is a comprehensive systematic analysis of benthic foraminiferal fauna providing paleoenvironmental interpretation based on multiple faunal parameters. The present paper contains brief taxonomic descriptions of 232 benthic foraminiferal species, including three new species, based on investigation of 665 samples from 12 boreholes and from two outcrops.

## Previous work

Paleogene benthic foraminifera of the HPB have been known to paleontologists since the late 19th century, when the Eocene–Oligocene transition sequences were first investigated by HANTKEN. The highly diverse and well-preserved benthic foraminiferal association from "Clavulina Szabói" layers, chiefly from Buda Mountains were published in 1868 and 1875. In these works HANTKEN described 97 and 213 species, from Kiscell Clay and from Buda Marl, respectively, including numerous new species. Further investigations in the Paleogene sequences were undertaken by VOGL (1910), who studied the micro- and macrofaunas of Piszke Marl Formation (= Buda Marl Formation) and listed 16 Upper Eocene benthic foraminifera species.

In 1956, SZÓTS published a synthesis of Eocene sediments in Hungary listing 356 species in a voluminous list without any description or illustration.

MAJZON (1966) listed 265 benthic species from the HPB, but did not include systematic analysis. However, he re-illustrated some species from HANTKEN's publications (1875).

VITÁLIS-ZILAHY (1967) studied the Upper Eocene benthic foraminiferal species from the Bükk Mountains. She published systematical descriptions and illustrations of 33 species from a borehole.

Further benthic species investigation by NYIRŐ (1970) provided systematical description and illustration of 58 species from the Bakony Mountains.

MAJZON (1972) published a systematic study of *Clavulinoides* from Hungary, including systematic descriptions and illustrations in two plates.

SZTRÁKOS (1982) described 30 foraminiferal species, including six new benthic foraminifera and illustrated 471 species from Buda Marl Formation. In 1987, SZTRÁKOS illustrated 483 species from Bartonian and Priabonian formations of Bakony Mountains and Buda Hills, but he did not offer paleoecological results.

HORVÁTH-KOLLÁNYI (1988) described 63 benthic foraminifera in detail, including one new species from shell-fill of gastropods from Dudar, Bakony Mountains collected by SZÓTS.

GELLAI-NAGY (1988) and HORVÁTH (2002, 2003) revised HANTKEN's collections; they illustrated 77 species and described 48 benthic foraminifera.

## Geological setting

Geographically, preserved sediments of the the HPB comprise a relatively narrow (less than 100 km wide) and about 300 km long SW to NE oriented belt, extending from the Zala Basin in the south-west to the Bükk Mountains in the north-east (Fig. 1). Geologically, the HPB belongs to Pelsonia Composite Terrane (KOVÁCS et al., 2000, which is part of the ALCAPA Mega-Terrane



(CSONTOS et al., 1992). The evolution of the HPB was understood as the product of transtensional (BÁLDI and BÁLDI-BEKE, 1985) or retroarc flexural tectonic processes (TARI et al., 1993) and more recently, as forearc basin migration (KÁZMÉR et al., 2003). Initiation of subsidence displays a conspicuous south-west to north-east shift from Early Lutetian to Priabonian. The Paleogene sedimentation in the HPB began in early Lutetian time with terrigenous clastic sequence interbedded with shallow marine limestone, calcareous marl (KECSKEMÉTI and VÖRÖS, 1975) and coal measures (KOPEK, 1980). The littoral and shelf sediments of the HPB has been studied intensively due to economical (coal, bauxite) and paleontological (outstanding subtropical shelf fauna) interest. These sequences pass upward into the shallow marine platform and sublittoral limestone with rich larger foraminifera fauna in the SW part of Bakony Mountains (VÖRÖS, 1989). Sedimentation grades into shallow-marine marl and calcareous marl in the NE part of Bakony Mountains and in Vértes Mountains. The shallow-marine or littoral sequences are overlain by shallow pelagic to bathyal glauconitic calcareous marl along the length of the basin, characterised by mass occurrence of coccoliths (BÁLDI and BÁLDI-BEKE, 1991).

## Localities

Four main subbasins were recognized in the HPB (Fig. 1). The *SW Bakony – Zala Paleogene Subbasin* includes a kilometre-thick calc-alkaline stratovolcanic sequence in the present-day Zala Subbasin which overlies undivided Eocene fossiliferous limestone and is interfingering with Eocene pelagic marl (KÁZMÉR et al., 2003). In this basin transgression began in Early Lutetian with thin conglomerate and paralic coal (*Darvastó Formation*). Sedimentation grades into shallow-marine carbonate rocks (*Szóc Limestone Formation*) and later into bathyal calcareous marl (*Padrag Marl Formation*). Sedimentation continued until the NP 19 (calcareous nannoplankton zone, presumably even to the NP 20 zone or later, but the Eocene sequences are topped by erosional unconformity. See Fig. 2 for detailed lithostratigraphy of this subbasin. Benthic foraminiferal assemblages and their taxonomy and paleoecology of 337 picked samples from four boreholes (Devecser 4: DV 4; Halimba 1: Hal 1; Somlóvásárhely 1: Sv 1 and Padrag 5: Pa 5) in *SW Bakony – Zala Paleogene Subbasin* were investigated (See Table 1).

Geographically, the *NE Bakony – Vértes Paleogene Subbasin* is between the *SW Bakony - Zala Paleogene Subbasin* and *Gerecse Paleogene Subbasin* (see Fig. 1). In this subbasin transgression began some five million years later than in the *NE Bakony – Vértes Paleogene Subbasin*, while the end of the sedimentation is unknown. Sedimentation started with alluvial and lacustrine sequences of coal seams (*Dorog Formation*) and continued as shallow marine marl, calcareous marl, and siltstone with rich subtropical coral, gastropod and bivalve fauna (*Csernye Formation*). This sequence passes upward into shallow marine clay marl, marl with mass occurrences of larger foraminifera (*Csolnok Clay Marl Formation*). Small, isolated, shallow-marine carbonate banks (*Szóc Limestone Formation*) interfinger with clay marl. A continuous, wide carbonate platform did not grow on the area. Shallow marine sequences are overlain by glauconitic, calcareous marl (*Padrag Marl Formation*). I have examined the benthic foraminiferal assemblages and their taxonomy and paleoecology in 250 samples from 6 boreholes (Balinka 285: Ba 285; Csetény 61: Cset 61; Dudar 240: D 240; Bakonyszentkirály 3: Bszk 3; Bakonycsernye 18: Bkcs 18 and Csákberény 89: Csbr 89) in the *NE Bakony - Vértes Paleogene Subbasin* (See Table 1).

The *Gerecse Paleogene Subbasin* situated in NW part of the HPB between the *NE Bakony – Vértes Paleogene Subbasin* in the SW and *North Hungarian Paleogene Subbasin* in the NE. This subbasin is represented by several tiny, isolated subbasins with individual subsidence and tectonic history (BERNHARDT, 1984). Generally, Paleogene sequences begin with paralic coal formations (*Dorog Formation*) in the uppermost Lutetian (NP 16 nannoplankton Zone). Similarly to the *NE Bakony – Vértes Paleogene Subbasin* numerous, small, isolated carbonate platforms built up on topographic highs (*Szóc Limestone Formation*) meanwhile the calcareous marl, siltstone with rich subtropical fauna was deposited (*Csernye Formation*) in topographic lows. Calcareous marl grade into shallow marine clay marl (*Csolnok Clay Marl Formation*).

Alternatively, fluvial sand and calcareous sandstone (*Tokod Formation*) are interbedded in clay marl. This marl and calcareous marl is overlain by calcareous marl (*Padrag Marl Formation*) on

the northern edges of *Gerecse Paleogene Subbasin*. Slumps, gravitational redeposition and mass flow deposits were described by SZTANÓ and FODOR (1997) within the uppermost part of the succession. I have examined the benthic foraminiferal assemblages and their taxonomy and paleoecology in 56 samples from 2 boreholes (Tarján 13: Tj 13 and Tarján 14: Tj 14) and from an open-cast mine of Csordakút Basin (See Table 1).

The *North Hungarian Paleogene Subbasin* includes numerous basins from the Buda Mountains to the southern edge of the Bükk Mountains. The northwestern border of subbasin is the Buda line, the southernmost part of the basin was sheared by Mid-Hungarian Line and the southward continuation of Paleogene sequences corresponds with the Paleogene succession of Slovenian Paleogene Basin (NAGYMAROSY, 1990). Sedimentation started with terrestrial breccias and conglomerates in the Buda Hills (western part of the *North Hungarian Paleogene Subbasin*) in the Priabonian stage and continued with shallow-marine carbonates (*Szép völgy Limestone Formation*). Sediments were deposited on the so-called Buda slope, the southeastward dipping flank of a complex upper Eocene anticline system, trending obliquely to the southern border fault of the escaping wedge (Fig. 10) (FODOR et al., 1992). The shallow water limestone is overlain by shallow pelagic to bathyal calcareous marl, marl which is characterised by mass occurrence of bryozoans (*Buda Marl Formation*) (ZÁGORŠEK & KÁZMÉR, 2001). I have examined the benthic foraminiferal fauna in 12 surface samples from a classical Upper Eocene section of Mátyás-hegy, Budapest (Fig. 19).

## Biostratigraphy

In this paper, the Paleogene biostratigraphy of the HPB is based on biochronological analysis of calcareous nannoplankton (BÁLDI-BEKE, 1972, 1977, 1984; NAGYMAROSY, 1983) and planktonic foraminifera (TOUMARKINE, 1971; SZTRÁKOS, 1974; HORVÁTH-KOLLÁNYI, 1983). Based on the published results of BÁLDI-BEKE and HORVÁTH-KOLLÁNYI the depositional time of Eocene sequences corresponds to NP14–21 nannoplankton zones. Fig. 2 summarises the stratigraphy of the HPB, with data from BÁLDI-BEKE (nannoplankton; 1984, HORVÁTH-KOLLÁNYI (planktonic foraminifera; 1983) and BERNHARDT et al. (lithostratigraphy; 1988).




Fig. 1. Position of the Hungarian Paleogene Basin in the Eastern Alpine-Western Carpathian junction. Base map: OZSVÁRT *in* KÁZMÉR et al. (2003).




Fig. 2. Eocene lithostratigraphy (after BÁLDI-BEKE, 1984; NAGYMAROSY and BÁLDI-BEKE, 1988; BERNHARDT et al., 1988 and CSÁSZÁR, G., 1997) and chronostratigraphy (after HARDENBOL et al., 1998) in the Hungarian Paleogene Basin.

## Methods

Benthic foraminifera from 12 boreholes and from two outcrops from the HPB were examined. All samples from boreholes were obtained from the Geological Institute of Hungary. Each sample from outcrops was collected by the author and approximately 250–300 g from each sample was cleaned using 25% hydrogen-peroxide. The residues were washed over a 50 µm sieve and dried. Samples were divided into the grain-size fractions of 50–500 µm and >500 µm. Faunal analysis was performed only on the 50–500 µm fraction. Prior to separation and counting, the samples were divided into parts of 1/2, 1/4, 1/8 etc., using a micro-splitter. For faunal analysis, ~200–250 specimens were picked, unless their numbers were less than 200, in which case all specimens were picked. For paleoecological and paleoceanographical studies, the Benthic Foraminiferal Number (BFN), Diversity  $H(S)$ , Dominance, Fisher alpha ( $\alpha$ ) index and faunal composition were calculated. The detailed description of calculation for  $H(S)$  is given in BUZAS and GIBSON (1969), for Dominance, Fisher index and faunal composition is given in MURRAY (1991). The BFN parameter is the number of specimens per gram sediments (SCHOTT, 1935) and gives an indication of the sedimentation rate, productivity and carbonate dissolution (DOUGLAS and WOODRUFF, 1981).

Foraminifera specimens were identified under a reflected light binocular microscope. All species were identified, described and photographed with a scanning electron microscope in Eötvös University of Budapest (ELTE).

## Sections

### *SW Bakony – Zala Paleogene Subbasin*

*Devecser 4 borehole*

The DV 4 borehole is located in the middle part of *SW Bakony – Zala Paleogene Subbasin* (Fig. 3). This borehole penetrated to 910 m depth, in which a 267 m thick Eocene succession was recovered. A 49.1 m continuous core from the top of the Eocene succession was investigated. Eighty-three samples yielded a total of 5136 specimens of 132 benthic foraminiferal species (see Table 2). The studied sequence is composed of clay, clay-marl and marl units. Diversity shows strong fluctuations (Fig. 4) in lower part ( $H(S) = 1-2$ ; Fisher ( $\alpha$ ) = 5-15) and continued with diversity increasing in middle and upper part with moderate fluctuation ( $H(S) = 2.5-3.5$  Fisher ( $\alpha$ ) = 10-30). More than 50% of benthic foraminifera test are agglutinated in the lower part of the section (between 95 m and 90 m). Upward this value varies between 0 and 10 % in the upper part of the section and hyaline tests dominate. The benthic foraminiferal fauna is distributed differently through the succession. Dominant species are: *Dentalina elegans*, *Gyroidionides soldanii*, *Lenticulina arcuatostrata*, *Nodosaria longiscata*, *Stilostomella elegans* (see Table 2).




Fig. 3. Position and hypothetical configuration of the *SW Bakony – Zala Paleogene Subbasin* in the Hungarian Paleogene Basin.




Fig. 4. Shannon-Wiener, Dominance, Fisher index and faunal composition of the Devceser 4 borehole.

#### Halimba 1 borehole

Hal 1 borehole penetrated 161.5 m thick Eocene succession, located on the south-eastern part of SW Bakony – Zla Paleogene Subbasin (Fig. 3). Sedimentation started with terrestrial breccia and conglomerate. Upwards this succession changes into shallow marine carbonates. These carbonates are overlain by the shallow pelagic to bathyal calcareous marl and marl. Thirty-five samples yielded 2813 specimens of 60 benthic foraminiferal species (see Table 2). The Shannon-Wiener H(S) diversity index of the benthic foraminiferal association increases continuously from the lower part to top of the succession (Fig. 5). Diversity varies between 0.67 and 3.15. Fisher ( $\alpha$ ) varies from 0.5 to 10. All foraminifera tests are hyaline along the whole section, except in the upper part (between 44 m and 60 m, where the ratio of agglutinated foraminifera increases (5–20%). Dominant benthic foraminifera are *Eponides polygonus*, *Heterolepa dutemplei*, *Cibicides* spp., *Dentalina* sp. and *Lenticulina* sp. (see Table 2).




Fig. 5. Shannon-Wiener, Dominance, Fisher index and faunal composition of the Halimba 1 borehole.

#### *Padrag 5 borehole*

The Pa 5 borehole is located in the central part of the subbasin (Fig. 3). A 58.2 m continuous core of the 244 m thick Eocene succession was investigated (see Fig. 6). 13 samples contained enough specimens for statistical analysis. 1705 specimens of 55 species were analysed (see Table 2). Diversity is low (Fig. 6), H(S) varies continuously (between 2.5 and 3; Fisher ( $\alpha$ ) = 5–10). At 34 m, the diversity is 0 and the dominant species is *Heterolepa simplex*. Calcareous, hyaline tests dominate the section. Benthic foraminifera assemblage is dominated by *Uvigerina eocaena*, *Lenticulina arcuatostrata* and *Heterolepa simplex* (see Table 2).




Fig. 6 – Shannon-Wiener, Dominance, Fisher-index and faunal composition of Padrag 5 borehole.

### Somlóvásárhely 1 borehole

Sv 1 borehole is located in the north-western part of *SW Bakony – Zala Paleogene Subbasin* (Fig. 3). A total of 306.8 m thick Eocene succession was drilled in this core, represented by continuous sequences between NP14 and NP 19 nannoplankton zones. The oldest Eocene sediments are terrestrial clay, variegated clay and clay marl. The sequence grades into shallow-marine carbonate (Szóc Limestone Formation). At 720 m the limestone grades into grey, greenish grey marl of the Padrag Formation. 223 samples contain 10,234 specimens of 64 species. The Shannon-Wiener H(S) diversity index of the benthic foraminiferal association shows strong fluctuation along the whole section (Fig. 7). Index values vary between 0 and 2 in the lower part of the section (between 837.8 m and 712 m). Diversity increases slightly ( $H(S) = 2.5-3$ ; Fisher ( $\alpha$ ) = 2–5) in the middle part of the section (between 580 m and 712 m, upper interval decreases suddenly ( $H(S) = 0.5-2$ , while Fisher ( $\alpha$ ) index increases at the top of the section. Calcareous tests are dominant (60–100%) in percentage, particularly the porcellaneous (60–100 % between 837.6 m and 828.0 m) and agglutinated tests (between 700.0 m to 800.0 m) are significant. The assemblage is chiefly composed of *Cibicides* sp., *Uvigerina eocaena*, *Dentalina elegans*, *Lenticulina arcuatostrata* and *Heterolepa dutemplei* (see Table 2).






Fig. 7. Shannon-Wiener, Dominance, Fisher index and faunal composition of Somlóvásárhely 1 borehole.

### *NE Bakony–Vértes Paleogene Subbasin*

#### *Bakonycsernye 18 borehole*

Bkcs 18 borehole was drilled in south-eastern part of *NE Bakony–Vértes Paleogene Subbasin* (Fig. 8). This core penetrated 93 m continuous Middle Eocene grey, greenish grey marl of Padrag Formation. Twenty-one samples yielded 2332 specimens of 81 species. Diversity index of the association increases slightly ( $H(S) = 1.5-3$ ; Fisher ( $\alpha$ ) = 1–10) up to 352 m from the bottom (Fig. 9). Diversity is low ( $H(S) = 0.5-1.5$ ; Fisher ( $\alpha$ )=2–3) in the middle part of the section and increases ( $H(S) = 1.5-3$ ; Fisher ( $\alpha$ ) = 4–6) in the upper interval. Calcareous tests are dominant, particularly agglutinated tests are significant in the upper interval. The dominant benthic foraminifera are *Lenticulina arcuostriata*, *Heterolepa dutemplei*, *Bulimina truncana* and *Heterolepa simplex* (see Table 2).




Fig. 8. Position and hypothetical configuration of the NE Bakony–Vértés Paleogene Subbasin in the Hungarian Paleogene Basin




Fig. 9. Shannon-Wiener, Dominance, Fisher index and faunal composition of Bakonycsernye 18 borehole.

### Bakonyszentkirály 3 borehole

Bszk 3 borehole contains 180 m thick Middle and Upper Eocene sediments of grey marl of *Padrag Formation*. 35 samples yielded 7324 specimens of 43 benthic foraminiferal species. Diversity is low (Fig. 10), ( $H(S)$  varies between 1 and 3; Fisher ( $\alpha$ ) = 2–14) and shows a decrease in the upper interval of the section. Calcareous tests are significant (40–100%, however the porcellaneous and agglutinated tests are also present in whole section (up to 60%). Dominant benthic foraminifera are *Lenticulina arcuatostrata*, *Heterolepa dutemplei*, *Bulimina truncana* and *Quinqueloculina* sp. (see Table 2).




Fig. 10. Shannon-Wiener, Dominance, Fisher index and faunal composition of Bakonyszentkirály 3 borehole.

### Balinka 285 borehole

Ba 285 borehole is located in the south-eastern part of the *NE Bakony-Vértes Paleogene Subbasin* (Fig. 8). This core penetrated a 100 m thick Eocene succession and 31 samples contain 3462 specimens of 56 benthic foraminiferal species.  $H(S)$  diversity index of the benthic foraminiferal association increases slightly ( $H(S) = 2.5-3$ ; Fisher ( $\alpha$ ) = 2–16) from the bottom to top of the investigated section (Fig. 11). In the lower part of the section (between 564.0 m to 580.0 m) porcellaneous tests dominate (60–80%), while agglutinated tests are present in 0–20%. Calcareous foraminifera tests become dominant from 564.0 m to top of the section. The benthic foraminiferal fauna are distributed differently through the succession, the dominant species are: *Heterolepa*

*dutemplei*, *Bulimina truncana*, *Quinqueloculina carinata*, *Cibicidoides* sp., and *Nodosaria* sp. (see Table 2).




Fig. 11. Shannon-Wiener, Dominance, Fisher index and faunal composition of Balinka 285 borehole.

### Csetény 61 borehole

Cset 61 borehole penetrated 543 m, in which a 245 m thick Eocene succession was recovered. The oldest Eocene sediments (Fig. 12) at depth of 515.4 m are Upper Lutetian (NP16) gray clay, variegated clay and clay marl layers, frequently intercalated by thin coal seams (*Dorog Formation*). Above 476.0 m this basal terrestrial sequence interfingers with calcareous marls and lumachelle of the Csernye Formation. At 470.0 m calcareous marls grade into grey, greenish grey marls of the Padrag Formation which is characterised by mass occurrence of coccoliths. Between 425.6 m and 420.6 m thin-bedded, laminated, fine-grained marls have been observed. Higher up (between 420.0 m and 401.0 m, stratification becomes more pronounced and carbonate content increases. The upper part of the section is lithologically uniform, but the carbonate content of marl layers is variable. The interval between 402.5 m and 356.0 m consists of thick-bedded marl, frequently intercalated with tuffaceous layers. Seventy samples yielded 3404 specimens of 102 benthic foraminiferal species. The Shannon-Wiener H(S) diversity index of the benthic foraminiferal association varies between 1 and 3 (Fig. 7). In the lower part of the section (between 432 m and 470 m) diversity exhibits higher values (between 2 and 3). In the middle part of the section (around 432 m) a drop in diversity is observed, and H(S) values show strong fluctuations between 1 and 2. Agglutinated and porcellaneous tests dominate in bottom part, but decrease their dominance progressively upwards. Calcareous tests are dominated from the 468.0 m. The dominant foraminiferal assemblage is composed of *Lenticulina platyptera*,

*Cibicidoides eoceanus*, *Lenticulina arcuatostrata*, *Eponides polygonus* and *Bulimina truncana* (see Table 2).




Fig. 12. Shannon-Wiener, Dominance, Fisher index and faunal composition of Csetény 61 borehole.

#### Dudar 240 borehole

D 240 borehole yield 168.5 m thick Middle and Upper Eocene (NP16–NP19 nannoplankton zones) sequences. The bottom interval is terrestrial clay, variegated clay and clay marl. The sequence grades into the shallow marine, marly limestone and limestone (*Szóc Limestone Formation*). This carbonate succession passes into grey, grey marls upward. Thirty-seven samples yielded 2773 specimens of 64 benthic foraminiferal species. H(S) index of the benthic foraminiferal association varies between 1.5 and 2.5, while the Fisher ( $\alpha$ ) varies between 2 and 8 (Fig. 13). The diversity indexes show strong fluctuation along the section. Calcareous tests are dominant in all samples. The dominant benthic foraminifera are *Lenticulina arcuatostrata*, *Heterolepa dutemplei*, *Bulimina truncana*, *Pararotalia inermis* and *Stilostomella* sp. (see Table 2).




Fig. 13. Shannon-Wiener, Dominance, Fisher index and faunal composition of Dudar 240 borehole.

### Csákberény 89 borehole

Csbr 89 borehole penetrated a 197.3 thick Lutetian (NP16 nannoplankton zone) succession. This consists of grey clay, clay marl, calcareous marl and marl. Seventy-six samples contain 4767 specimens of 98 benthic foraminiferal species. Diversity indexes show strong fluctuation in whole section ( $H(S) = 1-2.5$ ; Fisher ( $\alpha$ ) = 1-6). They show slightly decreasing (Fig. 14) from bottom to the middle interval, later increase up to relatively high values ( $H(S) = 1.5-3$ ; Fisher ( $\alpha$ ) = 6-10). Porcellaneous tests dominant (80-100%) up to 240.0 m, from that level calcareous tests become dominant. The benthic foraminiferal fauna is distributed differently through the succession. The dominant assemblage is composed of *Quinqueloculina carinata*, *Eponides haidingeri*, *Eponides umbonatus* and *Triloculina porvaensis* (see Table 2).




Fig. 14. Shannon-Wiener, Dominance, Fisher index and faunal composition of Csákberény 89 borehole.

### *Gerecse Paleogene Subbasin*

#### *Csordakút outcrop*

Csordakút is located in the southern part of the Gerecse Mountains (Fig. 15). The 35 m thick Middle Eocene sequence of the open-cast mine of Csordakút Basin is the following: very fine-grained, grey, reddish brown variegated clay covered by brown marl with Triassic dolomite boulders in the lower part of the bed. This sequence grades into grey, brownish-grey, well-stratified *Nummulites* marl and limestone. It is biomicrite with packstone (rudstone), wackestone (floatstone) and wackestone/packstone texture. This marl and limestone passes into light-coloured, unstratified, nodular “*Alveolina*” limestone. It is biomicrite with mudstone, wackestone and packstone texture. This formation is overlain by *Ostrea lumachelle*. It is poorly consolidated greenish grey argillaceous marl with rich mollusca fauna. Twenty-one samples yielded 5183 specimens of 41 benthic foraminiferal species. Diversity indexes show slightly increasing (Fig. 16) from bottom to upper interval of the section ( $H(S) = 0.5\text{--}2$ ; Fisher ( $\alpha$ ) = 1–4). Calcareous tests are dominant, particularly the porcellaneous and agglutinated tests are also significant in particular levels. The dominant benthic foraminifera are *Nonion scaphum*, *Pararotalia curry*, *Pyrgo simplex* and *Eponides polyganus* (see Table 2).




Fig. 15. Position and hypothetical configuration of *Gerecse Paleogene Subbasin* in the Hungarian Paleogene Basin.




Fig. 16. Shannon-Wiener, Dominance, Fisher index and faunal composition of Csordakút section.



Tarján 13 and Tarján 14 boreholes

Tj 13 and Tj 14 boreholes are located in the middle part of the *Gerecse Paleogene Subbasin*, close to each other (Table 1). Both boreholes represent the Middle Eocene (NP16–NP17 nannoplankton zones) sequences, and the investigated samples contain 2630 specimens of 43 individual benthic foraminiferal species (see Table 2). Diversity indices are low ( $H(S) = 1.5\text{--}2$ ; Fisher ( $\alpha$ ) = 1–5) and show decreasing trend in Tj 13 borehole (Fig. 17), and relatively high fluctuation in Tj 14 borehole (Fig. 18). Calcareous and porcellaneous tests are dominant in both sections. The benthic foraminiferal fauna is distributed differently through the succession. Dominant assemblages are composed of *Quinqueloculina carinata*, *Heterolepa dutemplei*, *Cibicides pygmeus*, *Cibicides lobatulus* and *Bulimina parisensis* (see Table 2).




Fig. 17. Shannon-Wiener (H(S)), Dominance, Fisher-index and faunal composition of the Tarján 13 borehole




Fig. 18. Shannon-Wiener (H(S)), Dominance, Fisher index and faunal composition of Tarján 14 borehole.

### North Hungarian Paleogene Subbasin

#### Mátyás-hegy outcrop

The section is situated in Budapest, District 3, at the fork of the Virág Benedek and Mátyáshegyi streets opposite the entrance of Pálvölgy Cave (Fig. 19). The quarry is part of the Buda Natural Protection Area. A nearly complete continuous Priabonian sequence is exposed in this quarry: paleoenvironments range from transgressive conglomerate (*Kosd Formation*) through neritic limestone (*Szép völgy Limestone*) and bryozoan marl to shallow bathyal globigerina marl (*Buda Marl Formation*). The sequence is subdivided into three units: hard compact limestone (0–10 m) containing large quantity of orthofragminids, in the lower 5 m in rock-forming quantity. This formation is overlain by marly limestone. This is covered by yellow, light-brown *Buda Marl Formation*. Twelve samples contain 1675 specimens of 100 benthic foraminiferal species. The Shannon-Wiener H(S) diversity index of the benthic foraminiferal association increases slightly (Fig. 20) from the bottom upwards (H(S) = 1–3.5; Fisher ( $\alpha$ ) = 1–22). Calcareous tests dominate, while agglutinated test are also significant in the upper interval. The dominant benthic foraminifera are *Cibicides oligocenicus*, *Heterolepa dutemplei*, and *Clavulinoides szaboi* (see Table 2).




Fig. 19. Position and hypothetical configuration of the *North Hungarian Paleogene Subbasin* in the Hungarian Paleogene Basin.




Fig. 20. H(S), Dominance, Fisher index and faunal composition of Mátyás-hegy section.

## Paleoecology

### *Benthic foraminiferal assemblages*

In total, 232 benthic foraminiferal species were picked from 665 samples. The assemblages are moderately diverse,  $H(S)$  varies between 1 and 3.5, principally ( $H(S_{\min}) = 0.17$  at 799.0 m in Sv 1 borehole and  $H(S_{\max}) = 3.66$  at top of Mátyás-hegy section), with 40 to 130 species, and 10 to 30 genera present in most sections. Commonly, the  $H(S)$  index varies between 1.5 and 2 and these values show strong fluctuations in all sections. Generally, foraminifera assemblages in different sections contain a few dominant species, except in such samples where extreme paleoenvironmental conditions might be assumed (typical in brackish or hypersaline conditions). The Fisher ( $\alpha$ ) index varies between 1 and 4, principally ( $F(\alpha_{\max}) = 64.34$  at 396.0 m in Cset 61 borehole and  $F(\alpha_{\min}) = 0.18$  at 294.45 m in Csbr 89 borehole). The most frequent Fisher ( $\alpha$ ) index values ( $<5$ ) during NP16 nannoplankton zone are typical for brackish or hypersaline, marginal marine environments. The Fisher ( $\alpha$ ) index varies between 5 and 15 from NP17 nannoplankton zone to NP19 nannoplankton zone which are typical for the inner and outer shelves and for bathyal (values between 1 and 23 (see Murray, 1991) region. Composition of benthic foraminiferal assemblages confirms shallow water conditions during the NP16 nannoplankton zone and the normal marine environment in the younger parts of the sections. Benthic foraminiferal faunas are chiefly dominated by hyaline, calcareous species, making up about between 60 to 100 % of the total fauna. The most common calcareous taxa include *Cibicides* spp., *Heterolepa* spp., *Lenticulina* spp., *Dentalina* spp., *Uvigerina* spp. The most common porcellaneous taxa are *Quinqueloculina* spp. and the most common agglutinated taxa are *Clavulinoides* spp., and *Textularia* spp. All these common taxa show strong fluctuations in relative abundance.

### *Water temperature and salinity*

Diversity and composition of benthic foraminiferal assemblages is strongly controlled by water temperature and salinity of water mass (Sen Gupta, 1999). Especially, in shallow and deeper region on shelves, where changes in temperature and salinity might pass off rapidly. Several observations of recent benthic foraminiferal faunas illustrate the temperature and salinity oscillations in shallow water region connect to seasonal changes (i. e. GOODAY, 1988, 1993; KITAZATO et al., 2000, while in abyssal region this procedure could be a long-term change. The dominant *Miliolina* spp. and *Cibicides* spp. and the low-diversity ( $H(S) = 0.5-1.5$ ; Fisher ( $\alpha$ ) = 2-4) indicate warm-temperate (18-23°) temperature and oscillated salinity in the SW Bakony – Zala Paleogene Subbasin up to NP16 nannoplankton zone. Significant temperature decreasing was recognized during early NP16 nannoplankton zone, where temperate and cold water foraminiferal species (*Lenticulina* spp., *Heterolepa* spp.) (after Murray, 1991) became dominant up to end of the NP18 nannoplankton zone. In NE Bakony–Vértes Paleogene Subbasin were warm water foraminiferal species dominated during the NP16 nannoplankton zone. Moreover presumably strongly oscillating salinity in warm sea-water. A significant cooling event was recognized by appearance of cold water foraminiferal species during NP17–NP18 nannoplankton zones. In Gerecse Paleogene Subbasin presumably warm and strong salinity fluctuation as indicated by low diversity ( $HS) = 0.5-1.5$ ; Fisher ( $\alpha$ ) = 2-3) during NP16–NP17 nannoplankton zones. Like in SW and NE Bakony Paleogene Subbasins there was a similar trend represented in North Hungarian Paleogene Subbasin during NP18–NP19 nannoplankton zones: warm-temperate water temperature without significant environmental stress; later benthic foraminiferal assemblages indicate temperate and cold bottom water.

## Acknowledgements

This study is part of a joint paleoecological and paleoceanographical analysis of the HPB during the Eocene time that was my doctoral project at Eötvös University, Budapest. My special thanks go to my supervisor, Miklós KÁZMÉR, for his enduring support, advice and encouragement throughout the project. I thank Katalin KOLLÁNYI (Geological Institute of Hungary) for providing access to their benthic foraminiferal collections. Special thanks are due to Mária HORVÁTH (Eötvös University) for reviewing an earlier version of my PhD manuscript, which included this systematic part. The draft manuscript has benefited from the constructive criticism and English correction of József PÁLFY (Hungarian Natural History Museum). The printing of this manuscript has been supported by a publishing grant from the Hungarian Academy of Sciences.

## Systematic paleontology

Benthic foraminifera taxonomy follows chiefly the systematic classification of LOEBLICH and TAPPAN (1988). Geographical and geological (age ranges) distribution is not given in taxonomy descriptions, instead the occurrences of taxa in the HPB are shown on Table 2. Holotypes and all illustrated material are deposited in the Department of Geology and Paleontology of Hungarian Natural History Museum, Budapest.

Phylum Protozoa GOLDFUSS, 1818  
Class Rhizopodea VON SIEBOLD, 1845  
Order Foraminifera EHRENBERG, 1930  
Suborder Textulariina DELAGE and HÉROUARD, 1896  
Superfamily Astrorhizacea BRADY, 1881  
Family Astrorhizidae BRADY, 1881  
Subfamily Astrorhizinae BRADY, 1881

Genus *Astrorhiza* SANDAHL, 1858

Type species: *Astrorhiza limicola* SANDAHL, 1858

*Astrorhiza bakonycseryensis* n. sp.  
Plate 1, Figure 1

Derivatio nominis: In reference to its occurrence in a borehole near Bakonycserye, Hungary.

Holotype: Figured specimen on Plate 1, Figure 1. (Inventory number: M 2008.116.1)

Dimension of holotype: diameter 0.89 mm.

Description: Test unilocular, spherical or oval in outline, very short radial arms visible on a single chamber; small oval or round apertures on arms without any sign of tooth; wall simple, imperforate, finely to coarsely agglutinated grains.

Type locality: Bakonycserye 18 (Bkcs 18) borehole.

Type strata: In Bkcs 18 borehole, at 365.0 m. Padrag Marl Formation (Middle Eocene, Bartonian).

Remarks: Generally, the specimens which belong to genus *Astrorhiza* are always compressed in various planes.

Family Bathysiphonidae AVNIMELECH, 1952

Genus *Bathysiphon* SARS, 1872

Type species: *Bathysiphon filiformis* Sars, 1872

*Bathysiphon eocenicus* CUSHMAN and HANNA, 1927  
Plate 1, Figures 2–3

- 1927 *Bathysiphon eocenicus* n. sp. – CUSHMAN and HANNA, p. 210, pl. 13, figs. 2–3.  
1936 *Bathysiphon eocenicus* CUSHMAN and HANNA – CUSHMAN and MCMASTERS, p. 508, pl. 74, fig. 1.  
1942 *Bathysiphon eocenicus* CUSHMAN and HANNA – CUSHMAN and SIEGFUS, p. 400, pl. 15, fig. 1.  
1944 *Bathysiphon eocenicus* CUSHMAN and HANNA – CUSHMAN and SIMONSON, p. 193, pl. 30, fig. 1.  
1947 *Bathysiphon eocenicus* CUSHMAN and HANNA – CUSHMAN et al., p. 97, pl. 12, figs. 1–2.  
1947 *Bathysiphon eocenicus* CUSHMAN and HANNA – CUSHMAN and STONE, p. 2, pl. 1, fig. 1.  
1949 *Bathysiphon eocenicus* CUSHMAN and HANNA – BERMÚDEZ, p. 47, pl. 1, figs. 1–2.  
1949 *Bathysiphon eocenicus* CUSHMAN and HANNA – CUSHMAN and STONE, p. 75, pl. 13, fig. 3.  
1951 *Bathysiphon eocenicus* CUSHMAN and HANNA – CUSHMAN and STAINFORTH, p. 142, pl. 25, fig. 4.  
1952 *Bathysiphon eocenicus* CUSHMAN and HANNA – TODD and KNIKER, p. 4, pl. 1, figs. 3–4.

- 1957 *Bathysiphon eocenica* CUSHMAN and HANNA – SMITH, p. 148, pl. 17, fig. 1.  
 1972 *Bathysiphon eocenica* CUSHMAN and HANNA – MCDUGALL, p. 33, pl. 1, fig. 1.  
 1975 *Bathysiphon eocenicus* CUSHMAN and HANNA – BRAGA et al., p. 102, fig. 5.  
 1982 *Bathysiphon eocenicus* CUSHMAN and HANNA – SZTRÁKOS, pl. 1, fig. 2.  
 1985 *Bathysiphon eocenicus* CUSHMAN and HANNA – GRÜNIG, p. 253, pl. 1, figs. 4–6.  
 1987 *Bathysiphon* sp. – SZTRÁKOS, pl. 1, figs. 2–3.  
 2005 *Bathysiphon eocenicus* CUSHMAN and HANNA – ANAN, p. 18, pl. 1, fig. 1.  
 2005 *Bathysiphon eocenica* CUSHMAN and HANNA – NARAYAN et al., p. 118, pl. 1, fig. 5.

Description: Test tubular, elongate, cylindrical, strongly compressed in apertural view; wall simple, finely agglutinated with amorphous material; oval aperture without any sign of tooth; length unidentified, breadth 0.4–0.6 mm.

*Bathysiphon saidi* (ANAN, 1994)

Plate 1, Figure 4

- 1927 *Rhabdammina eocenica* n. sp. – CUSHMAN and HANNA, p. 208, pl. 13, fig. 1.  
 1952 *Rhabdammina eocenica* CUSHMAN and HANNA – TODD and KNIKER, p. 4, pl. 1, figs. 1–2.  
 1982 *Rhabdammina eocenica* CUSHMAN and HANNA – SZTRÁKOS, pl. 1, fig. 1.  
 1994 *Rhabdammina saidi* n. sp. – ANAN, p. 218, fig. 8. 1.  
 1997 *Bathysiphon abbassi* n. sp. – HUSSEIN, p. 109, fig. 3. 2.  
 2005 *Bathysiphon saidi* (ANAN) – ANAN, p. 19, pl. 1, fig. 2.

Description: Test tubular, straight and cylindrical, mildly compressed in apertural view; wall simple, coarsely agglutinated; oval aperture without any sign of tooth; length unidentified, breadth 0.3 – 0.5 mm.

Remarks: CUSHMAN and HANNA (1927) described the new, straight and elongate species *Rhabdammina eocenica* from Coalinga, California, although shape of the genus *Rhabdammina* was defined as triradiate or quadriradiate. They described the *Bathysiphon eocenica* as an other new species from the same section. ANAN (2005) established the new species *Rhabdammina saidi*, which was placed in 2005 in the genus *Bathysiphon*. The species *Bathysiphon saidi* has straight, elongate test and its wall agglutinated by coarse fragmented grains as CUSHMAN and HANNA (1927) described for *Rhabdammina eocenica*. Thus, the genus name *Rhabdammina* is invalid for the elongate and straight test so it should be placed into the genus *Bathysiphon*. Unfortunately, the species name *eocenica* is also invalid, because the species *Bathysiphon eocenica* is a valid species name so having referred to International Code of Zoological Nomenclature (Article 57.3) the species *Rhabdammina eocenica* and the species *Bathysiphon eocenica* become secondary homonyms. Thus, the first valid synonym (*Bathysiphon saidi* in 2005) was used for this species. *Bathysiphon saidi* differs only from *B. eocenicus* in having rough agglutinated surface with coarse fragmented grains.

Family Rhabdamminidae BRADY, 1884

Subfamily Rhabdammininae BRADY, 1884

Genus *Rhabdammina* SARS, 1869

Type species: *Rhabdammina abyssorum* SARS, 1869

*Rhabdammina abyssorum* SARS, 1868

Plate 1, Figure 5

- 1868 *Rhabdammina abyssorum* n. sp. – SARS, p. 248.  
 1869 *Rhabdammina abyssorum* SARS – SARS in CARPENTER, p. 60.  
 1881 *Rhabdammina abyssorum* SARS – CARPENTER, p. 562, pl. 321c–d.  
 1884 *Rhabdammina abyssorum* SARS – BRADY, p. 266, pl. 21, figs. 1–3; 10–13.  
 1889 *Rhabdammina abyssorum* SARS – NEUMAYR, p. 173, pl. 17, fig. a.  
 1893 *Rhabdammina abyssorum* SARS – EGGER, p. 255, pl. 4, fig. 31.  
 1894 *Rhabdammina abyssorum* SARS – GOES, p. 19, pl. 4, figs. 67–68.  
 1896 *Rhabdammina abyssorum* SARS – GRZYBOWSKI, p. 275, pl. 8, figs. 1–4.  
 1901 *Rhabdammina abyssorum* SARS – SCHUBERT, p. 17, pl. 1, figs. 5–9.

- 1902 *Rhabdammina abyssorum* SARS – CHAPMAN, p. 125, pl. 6, fig. H.  
 1918 *Rhabdammina abyssorum* SARS – CUSHMAN, p. 15, pl. 6, fig. 1; pl. 7, fig. 1.  
 1921 *Rhabdammina abyssorum* SARS – CUSHMAN, p. 36, pl. 1, fig. 2.  
 1925 *Rhabdammina abyssorum* SARS – CUSHMAN, pl. 1, fig. 2.  
 1927 *Rhabdammina abyssorum* SARS – CUSHMAN, p. 7, pl. 1, fig. 4.  
 1928 *Rhabdammina abyssorum* SARS – CUSHMAN, p. 64, pl. 2, figs. 7–8.  
 1930 *Rhabdammina abyssorum* SARS – HOFKER, p. 107, pl. 42, figs. 5–11; pl. 43, figs. 3–4, 6.  
 1933 *Rhabdammina abyssorum* SARS – CUSHMAN, p. 69, pl. 1, fig. 9.  
 1933 *Rhabdammina abyssorum* SARS – GALLOWAY, p. 69, pl. 5, fig. 5.  
 1969 *Rhabdammina abyssorum* SARS – VILKS, p. 43, pl. 1, fig. 2.  
 1979 *Rhabdammina eocenica* CUSHMAN and HANNA – SZTRÁKOS, pl. 1, figs. 1–2.  
 1985 *Rhabdammina eocenica* CUSHMAN and HANNA – GRÜNIG, p. 253, pl. 1, figs. 1–2.

Description: Test triradiate, mildly compressed in apertural view; aperture terminal, mildly compressed; wall simple, coarsely agglutinated.

Superfamily Hormosinacea HAECKEL, 1894  
 Family Hormosinidae HAECKEL, 1894  
 Subfamily Reophacinae CUSHMAN, 1910

Genus *Reophax* De Montfort emend. Brönnimann and Whittaker, 1980

Type species: *Reophax arctica* BRADY, 1881

*Reophax harrisi* nomen novum  
 Plate 1, Figure 6

1951 *Reophax sabulosus* n. sp. – HARRIS and JOBE, p. 5, pl. 1, fig. 3.

Description: Test elongate, monoserial, four chambers, circular in apertural view; chambers increasing gradually in size, earlier chambers subspherical, final chamber inflated; sutures barely visible, mildly depressed; terminal aperture, circular, slightly produced; wall simple, coarsely agglutinated; length of test 1.53 mm, maximum breadth 0.5 mm.

Remarks: BRADY established the new species *Reophax sabulosa* in 1882 and HARRIS and JOBE (1951) described the new species *Reophax sabulosus* from the Paleocene Midway Formation from Hempstead County, Arkansas, USA. Thus, the *Reophax sabulosus* is invalid name, because it is junior homonym of *Reophax sabulosa* referred to International Code of Zoological Nomenclature (Article 60.1).

Etymology: Named after R. W. HARRIS who described the new species *Reophax sabulosus*.

Holotype: Original holotype deposited in the collections of Oklahoma, Norman, Oklahoma; no. PeM 10,002A.

Type locality: Exposure in a roadside ditch on the east side of State Highway 29 in Hempstead County, Arkansas, USA.

Type strata: Midway Formation, Paleocene.

Superfamily Lituolacea DE BLAINVILLE, 1827  
 Family Haplophragmoididae MAINC, 1952

Genus *Haplophragmoides* CUSHMAN emend. HÖGLUND, 1947

Type species: *Haplophragmoides canariense* D'ORBIGNY, 1889

*Haplophragmoides* sp.  
 Plate 1, Figure 8



Description: Test lenticular, planispiral, involute, circular and slightly lobulate in outline, biconvex in cross-section; chambers increasing in size; sutures barely visible; wall thick, finely agglutinated; aperture deeply impressed fissure on interiomarginal side.

Family Lituolidae DE BLAINVILLE, 1827

Genus *Ammomarginulina* WIESNER, 1931

Type species: *Ammomarginulina enis* WIESNER, 1931.

*Ammomarginulina* sp.  
Plate 1, Figure 7

Description: Test trochospiral in earlier three – four chambers, becoming monoserial in final chambers; length approximately two and one – half times width breadth; test elongate, mildly compressed in cross-section; trochospiral chambers increasing rapidly in size, monoserial chambers equal in size; sutures barely visible; aperture terminal, compressed from apertural view; wall simple, coarsely agglutinated.

Superfamily Spiroplectamminacea CUSHMAN, 1927

Family Spiroplectamminidae CUSHMAN, 1927

Subfamily Spiroplectammininae CUSHMAN, 1927

Genus *Bolivinopsis* JAKOVLEV emend. KISELMAN, 1964

Type species: *Bolivinopsis capitata* JAKOVLEV, 1891

*Bolivinopsis foliacea* (GRZYBOWSKI, 1898)  
Plate 1, Figure 9

1898 *Spiroplecta foliacea* n. sp. – GRZYBOWSKI, p. 294, pl. 12, figs. 14–15.

1952 *Spiroplectamina elgansoensis* n. sp. – TODD and KNIKER, p. 7, pl. 1, fig. 17.

1975 *Bolivinopsis foliacea* (GRZYBOWSKI) – PROTO DECIMA and DE BIASE, p. 91, pl. 3, fig. 10.

Description: Test strongly elongate, strongly compressed in cross-section; test planispiral in early four – five chambers, becoming biserial in adult portion; length approximately six – seven times width; initial chambers slowly increasing in size as added; sutures fairly distinct, slightly depressed, straight; wall finely agglutinated; aperture a narrow slit between the last two chambers.

Genus *Spiroplectamina* CUSHMAN emend. NØRVANG, 1966

Type species: *Spiroplectamina biformis* (PARKER and JONES, 1865)

*Spiroplectamina carinata* (D'ORBIGNY, 1846)  
Plate 1, Figures 10–11

1846 *Textularia carinata* n. sp. – D'ORBIGNY, p. 247, pl. 14, figs. 32–34.

1936 *Spiroplectamina carinata* n. sp. – SUBBOTINA, p. 6, pl. 1, figs. 8–11.

1951 *Spiroplectamina carinata* (D'ORBIGNY) – MARKS, p. 35, pl. 6, figs. 2 a–b.

1961 *Spiroplectamina carinata* (D'ORBIGNY) – KAASSCHIETER, p. 140, pl. 1, fig. 12.

1969 *Textularia carinata* (D'ORBIGNY) – RÖGL, p. 67, pl. 1, figs. 2a–b.

1985 *Spiroplectamina carinata* (D'ORBIGNY) – KORECZNE-LAKY and NAGYNÉ-GELLAI, pl. 6, fig. 11; pl. 84, fig. 3; pl. 104, figs. 4–5, 7, 15.

Description: Test planispiral in earlier three – four chambers, becoming biserial in adult part; seven to eight pairs of chambers in biserial part, increasing rapidly in size; mildly lobulate in outline; biconvex

and compressed in cross-section; length approximately two times width; periphery serrated; wall finely agglutinated; sutures curved, composed of coarser material than chambers; aperture interiomarginal fissure.

*Spiroplectammina subhaeringensis* (GRZYBOWSKI, 1896)  
Plate 1, Figure 12

- 1896 *Textularia subhaeringensis* n. sp. – GRZYBOWSKI, p. 285, pl. 9, figs. 13, 16.  
1926 *Textularia excolata* n. sp. – CUSHMAN, p. 585, pl. 15, figs. 9a–b.  
1929 *Textularia excolata* CUSHMAN – WHITE, p. 30, pl. 4, fig. 1.  
1946 *Spiroplectammina excolata* (CUSHMAN) – CUSHMAN, p. 27, pl. 5, figs. 9–10.  
1962 *Spiroplectammina excolata* (CUSHMAN) – HILLEBRANDT, p. 29, pl. 1, figs. 12–13.  
1966 *Spiroplectammina subhaeringensis* (GRZYBOWSKI) – HUSS, p. 39, pl. 6, figs. 15–17.  
1966 *Spiroplectammina excolata* (CUSHMAN) – HOFKER, p. 306, pl. 66, figs. 7–8.  
1970 *Spiroplectammina excolata* (CUSHMAN) – KIESEL, p. 194, pl. 3, fig. 24; pl. 16, fig. 1.  
1974 *Spiroplectammina subhaeringensis* (GRZYBOWSKI) – SZCZECZURA and POŻARYSKA, p. 31, pl. 3, figs. 16–17.  
1975 *Vulvulina haeringensis* (GÜMBEL) – PROTO PROTO DECIMA and DE BIASE, p. 91, pl. 1 fig. 23.

Description: Test planispiral in earlier three to four chambers, becoming biserial in adult part; four to five pairs of chambers in biserial part, increasing very rapidly in size; final chambers extremely inflated, overlapping earlier chambers; periphery subacute, slightly serrate; length approximately equal to width; sutures mildly compressed; wall finely agglutinated; aperture interiomarginal fissure.

Subfamily Vulvulininae SAIDOVA, 1981

Genus *Vulvulina* D'ORBIGNY emend. CICHA and ZAPLETALOVA, 1965

Type species: *Vulvulina capreolus* (DEFRANCE, 1826)

*Vulvulina advena* CUSHMAN, 1926  
Plate 1, Figure 13

- 1926 *Vulvulina advena* n. sp. – CUSHMAN, p. 32, pl. 4, fig. 9.  
1928 *Vulvulina advena* CUSHMAN – COLE, p. 206, pl. 1, fig. 24; pl. 3, fig. 17.  
1935 *Vulvulina advena* CUSHMAN – CUSHMAN, p. 9, pl. 2, figs. 1a–b.  
1949 *Vulvulina advena* CUSHMAN – CUVILLIER and SZAKÁLL, p. 18, pl. 6, fig. 3.  
1956 *Vulvulina haeringensis* (GÜMBEL) – HAGN, p. 115, pl. 9, figs. 7–8.  
1975 *Vulvulina haeringensis* (GÜMBEL) – PROTO DECIMA and DE BIASE, p. 91, pl. 1, fig. 4.  
1982 *Vulvulina haeringensis* (GÜMBEL) – SZTRÁKOS, pl. 2, fig. 7.  
1985 *Vulvulina haeringensis* (GÜMBEL) – GRÜNIG, p. 256, pl. 2, fig. 10.  
1987 *Vulvulina haeringensis* (GÜMBEL) – SZTRÁKOS, pl. 2, fig. 12.  
2006 *Vulvulina haeringensis* (GÜMBEL) – CIMERMAN et al., p. 16, pl. 1, fig. 6.  
2006 *Vulvulina advena* CUSHMAN – ORTIZ and THOMAS, p. 109, pl. 2, figs. 10–12. (non 9a–b).

Description: Test planispiral in juvenile stage, becoming biserial in adult part, last two or three chambers monoserial; triangular in outline; periphery subacute; biconvex, strongly compressed in cross-section; length approximately two times width; sutures limbate, slightly depressed; wall finely agglutinated; aperture interiomarginal fissure.

*Vulvulina jarvisi* CUSHMAN, 1932  
Plate 1, Figure 14

- 1932 *Vulvulina jarvisi* n. sp. – CUSHMAN, p. 84, pl. 10, fig. 20.  
1945 *Vulvulina jarvisi* CUSHMAN – CUSHMAN and STAINFORTH, p. 16, pl. 1, fig. 27.  
1953 *Vulvulina jarvisi* CUSHMAN – BECKMANN, p. 340, pl. 17, figs. 3–5.  
1982 *Vulvulina jarvisi* CUSHMAN – AGIP, pl. 3, fig. 2.  
1985 *Vulvulina haeringensis* (GÜMBEL) – GRÜNIG, p. 256, pl. 2, fig. 12.  
1998 *Vulvulina jarvisi* CUSHMAN – ROBERTSON, p. 16, pl. 1, fig. 3.

Description: Test planispiral in juvenile stage, becoming biserial in adult part, last two or three chambers monoserial, increasing rapidly in size, final chambers overlapping earlier chambers; oval in outline; periphery acute; biconvex, compressed in cross-section; length approximately two times width; sutures limbate, slightly depressed; wall finely agglutinated; aperture interiomarginal fissure.

Family Prolixoplectidae LOEBLICH and TAPPAN, 1985

Genus *Plectina* MARSSON, 1878

Type species: *Plectina ruthenica* (REUSS, 1851)

*Plectina dalmatina* (SCHUBERT, 1911)

Plate 1, Figure 15

1911 *Gaudryina dalmatina* n. sp. – SCHUBERT in LIEBUS, p. 939, pl. 3, fig. 5.

1950 *Plectina dalmatina* (SCHUBERT) – CITA, p. 86, pl. 6, fig. 10.

1972 *Plectina dalmatina* (SCHUBERT) – KUHN, p. pl. 3, figs. 22–25.

1975 *Plectina dalmatina* (SCHUBERT) – PROTO DECIMA and BIASE, p. 92, pl. 1, figs. 21–22. (non fig. 30.).

1987 *Plectina dalmatina* (SCHUBERT) – SZTRÁKOS, pl. 3, fig. 7.

Description: Test planispiral in juvenile stage, becoming biserial in adult part; chambers inflated, increasing rapidly in size; rounded subtriangular in outline; circular, oval in cross-section; wall finely agglutinated; aperture terminal.

*Plectina eocenica* CUSHMAN, 1936

Plate 1, Figure 16

1936 *Plectina eocenica* n. sp. – CUSHMAN, p. 43, pl. 3 figs. 1–3.

1975 *Plectina dalmatina* (SCHUBERT) – PROTO DECIMA and BIASE, p. 92, pl. 1, fig. 30.

Description: Test elongate, slender, triserial in juvenile portion, becoming biserial in adult portion; length approximately three times width; chambers inflated, distinct in adult portion, slightly overlapping; sutures barely visible, slightly depressed; wall coarsely agglutinated; aperture marginal.

Family Verneulinidae CUSHMAN, 1911

Subfamily Verneulininae CUSHMAN, 1911

Genus *Verneulina* D'ORBIGNY, 1839

Type species: *Verneulina tricarinata* D'ORBIGNY, 1840

*Verneulina* sp

Plate 1, Figure 17

Description: Test triserial; triangular in outline; triangular in cross-section; sutures barely visible; wall coarsely agglutinated; aperture terminal.

Superfamily Textulariacea EHRENBERG, 1838

Family Eggerellidae CUSHMAN, 1937

Subfamily Dorothisinae BALAKHMATOVA, 1972

Genus *Dorothia* PLUMMER emend. DESAI and BANNER, 1987

Type species: *Dorothia bulletta* (CARSEY, 1926)

*Dorothia textilaroides* (HANTKEN, 1875)

Plate 1, Figure 18

1875 *Gaudryina textilaroides* n. sp. – HANTKEN, p. 12, pl. 1, fig. 6.

1987 *Dorothia textilaroides* (HANTKEN) – SZTRÁKOS, pl. 2., fig. 22.

Description: Test trochospiral in juvenile stage, becoming biserial in adult part; chambers inflated, increasing rapidly in size; elongate in outline; circular, rounded in cross-section; sutures mildly depressed; wall coarsely agglutinated; aperture interiomarginal slit.

Subfamily Eggerellinae CUSHMAN, 1937

Genus *Martinottiella* CUSHMAN, 1933

Type species: *Martinottiella communis* (D'ORBIGNY, 1826)

*Martinottiella rhumbleri* (CUSHMAN, 1936)

Plate 1, Figure 19

1936 *Listerella rhumbleri* n. sp. – CUSHMAN, p. 38, pl. 6, fig. 4.

1979 *Martinottiella rhumbleri* (CUSHMAN) – SZTRÁKOS, pl. 6. fig. 4.

Description: Test elongate, slender, trochospiral in initial chambers, becoming monoserial in adult portion; chambers inflated, increasing in size regularly; circular in cross-section; wall coarsely agglutinated; aperture terminal, rounded.

Family Textulariidae EHRENBERG, 1838

Subfamily Textulariinae EHRENBERG, 1838

Genus *Textularia* DEFRANCE emend. NØRVANG, 1966

Type species: *Textularia sagittula* DEFRANCE, 1824

*Textularia crookshanki* HAQUE, 1956

Plate 1, Figure 20

1956 *Textularia crookshanki* n. sp. – HAQUE, p. 24, pl. 2, fig. 8.

1982 *Textularia crookshanki* HAQUE – SZTRÁKOS, pl. 2., figs. 5–6.

Description: Test biserial, slightly flaring towards apertural end; chambers two times as long as with, rapidly increasing in size; horn-like in outline; biconvex, mildly compressed, parallelogram-like in cross-section; periphery acute; sutures barely visible, mildly depressed, gently curved towards apertural end; wall finely agglutinated; aperture interiomarginal wide slit.

Remarks: Our specimens differ from the holotype of *Textularia crookshanki* in having more compressed test and stronger acute on periphery area.

*Textularia deperdita* D'ORBIGNY, 1846

Plate 2, Figures 1–2

1846 *Textularia deperdita* n. sp. – D'ORBIGNY, p. 244, pl. 14, figs. 23–25.

1985 *Textularia deperdita* D'ORBIGNY – PAPP and SCHMID, p. 84, pl. 78, figs. 7–9.

Description: Test biserial, slightly flaring towards apertural end; chambers two times as long as broad, rapidly increasing in size; conical in outline; biconvex, inflated in cross-section; periphery subacute; sutures distinct, straight; wall finely agglutinated; aperture interiomarginal wide slit.

*Textularia halkyardi* LALICKER, 1935

Plate 2, Figure 3

- 1875 *Textularia elongata* n. sp. – HANTKEN, p. 57, pl. 15, fig. 3.  
1935 *Textularia halkyardi* n. sp. – LALICKER, p. 45, pl. 2, figs. 5a–c.  
1962 *Bolivina elongata* (HANTKEN) – MAJZON, pl. 43, fig. 3.  
1982 *Textularia elongata* HANTKEN – SZTRÁKOS, p. 20, pl. 2, fig. 10.  
1987 *Textularia elongata* HANTKEN – SZTRÁKOS, pl. 1, fig. 16.  
1993 *Textularia elongata* HANTKEN – MATHÉLIN and SZTRÁKOS, p. 40, pl. 24, fig. 12.  
2002 *Textularia elongata* HANTKEN – HORVÁTH, p. 35, pl. 2, fig. 8.

Description: Test elongate, biserial, slightly flaring towards apertural end; chambers strongly inflated, rapidly increasing in size; conical in outline; biconvex, inflated, broadly oval in cross-section; periphery smooth; sutures barely visible, curved; wall finely agglutinated; aperture interiomarginal wide slit.

Remarks: HANTKEN established the species *Textularia elongata* in 1875. He described this species from the world-famous Upper Eocene and Lower Oligocene Buda Marl Formation from Budapest, Hungary. The species name *Textularia elongata* was described as new name at five times before 1875 (*Textularia elongata* HAGENOW, 1842; *Textularia elongata* CORNUEL, 1848; *Textularia elongata* JONES, 1850; *Textularia elongata* D'ORBIGNY, 1852 and *Textularia elongata* EHRENBERG, 1873) thus, this name is invalid after 1842. In my view, the first valid synonym of *Textularia elongata* (HANTKEN, 1875) is *Textularia halkyardi* LALICKER, 1932 who described it as a new species from the Upper Eocene of Biarritz, France.

*Textularia globosa* (HANTKEN, 1875)

Plate 2, Figures 4–5

- 1875 *Textularia globosa* n. sp. – HANTKEN, p. 58, pl. 15, figs. 5a–b.

Description: Test squattish, biserial; oval in outline; circular in cross-section; chambers mildly inflated; increasing in size; periphery mildly undulate; sutures distinct, strongly curved; wall finely agglutinated; aperture interiomarginal wide slit.

*Textularia* cf. *partschii* CZIZEK, 1848

Plate 2, Figures 6–7

- 1848 *Textularia partschii* n. sp. – CZIZEK, p. 148, pl. 13, figs. 22–24.  
1893 *Textularia partschii* CZIZEK – EGGER, p. 267, pl. 6, figs. 22–23.

Description: Test biserial, rapidly flaring towards apertural end; chambers inflated, rapidly increasing in size; triangular in outline; biconvex, circular in cross-section; periphery subacute; sutures barely visible, mildly depressed, strongly curved towards apertural end; wall finely agglutinated; aperture interiomarginal slit.

Remarks: Originally, *Textularia partschii* CZIZEK was described from the Badenian of Vienna Basin, Austria. The figured specimen is a bit wider and larger than the holotype. It differs from typical *Textularia partschii* in having larger, less elongate final chambers.

*Textularia lanceolata* (KARRER, 1868)

Plate 2, Figure 8

- 1868 *Plecanium lanceolatum* n. sp. – KARRER, p. 129, pl. 1, fig. 2.  
1979 *Textularia lanceolata* (KARRER) – SZTRÁKOS, pl. 4, figs. 1a–b.  
1982 *Textularia lanceolata* (KARRER) – SZTRÁKOS, pl. 3, fig. 1.  
1985 *Textularia lanceolata* (KARRER) – KORECZNÉ LAKY and NAGYNÉ GELLAI, pl. 91, fig. 2.

Description: Test strongly elongate, biserial; circular in cross-section; periphery subacute; sutures barely visible; wall finely agglutinated; aperture interiomarginal slit.

*Textularia pala* CZJZEK, 1848

Plate 2, Figure 9

1848 *Textularia pala* n. sp. – CZJZEK, p. 148, pl. 13, figs. 25–27.

1979 *Textularia pala* CZJZEK – SZTRÁKOS, pl. 4., fig. 3.

1985 *Textularia pala* CZJZEK – KORECZNÉ-LAKY and NAGYNÉ-GELLAI, pl. 6, fig. 14.

Description: Test squattish, biserial; oval in outline; biserial, strongly compressed in cross-section; chambers two times as long as broad, rapidly increasing in size; periphery mildly undulate; sutures distinct, strongly curved; wall finely agglutinated; aperture interiomarginal wide slit.

*Textularia* sp. 1.

Plate 2, Figures 10–11

Description: Test biserial, rapidly flaring towards apertural end; chambers inflated, rapidly increasing in size; triangular in outline; biconvex, oval in cross-section; sutures barely visible; wall finely agglutinated; aperture interiomarginal slit.

*Textularia* sp. 2.

Plate 2, Figure 12

Description: Test strongly elongate, biserial; final chambers elongate; circular in cross-section; periphery acute; sutures barely visible; wall finely agglutinated; aperture interiomarginal slit.

*Textularia* sp. 3.

Plate 2, Figures 13–14

Description: Test biserial, rapidly flaring towards apertural end; chambers inflated, rapidly increasing in size; triangular in outline; biconvex, mildly compressed in cross-section; periphery corrugated; sutures barely visible; wall finely agglutinated; aperture interiomarginal fissure.

Family Pseudogaudryinidae LOEBLICH and TAPPAN, 1985

Subfamily Pseudogaudryininae LOEBLICH and TAPPAN, 1985

Genus *Clavulinoides* CUSHMAN emend. BANNER and DESAI, 1985

Type species: *Clavulinoides trilatera* (CUSHMAN, 1926)

*Clavulinoides alpina* CUSHMAN, 1936

Plate 2, Figures 15–16

1936 *Clavulinoides alpina* n. sp. – CUSHMAN, p. 22, pl. 3, fig. 16.

1937 *Clavulinoides alpina* CUSHMAN – CUSHMAN, p. 127, pl. 18, figs. 13–15.

1975 *Tritaxia alpina* (CUSHMAN) – PROTO DECIMA and BIASE, p. 91, pl. 1, fig. 12.

1987 *Tritaxia alpina* (CUSHMAN) – SZTRÁKOS, pl. 2, fig. 19.

2005 *Tritaxia alpina* (CUSHMAN) – ANAN, p. 21, pl. 1, fig. 8.

Description: Test squattish, initially triserial, becoming monoserial; chambers rapidly flaring towards apertural end; length approximately two times as long as broad; triangular in cross-section; chambers slightly inflated; sutures invisible; wall finely agglutinated; aperture terminal.

*Clavulinoides lakiensis elongata* HAQUE, 1956

Plate 2, Figure 17

1956 *Clavulinoides lakiensis* HAQUE var. *elongata* n. ssp. – HAQUE, p. 45, pl. 21, figs. 13a–b.

1987 *Tritaxia dimidiata* (CUSHMAN and BERMÚDEZ) – SZTRÁKOS, pl. 2, fig. 21.

Description: Test elongate, initially triserial, becoming monoserial; chambers rapidly flaring towards apertural end; length approximately four times as long as broad; triangular in cross-section; chambers slightly inflated; sutures barely visible, slightly depressed; wall finely agglutinated; aperture terminal, rounded, extended.

*Clavulinoides procerus* n. sp.

Plate 2, Figure 20

Derivatio nominis: Named after long, elongate test.

Holotype: The specimen on Plate 2, Figure 20. (Inventory number: M 2008.117.1)

Dimension of holotype: length 0.95 mm, width 0.33 mm.

Description: Test elongate, initially triserial, becoming monoserial in adult portion; length approximately three times as long as broad; triserial portion extremely acute and sharp; mildly curved in outline; triangular in cross-section; periphery serrated; chambers increasing gradually in size; sutures barely visible, slightly depressed; wall coarsely agglutinated; aperture terminal, rounded, extended.

Type locality: Bkcs 18 borehole, Transdanubian Central Range, Bakony Mountains.

Type strata: In the B18 borehole, at 295.3 m. Padrag Marl Formation (Middle Eocene, Bartonian)

Remarks: *Clavulinoides procerus* differs from *Clavulinoides lakiensis elongata* HAQUE in having slender test, smaller and slender triserial portion and wider monoserial portion.

*Clavulinoides szabói* (HANTKEN, 1868)

Plate 2, Figures 18–19

- 1868 *Rhabdognium Szabói* n. sp. – HANTKEN, p. 90, pl. 1, fig. 18.  
1875 *Clavulina szabói* HANTKEN – HANTKEN, p. 13, pl. 1, figs. 9a–d.  
1903 *Clavulina szabói* HANTKEN – WOJCIK, p. 498, pl. 6, fig. 20.  
1932 *Clavulina szabói* HANTKEN – PROTESCU, p. 88, pl. 1, figs. 1–2.  
1937 *Clavulinoides szabói* (HANTKEN) – CUSHMAN, p. 133, pl. 18, figs. 33a–b, 34.  
1937 *Clavulinoides szabói* (HANTKEN) – CUSHMAN, p. 134, pl. 18, figs. 35a–b, 36.  
1946 *Clavulina szabói* HANTKEN – KAPTARENKO and CSERNUSZOVA, p. 229, pl. 2, fig. 10.  
1946 *Clavulinoides szabói* (HANTKEN) – VAN BELLEN, p. 86, pl. 13, fig. 16.  
1949 *Clavulinoides szabói* (HANTKEN) – CUVILLIER and SZAKALL, p. 24, pl. 10, fig. 4.  
1950 *Clavulinoides szabói* (HANTKEN) – CITA, p. 85, pl. 6, fig. 8.  
1953 *Clavulinoides szabói* (HANTKEN) – HAGN, p. 39, fig. 2.  
1956 *Clavulinoides szabói* (HANTKEN) – HAGN, p. 116, pl. 10, fig. 1.  
1956 *Clavulina szabói* HANTKEN – KAPTARENKO and CSERNUSZOVA, pl. 7, fig. 9.  
1972 *Clavulinoides szabói* (HANTKEN) – MAJZON, p. 114, pl. 1, figs. 3–9, 11–16, 18–19, 21–23; pl. 2, figs. 1–8, 14–20.  
1973 *Clavulinoides szabói* (HANTKEN) – NAGYNÉ GELLAI, p. 445, pl. 2, fig. 1.  
1975 *Tritaxia szabói* (HANTKEN) – BRAGA and GRÜNIG, p. 103, pl. 4, figs. 1–2.  
1979 *Tritaxia szabói* (HANTKEN) – SZTRÁKOS, pl. 5, fig. 8.  
1985 *Tritaxia szabói* (HANTKEN) – GRÜNIG, p. 257, pl. 2, figs. 22–23.  
1985 *Tritaxia szabói* (HANTKEN) – KORECZNÉ-LAKY and NAGYNÉ-GELLAI, pl. 7, figs. 9–10; pl. 22, figs. 1–2, 4.  
1988 *Clavulina szabói* HANTKEN – GELLAI – NAGY, pl. 4, figs. 1–7.  
1998 *Tritaxia szabói* (HANTKEN) – CÍCHA et al., p. 132, pl. 7, figs. 12–13.  
1999 *Tritaxia szabói* (HANTKEN) – OZSVART, p. 83, pl. 1, fig. 2.  
2002 *Tritaxia szabói* (HANTKEN) – HORVÁTH, p. 35, pl. 2, fig. 10.

Description: Test initially triserial, becoming monoserial in adult portion; length approximately two and one-half times as long as broad; triserial portion acute; triangular in cross-section; chambers increasing gradually in size; sutures barely visible, slightly depressed; wall coarsely agglutinated; aperture terminal, rounded.

Family Valvulaminidae LOEBLICH and TAPPAN, 1986

Genus *Clavulina* D'ORBIGNY, 1826

Type species: *Clavulina parisiensis* D'ORBIGNY, 1826

*Clavulina terterensis* KHALILOV, 1958  
Plate 2, Figure 21

1958 *Clavulina terterensis* n. sp. – KHALILOV, p. 9, pl. 2, figs. 1–2.

Description: Test elongate, initially triserial, becoming monoserial in adult portion; monoserial part three-four time longer as triserial part; four monoserial chambers; length approximately three-four times as long as broad; triserial chambers inflated, increasing gradually in size; monoserial chambers inflated, equal in size; final chamber stronger inflated; circular in cross-section; sutures distinct, strongly depressed; wall finely agglutinated; aperture terminal.

Remarks: This form was originally described from Caucasus, northeastern Azerbaidzhan. It occurs only in the Sv 1 borehole in Hungary.

Genus *Cylindroclavulina* BERMÚDEZ and KEY, 1952

Type species: *Cylindroclavulina bradyi* (CUSHMAN, 1911)

*Cylindroclavulina colomi* HAGN, 1956  
Plate 2, Figures 22–23

1956 *Cylindroclavulina colomi* n. sp. – HAGN, p. 122, pl. 11, fig. 14.

1987 *Cylindroclavulina colomi* HAGN – SZTRÁKOS, pl. 3, fig. 8.

Description: Test initially spherical, planispiral (?), becoming monoserial in final chambers; length two times as long as width; initially chambers increasing rapidly in size, monoserial chambers equal in size; final chambers width greater than height; final chamber slightly pointed; sutures distinct, depressed; wall finely agglutinated; wide terminal aperture.

*Cylindroclavulina rudislostia* (HANTKEN, 1875)  
Plate 2, Figure 24

1875 *Clavulina cylindrica* n. sp. – HANTKEN, p. 14, pl. 1, fig. 8.

1987 *Cylindroclavulina rudislostia* HANTKEN – SZTRÁKOS, pl. 3, fig. 9.

Description: Test squattish, initially spherical, planispiral (?), becoming monoserial in final chambers; length two times as long as width; chambers increasing rapidly in size; sutures invisible; wall finely agglutinated; aperture terminal on short neck.

Subfamily Tritaxilininae LOEBLICH and TAPPAN, 1986

Genus *Tritaxilina* CUSHMAN emend. MARIE 1941

Type species: *Tritaxilina caperata* (BRADY, 1881)

*Tritaxilina pupa* (GÜMBEL, 1868)  
Plate 2, Figures 25–27

1868 *Gaudryna pupa* n. sp. – GÜMBEL, p. 602, pl. 18, figs. 8–15.

1937 *Tritaxilina pupa* (GÜMBEL) – CUSHMAN, p. 156, pl. 3, fig. 5.

1975 *Tritaxilina pupa* (GÜMBEL) – PROTO DECIMA and DE BIASE, p. 93, pl. 1, figs. 24–27.

1982 *Tritaxilina hantkeni* CUSHMAN – SZTRÁKOS, pl. 4, figs. 13a–b.

1987 *Tritaxilina pupa* (GÜMBEL) – SZTRÁKOS, pl. 3, figs. 11–12.



Description: Test squattish, initially triserial, becoming biserial in adult portion; test rapidly flaring towards apertural end; triangular in outline; circular in cross-section; sutures mildly depressed; wall rather coarsely agglutinated; aperture interiomarginal slit.

*Tritaxilina* sp.  
Plate 3, Figure 1

Description: Test squattish, initially triserial, becoming biserial in adult portion; test rapidly flaring towards apertural end; triangular in outline; circular in cross-section; sutures distinct, strongly depressed; wall rather coarsely agglutinated; aperture interiomarginal slit.

Suborder Miliolina DELAGE and HÉROUARD, 1896  
Superfamily Cornuspiracea SCHULTZE, 1854  
Family Cornuspiridae SCHULTZE, 1854  
Subfamily Cornuspirinae SCHULTZE, 1854

Genus *Cornuspira* SCHULTZE, 1854

Type species: *Cornuspira foliacea* (PHILIPPI, 1844)

*Cornuspira involvens* (REUSS, 1850)  
Plate 3, Figures 2–3

- 1850 *Operculina involvens* n. sp. – REUSS, p. 370, pl. 46, fig. 20.
- 1864 *Cornuspira archimedis* (REUSS) – STACHE, p. 180, pl. 22, figs. 1a–b.
- 1875 *Cornuspira involvens* (REUSS) – HANTKEN, p. 16, pl. 2, fig. 2.
- 1884 *Cornuspira involvens* (REUSS) – BRADY, p. 200, pl. 60, figs. 1–3.
- 1926 *Cornuspira involvens* (REUSS) – CHAPMAN, p. 27, pl. 3, figs. 1–2.
- 1982 *Cornuspira involvens* (REUSS) – AGIP, pl. 7, figs. 10–10p.
- 1994 *Cornuspira involvens* (REUSS) – JONES, p. 26, pl. 11, figs. 1–3.

Description: Test planispiral, evolute, seven whorls visible; chambers unidentified; circular in outline; strongly compressed in cross-section; periphery smooth; wall calcareous, porcellaneous; aperture oval opening at end of final chamber; without visible tooth.

Superfamily Miliolacea EHRENBERG, 1839  
Family Spiroloculinidae WIESNER, 1920

Genus *Adelosina* D'ORBIGNY, 1826

Type species: *Adelosina laevigata* D'ORBIGNY, 1826

*Adelosina* sp.  
Plate 3, Figure 4

Description: Test circular in outline, axial periphery subacute, quinqueloculine coiling, oval or triangular in cross-section; chambers indistinct, curved, elongated and overlapped by the subsequent chambers; wall calcareous, porcellaneous, smooth; aperture an elongate opening at end of final chamber with tiny pores.

Genus *Spiroloculina* D'ORBIGNY, 1826

Type species: *Spiroloculina depressa* D'ORBIGNY, 1826

*Spiroloculina obscura* (CUSHMAN and TODD, 1944)

Plate 3, Figure 5

- 1852 *Spiroloculina grateloupi* n. sp. – D'ORBIGNY, p. 298, p. 161, pl. 1, fig. 3.  
1882 *Spiroloculina grateloupi* n. sp. – TERQUEM, p. 155, pl. 16, fig. 6.  
1905 *Spiroloculina grateloupi* TERQUEM – FORNASINI, p. 4, pl. 1, fig. 3.  
1944 *Spiroloculina obscura* nomen nudum – CUSHMAN and TODD, p. 20, pl. 3, figs. 24–25.

Description: Test planispiral; oval in outline; slightly compressed in cross-section; length approximately two times width; chambers barely visible in initial portion, in final whorl tubular, increasing rapidly in size, overlapping earlier chambers; periphery smooth; sutures barely visible, mildly depressed; wall calcareous, smooth, porcellaneous; aperture terminal, circular on short neck.

*Spiroloculina bicarinata* TERQUEM, 1882

Plate 3, Figures 6–7

- 1882 *Spiroloculina bicarinata* n. sp. – TERQUEM, p. 155, pl. 16, fig. 5.  
1944 *Spiroloculina bicarinata* TERQUEM – CUSHMAN and TODD, p. 8, pl. 2, fig. 12.

Description: Test planispiral, evolute; oval in outline; long, tubular chambers, increasing rapidly in size; periphery smooth, channelled; sutures distinct, slightly swollen; wall calcareous, smooth, porcellaneous; aperture terminal, circular on short neck.

*Spiroloculina jarvisi* CUSHMAN and TODD, 1944

Plate 3, Figures 8–9

- 1944 *Spiroloculina jarvisi* n. sp. – CUSHMAN and TODD, p. 14, pl. 3, fig. 9.  
1951 *Spiroloculina jarvisi* CUSHMAN and TODD – CUSHMAN and STAINFORTH, p. 145, pl. 25, fig. 36.  
1956 *Spiroloculina jarvisi* CUSHMAN and TODD – HOFKER, p. 91. Text-figs. 8e–f.

Description: Test planispiral, evolute; length approximately equal with width; circular in outline; mildly compressed in cross-section; tubular chambers, increasing rapidly in size; sutures distinct, slightly swollen; wall calcareous, smooth, porcellaneous; aperture terminal, circular on short neck.

Family Hauerinidae SCHWAGER, 1876

Subfamily Hauerininae SCHWAGER, 1876

Genus *Quinqueloculina* D'ORBIGNY, 1826

Type species: *Quinqueloculina seminula* (LINNE, 1758)

*Quinqueloculina buchiana* D'ORBIGNY, 1846

Plate 3, Figure 10

- 1846 *Quinqueloculina buchiana* n. sp. – D'ORBIGNY, p. 289, pl. 18, figs. 10–12.  
1974 *Quinqueloculina buchiana* D'ORBIGNY – ŁUCZKOWSKA, p. 46, pl. 4, figs. 1–4.  
1985 *Quinqueloculina buchiana* D'ORBIGNY – PAPP and SCHMID, p. 99, pl. 93, figs. 1–7.  
1985 *Quinqueloculina buchiana* D'ORBIGNY – KORECZNÉ-LAKY and NAGYNÉ-GELLAI, pl. 86, figs. 3a–b.

Description: Test nearly twice as long as broad, periphery subacute; trapezoidal in outline, triangular in cross-section with sharp edge; chambers increasing gradually in size, each chamber comprising one and half coil and overlapped by subsequent chambers; sutures distinct, mildly compressed; wall calcareous, smooth, porcellaneous; aperture terminal, circular.

Remarks: *Q. buchiana* differs from all other species of *Quinqueloculina* in having a sharp periphery of test.

*Quinqueloculina juleana* D'ORBIGNY, 1846

Plate 3, Figure 11

- 1846 *Quinqueloculina juleana* n. sp. – D'ORBIGNY, p. 298, pl. 20, figs. 1–3.  
1955 *Quinqueloculina juleana* D'ORBIGNY – BHATIA, p. 672, pl. 66, fig. 9.  
1958 *Quinqueloculina juleana* D'ORBIGNY – BATJES, p. 103, pl. 1, fig. 16.  
1961 *Quinqueloculina juleana* D'ORBIGNY – KAASSCHIETER, p. 149, pl. 2, figs. 14–15.  
1985 *Quinqueloculina juleana* D'ORBIGNY – PAPP and SCHMID, p. 104, pl. 100, figs. 1–4.  
1985 *Quinqueloculina juleana* D'ORBIGNY – KORECZNÉ-LAKY and NAGYNÉ-GELLAI, pl. 9, fig. 2.

Description: Test elongate in outline, oval in cross-section; wall calcareous, smooth, porcellaneous; aperture terminal, circular with single small tooth.

*Quinqueloculina seminula* (LINNÉ, 1758)

Plate 3, Figure 12

- 1758 *Serpula seminulum* n. sp. – LINNÉ, p. 786, pl. a, figs. 1a–c (fide: ELLIS and MESSINA).  
1949 *Quinqueloculina seminula* (LINNÉ) – CUVILLIER and SZAKALL, p. 37, pl. 17, fig. 4.  
1955 *Quinqueloculina seminula* (LINNÉ) – BHATIA, p. 674, pl. 67, fig. 8.  
1956 *Quinqueloculina seminula* (LINNÉ) – GULLENTOPS, p. 9, pl. 1, fig. 1.  
1958 *Quinqueloculina seminula* (LINNÉ) – BATJES, p. 102, pl. 1, fig. 15.  
1961 *Quinqueloculina seminula* (LINNÉ) – KAASSCHIETER, p. 147, pl. 2, figs. 5–6.  
1963 *Quinqueloculina seminula* (LINNÉ) – KÜMMERLE, p. 27, pl. 1, fig. 6.  
1970 *Quinqueloculina seminula* (LINNÉ) – LE CALVEZ, p. 40, pl. 46, figs. 6–7.

Description: Test large, oval or circular in outline, oval in cross-section; chambers increasing gradually in size, each chamber comprising one and half coil and overlapped by subsequent chambers; wall calcareous, smooth, porcellaneous; sutures barely visible; terminal aperture with two sharp teeth.

*Quinqueloculina* sp. 1.

Plate 3, Figure 13

Description: Test large, milioline, oval in lateral view, triangular in cross-section; periphery subacute; chambers increasing gradually in size, each chamber comprising one and half coil and overlapped by subsequent chambers; wall calcareous, smooth, porcellaneous; sutures barely visible; terminal aperture on short neck.

*Quinqueloculina* sp. 2.

Plate 3, Figure 14

Description: Test large, milioline, oval in lateral view, strongly compressed in cross-section; each chamber comprising one and half coil and overlapped by subsequent chambers; wall calcareous, smooth, porcellaneous; sutures barely visible; terminal aperture.

Subfamily Miliolinellinae VELLA, 1957

Genus *Pyrgo* DEFRANCE, 1824

Type species: *Pyrgo laevis* DEFRANCE, 1824

*Pyrgo bulloides* (D'ORBIGNY, 1826)

Plate 3, Figure 15

- 1826 *Biloculina bulloides* n. sp. – D'ORBIGNY, p. 13, pl. 1, fig. 1.  
1961 *Pyrgo bulloides* (D'ORBIGNY) – KAASSCHIETER, p. 167, pl. 5, fig. 18.  
1970 *Pyrgo bulloides* (D'ORBIGNY) – LE CALVEZ, p. 54, pl. 6, fig. 5.

Description: Test large, biloculine; oval in outline, subcircular in cross-section; two chambers visible in final whorl; wall calcareous, smooth, porcellaneous; aperture terminal with tooth.

Remarks: *Pyrgo bulloides* (D'ORBIGNY) is very common in Paleogene sediments of the HPB, especially in shallow water environments.

Subfamily Sigmoidinitinae ŁUCZKOWSKA, 1974

Genus *Articulina* D'ORBIGNY, 1825

Type species: *Articulina nitida* D'ORBIGNY, 1825

*Articulina curta* LE CALVEZ, 1947  
Plate 3, Figure 16

1947 *Articulina curta* n. sp. – LE CALVEZ, p. 39, pl. 4, fig. 87.

1970 *Articulina curta* LE CALVEZ – LE CALVEZ, p. 67, pl. 10, fig. 8.

Description: Test squattish, elongated in outline, oval in cross-section; initial chambers rather triloculine, later monoserial; sutures distinct, mildly depressed; wall calcareous, porcellaneous with numerous fine, longitudinal ribs; aperture circular, terminal with thickened lip.

*Articulina laevigata* TERQUEM, 1882  
Plate 3, Figure 17

1882 *Articulina laevigata* n. sp. – TERQUEM, p. 151, pl. 15, figs. 27–31.

1947 *Articulina laevigata* TERQUEM – KAASSCHIETER, p. 158, pl. 4, figs. 15–17.

Description: Test elongated, oval in cross-section; initial chambers rather triloculine or quinqueloculine, later monoserial; sutures distinct, mildly depressed; wall calcareous, porcellaneous with numerous fine, longitudinal ribs; aperture circular, terminal with thickened lip.

Remarks: *A. laevigata* TERQUEM is distinguished from all other species of *Articulina* by long, monoserial final chambers.

*Articulina nitida* D'ORBIGNY, 1826  
Plate 3, Figure 18

1826 *Articulina nitida* n. sp. – D'ORBIGNY, p. 300, fig. 22.

1961 *Articulina nitida* D'ORBIGNY – KAASSCHIETER, p. 157, pl. 4, fig. 11.

1970 *Articulina nitida* D'ORBIGNY – LE CALVEZ, p. 69, pl. 4, fig. 6.

Description: Test squattish, oval in outline, subcircular in cross-section; initial chambers triloculine, later monoserial; sutures distinct, barely sutures; wall calcareous, porcellaneous with numerous fine, longitudinal ribs; aperture circular, terminal with thickened lip.

Family Miliolidae EHRENBERG, 1839

Subfamily Miliolinae EHRENBERG, 1839

Genus *Miliola* LAMARCK, 1804

Type species: *Miliola trigonula* (LAMARCK, 1804)

*Miliola prisca* (D'ORBIGNY, 1826)  
Plate 3, Figures 19–20

1826 *Quinqueloculina prisca* n. sp. – D'ORBIGNY, p. 136, pl. 6, fig. 5.

1970 *Miliola prisca* (D'ORBIGNY) – LE CALVEZ, p. 45, pl. 6, figs. 1–2; pl. 46, figs. 4–5.

Description: Test large, milioline, oval in lateral view; large-size chambers increasing gradually in size, each chamber comprising one and half coil and overlapped by subsequent chambers; wall calcareous, smooth, porcellaneous; sutures barely visible; terminal aperture with many pores.

Remarks: The only difference between *Quinqueloculina* and *Miliola* is the aperture. *Quinqueloculina* have large, open aperture with or without any tooth. *Miliola* have closed aperture with many pores.

*Miliola strigillata* (D'ORBIGNY, 1850)

Plate 3, Figures 21–22

1850 *Triloculina strigillata* n. sp. – D'ORBIGNY, p. 409, pl. 1, figs. 7a–b.

1882 *Triloculina strigillata* D'ORBIGNY – TERQUEM, p. 169, pl. 17, fig. 25.

1905 *Miliola strigillata* (D'ORBIGNY) – FORNASINI, p. 60, pl. 1, fig. 7.

Description: Test large, milioline, subcircular in lateral view; oval in cross-section, large-size chambers increasing gradually in size, each chamber comprising one and half coil and overlapped by subsequent chambers; wall calcareous, porcellaneous with numerous fine, longitudinal ribs on periphery; sutures barely visible; terminal aperture with many pores.

Remarks: *Miliola strigillata* (D'ORBIGNY) is distinguished from *Miliola prisca* (D'ORBIGNY) by its numerous fine, longitudinal ribs on periphery.

Genus *Massilina* SCHLUMBERGER, 1893

Type species: *Massilina secans* (D'ORBIGNY, 1826)

*Massilina* sp. 1.

Plate 3, Figure 23

Description: Test large, milioline, oval in lateral view; triangular in cross-section; periphery acute, slightly carinate; chambers increasing gradually in size, each chamber comprising one and half coil and overlapped by subsequent chambers; wall calcareous, porcellaneous with minute pores; sutures distinct; terminal, circular aperture.

Family Peneroplidae SCHULTZE, 1854

Genus *Spirolina* LAMARCK, 1804

Type species: *Spirolina cylindrica* (LAMARCK, 1804)

*Spirolina mariei* LE CALVEZ, 1952

Plate 3, Figures 24–25

1952 *Spirolina mariei* n. sp. – LE CALVEZ, p. 25, pl. 2, figs. 17–18.

1970 *Spirolina mariei* LE CALVEZ – LE CALVEZ, p. 76, pl. 15, fig. 2.

Description: Test squattish, planispiral in initial five-six chambers, becoming monoserial in adult part (two or three chambers, circular in cross-section; length approximately two times of width; initial chambers slowly increasing in size; wall calcareous, perforate with numerous fine, longitudinal ribs; sutures distinct, strongly depressed; terminal aperture.

*Spirolina pedum* D'ORBIGNY, 1826

Plate 3, Figure 26

1826 *Spirolina pedum* n. sp. – D'ORBIGNY, p. 121, pl. 1, figs. 20–24.

1952 *Spirolina pedum* D'ORBIGNY – LE CALVEZ, p. 23, pl. 2, fig. 15.

1970 *Spirolina pedum* D'ORBIGNY – LE CALVEZ, p. 76, pl. 15, fig. 4.

Description: Test strongly elongate, planispiral in initial five-six chambers, becoming monoserial in adult part (six to nine chambers, circular in cross-section; length approximately six-seven times width; initial chambers slowly increasing in size; wall calcareous, perforate with numerous fine, longitudinal ribs; sutures distinct, strongly depressed; terminal aperture.

Remarks: *Spirolina pedum* D'ORBIGNY is distinguished from *Spirolina mariei* LE CALVEZ by its long monoserial part (six to nine chambers) of test.

*Spirolina* sp.  
Plate 3, Figure 27

Description: Test elongate, planispiral in initial five-six chambers, becoming monoserial in adult part, circular in cross-section; length approximately three times width; initial chambers slowly increasing in size; wall calcareous, perforate with numerous fine, longitudinal ribs; sutures distinct, strongly depressed; terminal aperture.

Suborder Lagenina DELAGE and HÉROUARD, 1896  
Superfamily Nodosariacea EHRENBERG, 1838  
Family Nodosariidae EHRENBERG, 1838  
Subfamily Nodosariinae EHRENBERG, 1838

Genus *Chrysalogonium* SCHUBERT, 1907

Type species: *Chrysalogonium polystoma* (SCHWAGER, 1866)

*Chrysalogonium tympanipectiformis* (SCHWAGER, 1866)  
Plate 4, Figure 1–2

1866 *Nodosaria tympanipectiformis* n. sp. – SCHWAGER, p. 215, pl. 5, fig. 34.

Description: Test elongate, monoserial, straight to slightly curved, circular in cross-section; four chambers increasing gradually in size, first chamber ovoidal, inflated, later chambers becoming more elongate, final chamber tubular; wall calcareous, smooth, finely perforate; sutures distinct; terminal aperture.

*Chrysalogonium* sp.  
Plate 4, Figure 3

Description: Test elongate, monoserial, straight, circular in cross-section; first chamber elongate, tubular, later chambers missing; wall calcareous, smooth, finely perforate; suture distinct; terminal aperture.

Genus *Dentalina* RISSO, 1826

Type species: *Dentalina cuvieri* (D'ORBIGNY, 1826)

*Dentalina aboleta* SCHWAGER, 1865  
Plate 4, Figure 6

1865 *Dentalina aboleta* n. sp. – SCHWAGER, p. 105, pl. 3, figs. 5, 8.

Description: Test squattish, monoserial, straight, subcircular in cross-section; ovoidal chambers increasing gradually in size; wall calcareous, hyaline, finely perforate; aperture terminal, radiate.

Remarks: *Dentalina aboleta* SCHWAGER has been reported from the Upper Jurassic of Northern Calcareous Alps.

*Dentalina* sp. cf. *D. acuta* D'ORBIGNY, 1846  
Plate 4, Figure 4

- 1846 *Dentalina acuta* n. sp. – D'ORBIGNY, p. 56, pl. 2, figs. 40–43.  
1949 *Dentalina acuta* D'ORBIGNY – CUVILLIER and SZAKÁLL, p. 75, pl. 28, figs. 3–4.  
1985 *Dentalina acuta* D'ORBIGNY – PAPP and SCHMID, p. 35, pl. 18, figs. 1–6.

Description: Test elongate, monoserial, straight, circular in cross-section; chambers increasing gradually in size; wall calcareous, hyaline, finely perforate with numerous fine, longitudinal ribs; aperture terminal, radiate.

*Dentalina antennula* D'ORBIGNY, 1846  
Plate 4, Figure 12

- 1846 *Dentalina antennula* n. sp. – D'ORBIGNY, p. 53, pl. 2, figs. 29–30.  
1851 *Dentalina philippii* n. sp. – REUSS, p. 60, pl. 3, fig. 5.  
1985 *Dentalina antennula* D'ORBIGNY – PAPP and SCHMID, p. 33, pl. 15, figs. 7–9.  
2004 *Dentalina antennula* D'ORBIGNY – EILAND and GUDMUNDSSON, p. 198, pl. 1, fig. P; pl. 2, fig. C.

Description: Test elongate, monoserial, straight, circular in cross-section; first and last chambers slightly inflated; wall calcareous, hyaline, finely perforate with numerous fine, longitudinal ribs; sutures barely visible; aperture terminal, radiate.

*Dentalina approximata* (REUSS, 1866)  
Plate 4, Figure 5

- 1866 *Nodosaria (Dentalina) approximata* n. sp. – REUSS, p. 134, pl. 2, fig. 22.  
1875 *Dentalina approximata* REUSS – HANTKEN, p. 26, pl. 3, fig. 5.  
1975 *Dentalina approximata* REUSS – SAMUEL, p. 122, pl. 69, figs. 5–6.

Description: Test elongate, monoserial, slightly curved, circular in cross-section; squattish chambers increasing gradually in size; wall calcareous, smooth, hyaline, finely perforate; aperture terminal, radiate.

*Dentalina cornicula* (D'ORBIGNY, 1826)  
Plate 4, Figures 7–8

- 1826 *Nodosaria (Dentalina) cornicula* n. sp. – D'ORBIGNY, p. 255, pl. 9, fig. 56.  
1949 *Dentalina cornicula* (D'ORBIGNY) – CUVILLIER and SZAKÁLL, p. 75, pl. 28, fig. 1.

Description: Test elongate, monoserial, straight, circular in cross-section; chambers increasing gradually in size, final chamber acute; wall calcareous, hyaline with numerous strong, longitudinal ribs; aperture terminal, radiate.

*Dentalina debilis* HANTKEN, 1868  
Plate 4, Figure 9

- 1868 *Dentalina debilis* n. sp. – HANTKEN, pl. 2, fig. 27.  
1875 *Dentalina debilis* HANTKEN – HANTKEN, p. 28, pl. 13, fig. 10.

Description: Test elongate, monoserial, straight to slightly curved, circular in cross-section; chambers increasing gradually in size; wall calcareous, hyaline, finely perforate; sutures distinct; aperture terminal, radiate.

*Dentalina elegans* D'ORBIGNY, 1846  
Plate 4, Figure 10

1846 *Dentalina elegans* n. sp. – D'ORBIGNY, p. 45, pl. 1, figs. 52–56.  
1985 *Dentalina elegans* D'ORBIGNY – PAPP and SCHMID, p. 28, pl. 10, figs. 1–5.

Description: Test elongate, monoserial, straight to slightly curved, circular in cross-section; chambers decreasing gradually in size; wall calcareous, hyaline, finely perforate; sutures distinct; aperture terminal, radiate.

*Dentalina fissicostata* GÜMBEL, 1868  
Plate 4, Figure 11

1868 *Dentalina fissicostata* n. sp. – GÜMBEL, p. 48, pl. 1, fig. 46.  
1966 *Dentalina fissicostata* GÜMBEL – HOFKER, p. 217, pl. 41, fig. 21.

Description: Test elongate, monoserial, straight, circular in cross-section; large-size, inflated chambers increasing gradually in size; wall calcareous, hyaline, finely perforate with numerous strong, longitudinal ribs; sutures strongly depressed; aperture terminal, radiate.

*Dentalina inornata* D'ORBIGNY, 1846  
Plate 4, Figure 13

1846 *Dentalina inornata* n. sp. – D'ORBIGNY, p. 44, pl. 1, figs. 50–51.  
1961 *Dentalina inornata* D'ORBIGNY – KAASSCHIETER, p. 176, pl. 7, figs. 18–19.  
1977 *Dentalina inornata* D'ORBIGNY – POŻARYSKA, p. 23, pl. 2, fig. 25.

Description: Test elongate, monoserial, slightly curved, circular in cross-section; chambers increasing gradually in size; wall calcareous, hyaline, finely perforate; sutures distinct; aperture terminal, radiate.

*Dentalina multilineata* BORNEMANN, 1855  
Plate 4, Figure 14

1855 *Dentalina multilineata* n. sp. – BORNEMANN, p. 325, pl. 13, fig. 12.  
1932 *Dentalina multilineata* BORNEMANN – NUTTALL, pl. 3, fig. 5.  
1987 *Dentalina multilineata* BORNEMANN – SZTRÁKOS, pl. 4, fig. 14.

Description: Test elongate, monoserial, slightly curved, circular in cross-section; chambers increasing gradually in size; wall calcareous, hyaline, finely perforate with numerous fine, longitudinal ribs; sutures distinct; aperture terminal, radiate.

*Dentalina karreri* (HANTKEN, 1868)  
Plate 4, Figure 15

1868 *Nodosaria karreri* n. sp. – HANTKEN, p.85, pl. 1, fig. 8.

Description: Test elongate, monoserial, straight, circular in cross-section; chambers increasing gradually in size, final chamber inflated; wall calcareous, hyaline, finely perforate; sutures distinct; aperture terminal, radiate.

*Dentalina havanensis* CUSHMAN and BERMÚDEZ, 1937  
Plate 4, Figure 16

1937 *Dentalina havanensis* n. sp. – CUSHMAN and BERMÚDEZ, p. 11, pl. 1, figs. 39–40.

Description: Test elongate, monoserial, slightly curved, circular in cross-section; chambers increasing gradually in size; wall calcareous, hyaline, finely perforate; sutures invisible; aperture terminal, radiate.

*Dentalina roemeri* NEUGEBOREN, 1856  
Plate 4, Figure 17



- 1856 *Dentalina roemeri* n. sp. – NEUGEBOREN, p. 82, pl. 2, figs. 13–17.  
1949 *Dentalina roemeri* NEUGEBOREN – CUVILLIER and SZAKÁLL, p. 79, pl. 28. fig. 31.

Description: Test elongate, monoserial, slightly curved, circular in cross-section; chambers increasing gradually in size; wall calcareous, hyaline, finely perforate; sutures slightly depressed; aperture terminal, radiate.

*Dentalina subtilis* NEUGEBOREN, 1856  
Plate 4, Figure 18

- 1856 *Dentalina subtilis* n. sp. – NEUGEBOREN, p. 9, pl. 3, fig. 4.  
1875 *Dentalina subtilis* NEUGEBOREN – HANTKEN, p. 28, pl. 3, fig. 13.

Description: Test elongate, monoserial, slightly curved, circular in cross-section; chambers increasing gradually in size; wall calcareous, hyaline, finely perforate; sutures barely visible; aperture terminal, radiate.

Genus *Nodosaria* LAMARCK, 1812

Type species: *Nodosaria radricula* (LINNÉ, 1758)

*Nodosaria acuminata* HANTKEN, 1875  
Plate 4, Figures 25–26

- 1875 *Nodosaria acuminata* n. sp. – HANTKEN, p. 28, pl. 2, fig. 9, pl. 13, fig. 5.  
1975 *Nodosaria acuminata* HANTKEN – SAMUEL, p. 119, pl. 70, fig. 3.  
1985 *Nodosaria acuminata* HANTKEN – KORECZNÉ LAKY and NAGYNÉGELLAI, pl. 10, fig. 14.

Description: Test elongate, monoserial, straight, circular in cross-section; chambers increasing gradually in size, final chamber acute; wall calcareous, hyaline, finely perforate with numerous strong, longitudinal ribs; aperture terminal, central, circular.

*Nodosaria affinis* REUSS, 1845  
Plate 4, Figure 19

- 1845 *Nodosaria affinis* n. sp. – REUSS, p. 26, pl. 13, fig. 16.  
1968 *Nodosaria affinis* REUSS – HOFKER, p. 179, pl. 34, figs. 88–89.

Description: Test elongate, monoserial, straight, circular in cross-section; first chamber inflated; wall calcareous, hyaline, finely perforate with numerous strong, longitudinal ribs; aperture terminal, central, circular.

*Nodosaria badenensis* (D'ORBIGNY, 1846)  
Plate 4, Figure 20

- 1846 *Nodosaria spinicosta* n. sp. – D'ORBIGNY, p. 37, pl. 1, figs. 32–33.  
1985 *Nodosaria badenensis* (D'ORBIGNY) – PAPP and SCHMID, p. 26, pl. 7, figs. 1–3.  
1985 *Nodosaria badenensis* (D'ORBIGNY) – KORECZNÉ-LAKY and NAGYNÉ-GELLAI, pl. 99, fig. 11.

Description: Test elongate, monoserial, straight, circular in cross-section; chambers increasing gradually in size, final chamber inflated; wall calcareous, hyaline, finely perforate with numerous strong, longitudinal ribs; aperture terminal, central, circular.

Remarks: *Nodosaria badenensis* (D'ORBIGNY) has been reported from the Miocene of Vienna Basin, from the Lower Miocene of France and Hungary.

*Nodosaria crassa* HANTKEN, 1868  
Plate 4, Figure 21

- 1868 *Nodosaria crassa* n. sp. – HANTKEN, p. 52, pl. 1, fig. 15.

- 1875 *Nodosaria crassa* HANTKEN – HANTKEN, p. 23, pl. 13, fig. 4.  
1985 *Nodosaria crassa* HANTKEN – KORECZNÉ-LAKY and NAGYNÉ-GELLAI, pl. 35, fig. 3.

Description: Test squattish, monoserial, straight, circular in cross-section; two or three chambers, wall calcareous, hyaline, finely perforate with numerous strong, longitudinal ribs; aperture terminal, central, circular.

*Nodosaria elegans* (HANTKEN, 1875)  
Plate 4, Figure 22

- 1875 *Dentalina elegans* D'ORBIGNY – HANTKEN, p. 25, pl. 3, fig. 7.

Description: Test elongate, monoserial, slightly curved, circular in cross-section; chambers increasing gradually in size, wall calcareous, smooth, hyaline, finely perforate; aperture terminal, central, circular.

*Nodosaria exilis* NEUGEBOREN, 1852  
Plate 4, Figure 23

- 1852 *Nodosaria exilis* n. sp. – NEUGEBOREN, p. 51, pl. 1, figs. 25–26.  
1975 *Nodosaria exilis* NEUGEBOREN – SAMUEL, p. 120, pl. 69, figs. 12a–b, 13–14.  
1985 *Nodosaria exilis* NEUGEBOREN – KORECZNÉ-LAKY – NAGYNÉ-GELLAI, pl. 9, fig. 14.

Description: Test elongate, monoserial, straight, circular in cross-section; first chamber inflated, spherical; wall calcareous, hyaline, finely perforate; aperture terminal, central, circular.

*Nodosaria intermedia* (HANTKEN, 1875)  
Plate 4, Figure 24

- 1875 *Dentalina intermedia* n. sp. – HANTKEN, p. 25, pl. 3, figs. 4, 8.

Description: Test elongate, monoserial, slightly curved, circular in cross-section; chambers increasing gradually in size, final three chambers slightly inflated; wall calcareous, hyaline, smooth, finely perforate; aperture terminal, central, circular.

*Nodosaria longiscata* D'ORBIGNY, 1846  
Plate 4, Figure 27

- 1846 *Nodosaria longiscata* n. sp. – D'ORBIGNY, p. 32, pl. 1, figs. 10–12.  
1952 *Nodosaria longiscata* D'ORBIGNY – TODD and KNIKER, p. 16, pl. 3, figs. 9–10.  
1985 *Nodosaria longiscata* D'ORBIGNY – PAPP and SCHMID, p. 23, pl. 3, figs. 1–5.  
1985 *Nodosaria longiscata* D'ORBIGNY – KORECZNÉ-LAKY and NAGYNÉ-GELLAI, pl. 99, fig. 1.

Description: Test elongate, monoserial, straight, circular in cross-section; first chamber inflated, chambers increasing gradually in size; wall calcareous, hyaline, finely perforate with numerous strong, longitudinal ribs; invisible sutures; aperture terminal, central, circular.

*Nodosaria pyrula* D'ORBIGNY, 1826  
Plate 4, Figure 28.

- 1826 *Nodosaria pyrula* n. sp. – D'ORBIGNY, p. 253, pl. 9, fig. 37.  
1846 *Nodosaria mariae* n. sp. – D'ORBIGNY, p. 33, pl. 1, figs. 15–16.  
1985 *Nodosaria pyrula* D'ORBIGNY – PAPP and SCHMID, p. 24, pl. 4, figs. 2–3.  
1985 *Nodosaria pyrula* D'ORBIGNY – KORECZNÉ-LAKY and NAGYNÉ-GELLAI, pl. 9, figs. 12–13; pl. 35, fig. 5; pl. 99, fig. 21.

Description: Test elongate, monoserial, straight, circular in cross-section; chambers increasing gradually in size, final chamber inflated; wall calcareous, smooth, hyaline, finely perforate; strongly depressed sutures; aperture terminal on long neck with thickened lip.

*Nodosaria radricula* (LINNÉ, 1758)

Plate 4, Figure 29

1758 *Nautilus radricula* n. sp. – LINNÉ, p. 711, pl. 1, figs. 5a–c (fide ELLIS and MESSINA).

1982 *Nodosaria radricula* LINNÉ – SZTRÁKOS, pl. 5, fig. 19.

1985 *Nodosaria radricula* LINNÉ – KORECZNÉ-LAKY and NAGYNÉ-GELLAI, pl. 14, figs. 3–4; pl. 103, fig. 3.

Description: Test squattish, monoserial, straight, circular in cross-section; chambers increasing gradually in size, final chamber inflated; wall calcareous, smooth, hyaline; aperture terminal, central, circular.

Genus *Pseudonodosaria* BOOMGAART, 1949

Type species: *Pseudonodosaria discreta* (REUSS, 1850)

*Pseudonodosaria discreta* (REUSS, 1850)

Plate 4, Figure 30

1850 *Glandulina discreta* n. sp. – REUSS, p. 372, pl. 47, fig. 9.

1987 *Pseudonodosaria discreta* (REUSS) – SZTRÁKOS, pl. 5, fig. 28.

Description: Test squattish, monoserial, straight, circular in cross-section; chambers increasing gradually in size, final chamber inflated; wall calcareous, hyaline, finely perforate smooth; aperture terminal, circular.

Subfamily Lingulininae LOEBLICH and TAPPAN, 1961

Genus *Gonatosphaera* GUPPY, 1894

Type species: *Gonatosphaera prolata* Guppy, 1894

*Gonatosphaera inflata* BERMÚDEZ, 1949

Plate 4, Figures 31–32

1949 *Gonatosphaera inflata* n. sp. – BERMÚDEZ, p. 232, pl. 14, figs. 50–51.

1979 *Gonatosphaera inflata* BERMÚDEZ – SZTRÁKOS, pl. 16, fig. 3.

Description: Test squattish, monoserial, straight, circular in cross-section; chambers inflated and increasing gradually in size, final chamber strongly inflated; wall calcareous, hyaline, finely perforate smooth; aperture terminal, central, circular.

Genus *Frondicularia* DEFRANCE emend. NORLING, 1972

Type species: *Frondicularia complanata* (DEFRANCE, 1824)

*Frondicularia budensis* (HANTKEN, 1875)

Plate 5, Figure 2

1875 *Flabellina budensis* n. sp. – HANTKEN, p. 37, pl. 4, fig. 17.

1949 *Frondicularia budensis* (HANTKEN) – CUVILLIER and SZAKÁLL, p. 84, pl. 30, fig. 14.

1953 *Frondicularia budensis* (HANTKEN) – SUBBOTINA, p. 199, pl. 7, figs. 6–7.

1979 *Palmula budensis* (HANTKEN) – SZTRÁKOS, pl. 33, figs. 9–10.

1985 *Frondicularia budensis* (HANTKEN) – GRÜNIG, p. 261, pl. 4, fig. 21.

1985 *Frondicularia budensis* (HANTKEN) – KORECZNÉ-LAKY and NAGYNÉ-GELLAI, pl. 11, fig. 24; pl. 25, figs. 1–2.

1992 *Palmula budensis* (HANTKEN) – DARAKCHIEVA and JURANOV, p. 12, pl. 1, figs. 9–10.

1993 *Palmula budensis* (HANTKEN) – SZTRÁKOS, p. 76, pl. 9, fig. 10.

2006 *Palmula budensis* (HANTKEN) – CIMERMAN et al., p. 24, pl. 5, fig. 7.

Description: Test monoserial, lanceolate in outline, strongly compressed in cross-section; length approximately two times width breadth; chambers increasing gradually in size, wall calcareous, finely perforate with numerous V-shaped elevated sutures; terminal aperture.

*Frondicularia semicosta* KARRER, 1877  
Plate 5, Figure 1

- 1877 *Frondicularia semicosta* n. sp. – KARRER, p. 380, pl. 16b, fig. 26.  
1954 *Plectofrondicularia semicosta* (KARRER) – COLOM, p. 260, pl. 29, figs. 24–25.  
1982 *Plectofrondicularia semicosta* (KARRER) – AGIP, pl. 24, fig. 6.

Description: Test monoserial, lanceolate in outline, strongly compressed in cross-section; length approximately three and a half times larger than width; chambers increasing gradually in size, wall calcareous, finely perforate with numerous fine, longitudinal ribs, aperture terminal.

*Frondicularia tenuissima* HANTKEN, 1875  
Plate 5, Figures 3–4

- 1875 *Frondicularia tenuissima* n. sp. – HANTKEN, p. 43, pl. 13, fig. 11.  
1927 *Frondicularia tenuissima* HANTKEN – CUSHMAN, p. 111, pl. 22, fig. 11.  
1949 *Frondicularia tenuissima* HANTKEN – CUVILLIER and SZAKÁLL, p. 85, pl. 30, fig. 15.  
1951 *Frondicularia tenuissima* HANTKEN – FRIESE, p. 27, pl. 11, fig. 22.  
1952 *Flabellinella tenuissima* (HANTKEN) – HAGN, p. 156, pl. 4, fig. 20.  
1979 *Frondicularia tenuissima* HANTKEN – SZTRÁKOS, pl. 12, fig. 1.  
1982 *Plectofrondicularia tenuissima* (HANTKEN) – AGIP, pl. 24, fig. 8.  
1985 *Frondicularia tenuissima* HANTKEN – KORECZNÉ-LAKY and NAGYNÉ-GELLAI, pl. 11, fig. 25; pl. 25, figs. 3–4.  
1987 *Flabellinella tenuissima* (HANTKEN) – REISER, p. 72, pl. 5, figs. 29–30.  
1993 *Frondivaginulina tenuissima* (HANTKEN) – MATHLEIN and SZTRÁKOS, p. 76, pl. 9, fig. 8.  
2006 *Palmula tenuissima* (HANTKEN) – CIMERMAN et al., p. 24, pl. 5, figs. 8–9.

Description: Test monoserial, lanceolate in outline, strongly compressed in cross-section; length approximately three times larger than width; wall calcareous, finely perforate with numerous fine, longitudinal ribs; terminal aperture.

Family Vaginulinidae REUSS, 1860  
Subfamily Lenticulininae CHAPMAN, PARR and COLLINS, 1934

Genus *Lenticulina* LAMARCK, 1804

Type species: *Lenticulina rotula* (LAMARCK, 1804)

*Lenticulina arcuatostrata* (HANTKEN, 1868)  
Plate 5, Figures 5–6

- 1868 *Cristellaria (Robulina) arcuato striata* n. sp. – HANTKEN, p. 93, pl. 2, fig. 30.  
1875 *Robulina arcuatostrata* (HANTKEN) – HANTKEN, p. 48, pl. 7, fig. 2.  
1949 *Robulus arcuatostratus* (HANTKEN) – CUVILLIER and SZAKÁLL, p. 51, pl. 23, fig. 9.  
1950 *Robulus arcuatostratus* (HANTKEN) – RUIZ DE GAONA and COLOM, p. 402, pl. 6, figs. 1–9.  
1956 *Robulus arcuatostratus* (HANTKEN) – HAGN, p. 127, pl. 11, fig. 4.  
1982 *Lenticulina arcuatostrata* (HANTKEN) – SZTRÁKOS, pl. 9, fig. 2.  
1985 *Lenticulina arcuatostrata* (HANTKEN) – GRÜNIG, p. 261, pl. 4, fig. 19.  
1985 *Lenticulina arcuatostrata* (HANTKEN) – KORECZNÉ-LAKY and NAGYNÉ-GELLAI, pl. 12, fig. 1.  
2006 *Lenticulina arcuatostrata* (HANTKEN) – CIMERMAN et al., p. 22, pl. 4, fig. 13.

Description: Test planispiral, involute, circular in outline, biconvex in cross-section; chambers increasing moderately in size; sutures distinct, slightly curved, wall calcareous, smooth, finely perforate; aperture terminal, radiate.

*Lenticulina excisa* (BORNEMANN, 1855)

Plate 5, Figure 7

1855 *Cristellaria excisa* n. sp. – BORNEMANN, p. 328, pl. 13, figs. 19–20.

1979 *Lenticulina excisa* (BORNEMANN) – SZTRÁKOS, pl. 12, fig. 11.

Description: Test planispiral, involute, oval in outline, biconvex in cross-section; periphery acute; chambers increasing moderately in size; sutures distinct, slightly curved; wall calcareous, smooth, finely perforate; aperture terminal, radiate.

*Lenticulina falcifer* (STACHE, 1865)

Plate 5, Figure 8

1865 *Cristellaria* (*Cristellaria*) *falcifer* n. sp. – STACHE, p. 240, pl. 23, figs. 19a–b.

1979 *Lenticulina falcifer* (STACHE) – SZTRÁKOS, pl. 12, fig. 12.

Description: Test planispiral, involute, circular in outline, biconvex in cross-section; chambers increasing moderately in size; sutures indistinct; wall calcareous, smooth, finely perforate; aperture terminal, radiate.

*Lenticulina granulata* (HANTKEN, 1875).

Plate 5, Figure 9

1875 *Robulina granulata* n. sp. – HANTKEN, p. 49, pl. 14, fig. 15.

1987 *Lenticulina granulata* (HANTKEN) – SZTRÁKOS, pl. 4, fig. 35.

Description: Test planispiral, involute, circular in outline, biconvex in cross-section; periphery acute; chambers increasing moderately in size; sutures indistinct, wall calcareous with many irregularly arranged nodes, finely perforate; aperture terminal, radiate.

*Lenticulina gutticastata* (GÜMBEL, 1868)

Plate 5, Figure 10

1868 *Robulina gutticastata* n. sp. – GÜMBEL, p. 643, pl. 1, figs. 74a–b.

1875 *Robulina gutticastatus* (GÜMBEL) – HANTKEN, p. 48, pl. 6, fig. 10.

1985 *Lenticulina cassis* (FICHEL and MOLL) – GRÜNIG, p. 261, pl. 4, fig. 20.

Description: Test planispiral, involute, circular in outline, biconvex in cross-section; chambers increasing moderately in size; sutures distinct with small oval to circular beads; wall calcareous, smooth, finely perforate; aperture terminal, radiate.

*Lenticulina platyptera* (REUSS, 1870)

Plate 5, Figure 11

1870 *Cristellaria* (*Robulina*) *platyptera* n. sp. – REUSS, p. 482, pl. 19, figs. 7–8.

1987 *Lenticulina platyptera* (REUSS) – SZTRÁKOS, pl. 4, fig. 37.

Description: Test planispiral, involute, circular in outline, biconvex in cross-section; periphery acute; chambers increasing moderately in size; sutures distinct, straight to slightly curved; wall calcareous, smooth, finely perforate; aperture terminal, radiate.

*Lenticulina* sp.

Plate 5, Figure 12

Description: Test planispiral, involute, dropshape in outline, biconvex in cross-section; chambers increasing moderately in size; sutures indistinct; wall calcareous, smooth, finely perforate; aperture terminal, radiate.

Genus *Marginulinopsis* SILVESTRI, 1904

Type species: *Marginulinopsis densicostata* THALMANN, 1937

*Marginulinopsis porvaensis* (HANTKEN, 1875)

Plate 5, Figure 13

- 1875 *Cristellaria porvaensis* n. sp. – HANTKEN, p. 42, pl. 14, fig. 1.  
1950 *Marginulina porvaensis* (HANTKEN) – RUIZ DE GAONA and COLOM, p. 387, pl. 11, figs. 1–14.  
1987 *Marginulinopsis porvaensis* (HANTKEN) – SZTRÁKOS, pl. 5, fig. 20.  
1993 *Astaculus porvaensis* (HANTKEN) – MATHELIN and SZTRÁKOS, p. 76, pl. 26, fig. 13.  
2006 *Marginulinopsis porvaensis* (HANTKEN) – CIMERMAN et al., p. 24, pl. 5, figs. 2–3.

Description: Test planispiral, involute in earlier three – four chambers, becoming monoserial later (five to seven chambers, oval in cross-section; length approximately four times width breadth; chambers increasing gradually in size; sutures distinct, straight, slightly elevated; wall calcareous, smooth, finely perforated; aperture terminal, circular on short neck.

*Marginulinopsis* sp.

Plate 5, Figure 14

Description: Test planispiral, involute in earlier three – four chambers, becoming monoserial later, oval in cross-section; length approximately four times width breadth; chambers increasing gradually in size; sutures distinct, straight, slightly elevated; wall calcareous, smooth, finely perforated with numerous fine, longitudinal ribs on earlier chambers; aperture terminal, circular on short neck.

Genus *Saracenaria* DEFRANCE, 1824

Type species: *Saracenaria italica* DEFRANCE, 1824

*Saracenaria hantkeni* CUSHMAN, 1933

Plate 5, Figures 15–16

- 1933 *Saracenaria arcuata* D'ORBIGNY var. *hantkeni* – CUSHMAN, p. 4, pl. 1, figs. 11–12.  
1953 *Saracenaria hantkeni* CUSHMAN – BECKMANN, p. 353, pl. 19, fig. 18.  
1956 *Saracenaria hantkeni* CUSHMAN – HAGN, p. 138, pl. 13, fig. 4.  
1977 *Saracenaria arcuata* (D'ORBIGNY) – POŻARYSKA, p. 25, pl. 2, figs. 22a–b.  
1987 *Saracenaria hantkeni* CUSHMAN – SZTRÁKOS, pl. 6, fig. 3.  
1993 *Saracenaria hantkeni* CUSHMAN – MATHELIN and SZTRÁKOS, p. 76, pl. 9, fig. 6.  
2006 *Saracenaria hantkeni* CUSHMAN – CIMERMAN et al., p. 24, pl. 5, fig. 6.

Description: Test planispiral in earlier chambers, involute, becoming monoserial in final two or three chambers; dropshape in outline, rounded triangular in cross-section; chambers increasing gradually in size, final chambers strongly inflated; wall calcareous, smooth, finely perforated; sutures invisible; aperture terminal, radiate.

Subfamily Marginulininae WEDEKIND, 1937

Genus *Marginulina* D'ORBIGNY, 1826

Type species: *Marginulina raphanus* LINNÉ, 1758

*Marginulina behmi* (REUSS, 1866)

Plate 5, Figure 17

- 1866 *Cristellaria behmi* n. sp. – REUSS, p. 138, pl. 2, fig. 37.  
1868 *Cristellaria behmi* REUSS – GÜMBEL, p. 55, pl. 1, fig. 61.  
1875 *Marginulina behmi* (REUSS) – HANTKEN, p. 48, pl. 5, figs. 1–2.

- 1947 *Marginulina behmi* (REUSS) – SAMOILOVA, p. 82, pl. 3, fig. 9.  
 1949 *Marginulina behmi* (REUSS) – CUVILLIER and SZAKÁLL, p. 70, pl. 27, fig. 7.  
 1953 *Marginulina behmi* (REUSS) – SUBBOTINA, p. 65, pl. 4, figs. 4–5.  
 1956 *Marginulina behmi* (REUSS) – HAGN, p. 131, pl. 11, fig. 11.  
 1950 *Marginulina behmi* (REUSS) – RUIZ DE GAONA and COLOM, p. 384, pl. 11, figs. 18–26.  
 1956 *Marginulina behmi* (REUSS) – HAGN, p. 131, pl. 11, fig. 11.  
 1965 *Marginulina behmi* (REUSS) – KRAYEVA, p. 4, pl. 21, figs. 1–2.  
 1969 *Marginulina behmi* (REUSS) – KRAYEVA and ZARNECKIJ, p. 60, pl. 21, figs. 5–6.  
 1975 *Marginulina behmi* (REUSS) – BRAGA et al., p. 105, pl. 4, fig. 13.  
 1975 *Marginulina behmi* (REUSS) – SAMUEL, p. 126, pl. 70, fig. 11.  
 1977 *Marginulina behmi* (REUSS) – POŻARYSKA, p. 24, pl. 2, figs. 27a–b; pl. 3, figs. 1–9; pl. 8, figs. 1a–c.  
 1982 *Marginulina behmi* (REUSS) – AGIP, pl. 11, figs. 10–10p.  
 1985 *Marginulina behmi* (REUSS) – GRÜNIG, p. 262, pl. 5, figs. 17–19.  
 1985 *Marginulina behmi* (REUSS) – KORECZNÉ-LAKY and NAGYNÉ-GELLAI, pl. 13, fig. 1.  
 1992 *Marginulina behmi* (REUSS) – DARAKCHIEVA and JURANOV, p. 13, pl. 2, figs. 4–5.  
 1993 *Marginulina behmi* (REUSS) – MATHELIN and SZTRÁKOS, p. 43, pl. 26, figs. 14–16.  
 2006 *Marginulinopsis behmi* (REUSS) – CIMERMAN et al., p. 24, pl. 5, fig. 1.

Description: Test elongate, planispiral, involute in earlier chambers, becoming monoserial later (four or five chambers, oval in cross-section; chambers increasing gradually in size; sutures distinct, slightly depressed; wall calcareous, smooth, finely perforated with numerous strong, longitudinal ribs; aperture terminal, circular with thickened lip on short neck.

*Marginulina fragaria texasensis* (CUSHMAN and APPLIN, 1926)

Plate 5, Figure 18

- 1926 *Cristellaria fragaria* (GÜMBEL) var. *texasensis* n. subsp. – CUSHMAN and APPLIN, p. 171, pl. 8, figs. 5–7.  
 1932 *Lenticulina fragaria* (GÜMBEL) var. *texasensis* CUSHMAN and APPLIN – HOWE and WALLACE, p. 32, pl. 5, figs. 3–5.  
 1933 *Marginulina fragaria* (GÜMBEL) var. *texasensis* CUSHMAN and APPLIN – ELLISOR, pl. 2, fig. 4.  
 1935 *Marginulina fragaria* (GÜMBEL) var. *texasensis* CUSHMAN and APPLIN – CUSHMAN, p. 19, pl. 7, figs. 8a–b, 10a–b (non figs. 9a–b).

Description: Test elongate, planispiral, involute in earlier chambers, becoming monoserial later (four or five chambers, oval in cross-section; chambers increasing gradually in size, final chamber smaller than previous; sutures distinct, slightly depressed; wall calcareous, smooth, finely perforated with numerous horizontal beads; aperture terminal, circular with thickened lip on short neck.

*Marginulina hantkeni* BANDY, 1949

Plate 5, Figures 19–20

- 1875 *Marginulina subbullata* n. sp. – HANTKEN, p. 46, pl. 4, figs. 9–10; pl. 5, fig. 9.  
 1949 *Marginulina hantkeni* nomen novum – BANDY, p. 46, pl. 6, fig. 9.  
 1975 *Marginulina hantkeni* BANDY – POPESCU, p. 56, pl. 20, fig. 4; pl. 21, fig. 8.  
 1982 *Marginulina subbullata* HANTKEN – AGIP, pl. 13, fig. 1.  
 1982 *Marginulina hantkeni* BANDY – SZTRÁKOS, pl. 11, fig. 6.  
 1987 *Marginulina hantkeni* BANDY – SZTRÁKOS, pl. 5, fig. 14.

Description: Test squattish, planispiral in earlier chambers, becoming monoserial in final two or three chambers; oval in outline, circular in cross-section; chambers increasing gradually in size, final chambers strongly inflated; wall calcareous, smooth, finely perforated; sutures distinct, strongly depressed; aperture terminal, radiate on short neck.

*Marginulina pediformis* BORNEMANN, 1855

Plate 5, Figure 21

- 1855 *Marginulina pediformis* n. sp. – BORNEMANN, p. 326, pl. 13, fig. 13.  
 1949 *Marginulina pediformis* BORNEMANN – CUVILLIER and SZAKÁLL, p. 70, pl. 27, fig. 5.  
 1987 *Marginulina pediformis* BORNEMANN – SZTRÁKOS, pl. 5, fig. 13.

Description: Test elongate, slightly curved, planispiral in earlier chambers, becoming monoserial in final three or four chambers; chambers mildly inflated and increasing gradually in size; sutures distinct, strongly depressed; wall calcareous, smooth, finely perforated

*Marginulina propinqua* HANTKEN, 1883  
Plate 5, Figure 22

1883 *Marginulina propinqua* n. sp. – HANTKEN, p. 26, pl. 2, figs. 4a–b.  
1987 *Marginulina propinqua* HANTKEN – SZTRÁKOS, pl. 5, fig. 16.

Description: Test squattish, planispiral, involute in earlier chambers, becoming monoserial later, oval in cross-section; chambers increasing gradually in size; sutures barely visible; wall calcareous, smooth, finely perforated with numerous fine, longitudinal ribs on planispiral chambers; aperture terminal, circular on short neck.

*Marginulina tumida* REUSS, 1851  
Plate 5, Figure 23

1851 *Marginulina tumida* n. sp. – REUSS, p. 64, pl. 3, figs. 14a–b.  
1979 *Marginulina tumida* REUSS – SZTRÁKOS, pl. 14, fig. 3.

Description: Test mildly elongate, planispiral, involute in earlier chambers, becoming monoserial later, oval in cross-section; chambers increasing gradually in size; sutures barely visible; wall calcareous, smooth, finely perforated; aperture terminal, radiate.

*Marginulina* sp. 1.  
Plate 5, Figure 24

Description: Test squattish, planispiral, involute in earlier chambers, becoming monoserial later, oval to circular in cross-section; chambers increasing gradually in size; sutures barely visible; wall calcareous, smooth, finely perforated with numerous strong longitudinal ribs; aperture terminal, radiate.

Genus *Vaginulinopsis* SILVESTRI, 1904

Type species: *Vaginulinopsis inversa carinata* SILVESTRI, 1904

*Vaginulinopsis hagni* n. sp.  
Plate 5, Figures 25–27

1956 *Vaginulinopsis cumulicostata* (GÜMBEL) – HAGN, p. 135, pl. 13, fig. 3.  
1975 *Vaginulinopsis cumulicostata* (GÜMBEL) – BRAGA et al, p. 105, pl. 4, fig. 8.  
1985 *Vaginulinopsis cumulicostata* (GÜMBEL) – GRÜNIG, p. 263, pl. 6, fig. 6.  
1987 *Vaginulinopsis cumulicostata* (GÜMBEL) – SZTRÁKOS, pl. 6, fig. 12.  
1993 *Vaginulinopsis cumulicostatus* (GÜMBEL) – MATHELIN and SZTRÁKOS, p. 76, pl. 10, fig. 12.  
2006 *Vaginulinopsis cumulicostata* (GÜMBEL) – CIMERMAN et al., p. 26, pl. 5, figs. 13–14.

Derivatio nominis: In honour of HAGN, H. for his outstanding work on Tertiary benthic foraminifera

Holotype: The specimen on Plate 5, Figure 26. (Inventory number: M 2008.118.1)

Dimension of holotype: length 0.7 mm, width 0.34 mm.

Description: Test elongate, planispiral, evolute in earlier chambers, becoming monoserial later (four or five chambers); oval in cross-section; chambers increasing gradually in size; sutures distinct, slightly depressed; wall calcareous, smooth, finely perforated with three or four horizontal ribs; aperture terminal, circular on short neck

Remarks: GÜMBEL (1868) described the species *Cristellaria cumulicostata* GÜMBEL, 1868 from Upper Eocene of Northern Calcareous Alps. An absolutely different specimen was assigned by HAGN (1956) to *Vaginulinopsis cumulicostata* (GÜMBEL) as the synonym of *Cristellaria cumulicostata*



GÜMBEL, 1868 from Tertiary of Monte Brione, Italy. This form is not related to *Vaginulinopsis cumulicostata* (GÜMBEL) which has a broad planispiral portion and long monoserial part with numerous horizontal beads on sutures. Many authors used the name *Vaginulinopsis cumulicostata* (GÜMBEL) refer to HAGN (1956) none the less the *Vaginulinopsis hagni* n. sp. has significantly different shape and different ribs.

Type locality: In borehole Devecser 4.

Type strata: In borehole Devecser 4 borehole, at 81.0 m. Padrag Marl Formation (Middle Eocene, Bartonian)

*Vaginulinopsis hantkeni* (HANTKEN, 1875)

Plate 5 Figure 28

1875 *Cristellaria ornata* n. sp. – HANTKEN, p. 77, pl. 13, fig. 19.

Description: Test squattish, planispiral, involute in earlier chambers, becoming monoserial later; corcular to oval in cross-section; chambers increasing gradually in size; sutures barely visible; wall calcareous, smooth, finely perforated; aperture terminal, circular on short neck.

*Vaginulinopsis fragaria* (GÜMBEL, 1868)

Plate 6, Figure 1

- 1868 *Marginulina fragaria* n. sp. – GÜMBEL, p. 57, pl. 1, figs. 58a–c.  
1875 *Cristellaria fragaria* (GÜMBEL) – HANTKEN, p. 53, pl. 6, figs. 1–3.  
1947 *Lenticulina fragaria* (GÜMBEL) – SUBBOTINA, p. 85, pl. 8, figs. 6–7.  
1953 *Marginulina fragaria* GÜMBEL – SUBBOTINA, p. 162, pl. 4, figs. 1–3.  
1949 *Marginulina fragaria* (GÜMBEL) – CUVILLIER and SZAKÁLL, p. 69, pl. 26, figs. 15, 17–18.  
1950 *Marginulina fragaria* (GÜMBEL) – RUIZ DE GAONA and COLOM, p. 386, pl. 11, figs. 15–17.  
1956 *Marginulinopsis fragaria* (GÜMBEL) – HAGN, p. 134, pl. 12, figs. 3, 8, 10.  
1958 *Marginulina fragaria* (GÜMBEL) – NYIRÓ, p. 34, fig. 8.  
1975 *Marginulina fragaria* (GÜMBEL) – PROTO DECIMA and DE BIASE, p. 93, pl. 1, fig. 35.  
1982 *Marginulinopsis fragaria* (GÜMBEL) – AGIP, pl. 12, figs. 6–6p.  
1982 *Marginulinopsis fragaria* (GÜMBEL) – SZTRÁKOS, pl. 12, fig. 5.  
1985 *Marginulinopsis fragaria* (GÜMBEL) – GRÜNIG, p. 161, pl. 5, figs. 24–28.  
1991 *Marginulinopsis fragaria* (GÜMBEL) – BARBIN and KELLER-GRÜNIG, p. 240, pl. 2, figs. 6–12.  
1992 *Marginulina fragaria* (GÜMBEL) – DARAKCHIEVA and JURANOV, p. 13, pl. 2, fig. 10.  
1993 *Percultazonaria fragaria* (GÜMBEL) – MATHELIN and SZTRÁKOS, p. 76, pl. 9, figs. 1–2.  
2006 *Percultazonaria fragaria* (GÜMBEL) – CIMERMAN et al., p. 24, pl. 5, figs. 4–5.

Description: Test elongate, planispiral, involute in earlier chambers, becoming monoserial later (six or seven chambers); oval in cross-section; chambers increasing gradually in size; sutures distinct, slightly depressed; wall calcareous, smooth, finely perforated with numerous fine longitudinal beads; aperture terminal, circular on short neck.

*Vaginulinopsis minimus* (HANTKEN, 1875)

Plate 5, Figure 29

- 1875 *Cristellaria minima* n. sp. – HANTKEN, p. 77, pl. 13, fig. 21.  
1962 *Marginulina (Marginulinopsis) minima* (HANTKEN) – MAJZON, pl. 41, fig. 21.  
1973 *Marginulina minima* (HANTKEN) – NAGYNÉ GELLAI, p. 453, pl. 3, fig. 12.  
1979 *Astacolus minima* (HANTKEN) – SZTRÁKOS, pl. 10, figs. 6a–b.  
1982 *Astacolus minima* (HANTKEN) – SZTRÁKOS, pl. 6, figs. 10a–b.  
1987 *Astacolus minimus* (HANTKEN) – REISER, p. 66, pl. 4, figs. 3, 8.  
2003 *Saracenaria minima* (HANTKEN) – HORVÁTH, p. 16, pl. 5, fig. 10. (HANTKEN's draw)

Description: Test squattish, planispiral, involute in earlier chambers, becoming monoserial later; mildly compressed in cross-section; chambers increasing gradually in size; sutures barely visible; wall calcareous, smooth, finely perforated; aperture terminal, radiate.

*Vaginulinopsis pseudodecorata* HAGN, 1956

Plate 6, Figure 2

1956 *Vaginulinopsis pseudodecorata* n. sp. – HAGN, p. 83, pl. 6, fig. 5.

1979 *Vaginulinopsis pseudodecorata* HAGN – SZTRÁKOS, pl. 15, fig. 10.

Description: Test elongate, planispiral, involute in earlier chambers, becoming monoserial later; oval in cross-section; chambers increasing gradually in size; sutures distinct; wall calcareous, smooth, finely perforated with numerous horizontal beads; aperture terminal, circular.

*Vaginulinopsis* sp.

Plate 6, Figure 3

Description: Test squattish, planispiral, involute in earlier chambers, becoming monoserial later; oval in cross-section; chambers increasing gradually in size; sutures barely visible; wall calcareous, smooth, finely perforated with curved rib on earlier chambers; aperture terminal, radiate.

#### Subfamily Vaginulininae REUSS, 1860

#### Genus *Planularia* DEFRANCE, 1826

Type species: *Planularia auris* (DEFRANCE), 1824

*Planularia* sp. 1

Plate 6, Figures 4–5

Description: Test small, planispiral, evolute, strongly compressed in cross-section; chambers increasing gradually in size; sutures barely visible; wall calcareous, smooth, finely perforated; aperture terminal, radiate.

*Planularia* sp. 2

Plate 6, Figure 6

Description: Test large, planispiral, evolute in earlier chambers, becoming monoserial in final two or three chambers; strongly compressed in cross-section; periphery with keel; chambers increasing gradually in size, final chamber acute; wall calcareous, smooth, finely perforated; sutures barely visible; aperture terminal, radiate.

#### Genus *Vaginulina* D'ORBIGNY, 1826

Type species: *Vaginulina legumen* (LINNÉ, 1758)

*Vaginulina legumen* (LINNÉ, 1758)

Plate 6, Figures 7–8

1758 *Nautilus legumen* n. sp. – LINNÉ, p. 711, pl. 1, figs. 7g–i (fide ELLIS and MESSINA).

1949 *Vaginulina legumen* (LINNÉ) – CUVILLIER and SZAKÁLL, p. 82, pl. 29, figs. 28–29.

1982 *Vaginulina legumen* (LINNÉ) – AGIP, pl. 15, figs. 7–7p.

1985 *Vaginulina legumen* (LINNÉ) – KORECZNÉ-LAKY and NAGYNÉ-GELLAI, pl. 118, figs. 3–4.

Description: Test elongate, planispiral, involute in earlier chambers, becoming monoserial later (six to eight chambers); oval, strongly compressed in cross-section; length approximately three-four times as long as broad; chambers increasing gradually in size in planispiral portion, later moderately increasing in size; sutures distinct; wall calcareous, smooth, finely perforated with numerous strong horizontal ribs on central portion of each chamber; aperture terminal, radiate.

*Vaginulina* sp. cf. *V. ex gr. mexicana* NUTTALL, 1932

Plate 6, Figure 9

Description: Test elongate, planispiral, involute in earlier chambers, becoming monoserial later (four to six chambers); oval, strongly compressed in cross-section; periphery with keel; length approximately two-three times as long as broad; chambers increasing gradually in size; sutures distinct; wall calcareous, smooth, finely perforated with numerous strong horizontal ribs; aperture terminal, radiate.

Family Lagenidae REUSS, 1862

Genus *Lagena* WALKER and BOYS 1784 emend. SILVESTRI, 1902

Type species: *Lagena sulcata* WALKER and JACOB, 1798

*Lagena globosa* (WALKER and BOYS, 1784)

Plate 6, Figure 10

- 1784 *Serpula (Lagena) laevis globosa* n. sp. – WALKER and BOYS, p. 2, pl. 1, fig. 8. (fide: ELLIS and MESSINA).  
1884 *Oolina globosa* (MONTAGU) – BRADY, p. 452, pl. 56, figs. 1–3.  
1970 *Oolina globosa* (WALKER and BOYS) – LE CALVEZ, p. 101, pl. 17, fig. 1.  
1979 *Oolina globosa* (MONTAGU) – SZTRÁKOS, pl. 18, fig. 6.  
1987 *Oolina globosa* (MONTAGU) – REISER, p. 77, pl. 6, fig. 23.

Description: Test small, squattish, unilocular, circular in outline, circular in cross-section; wall calcareous, hyaline, smooth, finely perforated; aperture terminal, circular on short neck.

*Lagena hexagona* (WILLIAMSON, 1848)

Plate 6, Figure 11

- 1848 *Entosolenia squamosa* var. *hexagona* – WILLIAMSON, p. 20, pl. 2, fig. 23.  
1884 *Lagena hexagona* (WILLIAMSON) – BRADY, p. 472, pl. 58, figs. 32–33.  
1932 *Lagena hexagona* (WILLIAMSON) – HOWE and WALLACE, p. 6, fig. 14.  
1935 *Lagena hexagona* (WILLIAMSON) – CUSHMAN, p. 23, pl. 9, fig. 10.  
1945 *Lagena hexagona* (WILLIAMSON) – CUSHMAN and TODD, p. 33, pl. 5, fig. 14.  
1956 *Lagena hexagona* (WILLIAMSON) – HAGN, p. 141, pl. 10, fig. 22.  
1960 *Oolina hexagona* (WILLIAMSON) – BARKER, p. 120, pl. 58, figs. 32–33.  
1961 *Oolina hexagona* (WILLIAMSON) – ANDERSEN, p. 98, pl. 20, figs. 19a–b.  
1962 *Lagena hexagona* (WILLIAMSON) – KIESEL, p. 41, pl. 6, fig. 18.  
1970 *Oolina hexagona* (WILLIAMSON) – LE CALVEZ, p. 101, pl. 16, figs. 2–3.  
1975 *Lagena hexagona* (WILLIAMSON) – SAMUEL, p. 124, pl. 65, figs. 4 a–d.  
1982 *Lagena hexagona* (WILLIAMSON) – SZTRÁKOS, pl. 8, fig. 15.  
1985 *Oolina hexagona* (WILLIAMSON) – KOHL, p. 57, pl. 16, figs. 16a–c.  
1985 *Lagena hexagona* (WILLIAMSON) – KORECZNÉ-LAKY and NAGYNÉ-GELLAI, pl. 29, figs. 3–4; pl. 122, figs. 1–4.  
1987 *Oolina hexagona* (WILLIAMSON) – SZTRÁKOS, pl. 16, fig. 6.  
1987 *Lagena hexagona* (WILLIAMSON) – WENGER, p. 260, pl. 5, fig. 9.  
1992 *Lagena hexagona* (WILLIAMSON) – DARAKCHIEVA and JURANOV, p. 15, pl. 2, fig. 6.  
1998 *Oolina hexagona* (WILLIAMSON) – ROBERTSON, p. 100, pl. 37, figs. 2a–b.

Description: Test small, squattish, unilocular, circular in outline, circular in cross-section; wall calcareous, hyaline, smooth, finely perforated with hexagonal, slightly elevated frames; aperture terminal.

*Lagena laevis* (WALKER and BOYS, 1784)

Plate 6, Figure 12

- 1784 *Serpula laevis ovalis* n. sp. – WALKER and BOYS, p. 2, pl. 1, fig. 9. (fide: ELLIS and MESSINA).  
1858 *Lagena laevis* (MONTAGU) – WILLIAMSON, p. 12, pl. 1, figs. 1–2.  
1962 *Lagena laevis* (MONTAGU) – KIESEL, p. 42, pl. 6, fig. 14.  
1970 *Lagena laevis* (MONTAGU) – LE CALVEZ, p. 81, pl. 16, fig. 4.  
1982 *Lagena laevis* (MONTAGU) – SZTRÁKOS, pl. 8, figs. 14a–b.

1985 *Lagena laevis* (MONTAGU) – KORECZNÉ-LAKY and NAGYNÉ-GELLAI, pl. 91, fig. 17; pl. 93, fig. 2.

Description: Test elongate, unilocular, oval in outline, circular in cross-section; wall calcareous, hyaline, smooth, finely perforated; aperture terminal, circular on short neck.

*Lagena sulcata* (WALKER and JACOB, 1784)

Plate 6, Figure 14

1784 *Serpula (Lagena) striata sulcata rotunda* n. sp. – WALKER and BOYS, p. 2, pl. 1, fig. 6. (fide: ELLIS and MESSINA).

1798 *Serpula (Lagena) sulcata* n. sp. – WALKER and JACOB in KANMACHER, p. 634, pl. 14, fig. 5.

1839 *Oolina isabella* n. sp. – D'ORBIGNY, p. 21, pl. 5, fig. 1.

1961 *Lagena isabella* (D'ORBIGNY) – KAASSCHIETER, p. 178, pl. 7, fig. 25.

1975 *Lagena isabella* (D'ORBIGNY) – SAMUEL, p. 125, pl. 66, fig. 6.

1985 *Lagena sulcata* (WALKER and JACOB) – KORECZNÉ-LAKY and NAGYNÉ-GELLAI, pl. 11, fig. 11; pl. 91, fig. 18; pl. 121, figs. 1–4.

1992 *Lagena* ex gr. *striata* (D'ORBIGNY) – DARAKCHIEVA and JURANOV, p. 15, pl. 2, fig. 9.

Description: Test small, squattish, unilocular, circular in outline, circular in cross-section; wall calcareous, hyaline, smooth, finely perforated with numerous longitudinal ribs; aperture terminal, circular on short neck.

*Lagena tenuis ornata* REUSS, 1863

Plate 6, Figure 13

1863 *Ovulina tenuis* (BORNEMANN) var. *ornata* n. ssp. – REUSS, 35, pl. 3, figs. 33–39.

1958 *Lagena tenuis* (BORNEMANN) – BATJES, p. 119, pl. 3, fig. 23.

1975 *Lagena tenuis ornata* – SAMUEL, p. 125, pl. 66, figs. 3a–b.

1985 *Lagena tenuis ornata* – KORECZNÉ-LAKY and NAGYNÉ-GELLAI, pl. 31, figs. 1–5.

Description: Test small, squattish, unilocular, drop-shaped in outline, circular in cross-section; wall calcareous, hyaline, smooth, finely perforated; aperture terminal, circular on short neck.

Family Polymorphinidae D'ORBIGNY, 1839

Subfamily Polymorphininae D'ORBIGNY, 1839

Genus *Globulina* D'ORBIGNY, 1839

Type species: *Globulina gibba* (D'ORBIGNY, 1839)

*Globulina guttula* REUSS, 1851

Plate 6, 15

1851 *Globulina guttula* n. sp. – REUSS, p. 82, pl. 6, fig. 46.

1979 *Globulina guttula* REUSS – SZTRÁKOS, pl. 17, fig. 4.

Description: Test slightly elongate, drop-shaped in outline, circular in cross-section; invisible chambers, earlier chambers overlapped by final chamber; length approximately two times maximum width; sutures invisible; wall calcareous, hyaline, smooth, finely perforated; aperture terminal, radiate.

*Globulina minuta* (ROEMER, 1838)

Plate 6, Figure 16

1838 *Polymorphina minuta* n. sp. – ROEMER, p. 386, pl. 3, figs. 35a–b. (fide ELLIS and MESSINA).

1987 *Globulina minuta* (ROEMER) – SZTRÁKOS, pl. 6, fig. 19.

Description: Test squattish, oval in outline, circular in cross-section; invisible chambers, earlier chambers overlapped by final chamber; length approximately two times maximum width; sutures invisible; wall calcareous, hyaline, smooth, finely perforated; aperture terminal, radiate.

Subfamily Ramulininae BRADY, 1884

Genus *Ramulina* JONES, 1875

Type species: *Ramulina laevis* JONES, 1875

*Ramulina* sp.  
Plate 6, Figure 17

Description: Test represented by a single fragment, unilocular, six to eight irregularly arranged branches; wall calcareous, hyaline, smooth, finely perforated.

Family Ellipsolagenidae SILVESTRI, 1923  
Subfamily Ellipsolageninae SILVESTRI, 1923

Genus *Fissurina* REUSS, 1850

Type species: *Fissurina laevigata* REUSS, 1850

*Fissurina orbignyana* SEGUENZA, 1862  
Plate 6, Figure 18

1862 *Fissurina orbignyana* n. sp. – SEGUENZA, p. 66, pl. 2, figs. 25–26.  
1961 *Fissurina orbignyana* SEGUENZA – KAASSCHIETER, p. 180, pl. 7, fig. 29.

Description: Test small, unilocular, oval, mildly compressed in cross-section; periphery with pair of keels, surrounding test and extending to apertural end; wall calcareous, hyaline, smooth, finely perforated; aperture terminal fissure on short neck.

*Fissurina* sp. cf. *F. orbignyana praeclara* (CUSHMAN and RENZ, 1946)  
Plate 6, Figure 19

1946 *Entosolenia orbignyana* var. *praeclara* n. subsp. – CUSHMAN and RENZ, p. 38, pl. 6, fig. 18.

Description: Test small, unilocular, oval, mildly compressed in cross-section; periphery with triple keels, surrounding test and extending to apertural end and with single wide keel surrounding test at aequatorial plane and extending to apertural end; wall calcareous, hyaline, smooth, finely perforated with numerous longitudinal ribs; aperture terminal, circular on long, straight neck.

*Fissurina tricincta* (GÜMBEL, 1868)  
Plate 6, Figures 20–22

1868 *Lagena tricincta* n. sp. – GÜMBEL, p. 606, pl. 1, figs. 8a–b.  
1883 *Lagena scarenaensis* n. sp. – HANTKEN, p. 24, pl. 1, fig. 9.  
1987 *Lagena tricincta* GÜMBEL – SZTRÁKOS, pl. 6, fig. 25.

Description: Test large, unilocular, circular in cross-section; periphery with pair of interior keels, surrounding test and extending to apertural end, with single keel surrounding test at aequatorial plane and extending to apertural end; wall calcareous, hyaline, smooth, finely perforated with numerous circular to oval, densely arranged nodes; aperture terminal, circular on long, straight neck.

*Fissurina* sp.

Plate 6, Figure 23

Description: Test small, unilocular, mildly compressed in cross-section; periphery with pair of interior keels, surrounding test and extending to apertural end, with single keel surrounding test at aequatorial planw and extending to apertural end; wall calcareous, hyaline, smooth, finely perforated with circular to oval, sparsely arranged nodes; aperture terminal, circular on long, straight neck.

Family Glandulinidae REUSS, 1860  
Subfamily Glandulininae REUSS, 1860

Genus *Glandulina* D'ORBIGNY, 1839

Type species: *Glandulina laevigata* D'ORBIGNY, 1826

*Glandulina aequalis* REUSS, 1863  
Plate 6, Figure 24

- 1863 *Glandulina aequalis* n. sp. – REUSS, p. 61, pl. 2, fig. 5.  
1969 *Glandulina aequalis* REUSS – BATJES, p. 123, pl. 4, figs. 5–6.  
non 1975 *Glandulina aequalis* REUSS – SAMUEL, p. 128, pl. 71, fig. 7.  
1987 *Pseudonodosaria aequalis* (REUSS) – SZTRÁKOS, pl. 5., fig. 27.

Description: Test squattish, oval in outline, circular in cross-section, monoserial; length approximately two times maximum width; chambers increasing gradually in size; sutures distinct; aperture terminal, radiate.

*Glandulina hantkeni* (FRANZENAU, 1894)  
Plate 6, Figure 27

- 1894 *Glandulina cuspidata* n. sp. – FRANZENAU, p. 11, pl. 5, fig. 5.  
1987 *Pseudonodosaria hantkeni* (FRANZENAU) – SZTRÁKOS, pl. 6. fig. 2.

Description: Test slightly elongate, monoserial, circular in cross-section; length approximately two times maximum width; chambers increasing gradually in size; strongly inflated final chamber; sutures barely visible; wall calcareous, hyaline, smooth, finely perforated; aperture terminal, radiate.

*Glandulina inflata* COSTA, 1856  
Plate 6, Figures 28

- 1856 *Glandulina inflata* n. sp. – COSTA, p. 126, pl. 11, fig. 21.

Description: Test elongate, monoserial, oval in outline, circular in cross-section; length approximately two times maximum width; mildly inflated chambers; sutures barely visible; wall calcareous, hyaline, smooth, finely perforated; aperture terminal, radiate.

*Glandulina obtusissima* REUSS, 1863  
Plate 6, Figures 25–26

- 1863 *Glandulina obtusissima* n. sp. – REUSS, p. 66, pl. 8, figs. 92–93.  
1949 *Pseudoglandulina obtusissima* REUSS – CUVILLIER and SZAKÁLL, p. 80, pl. 29. fig. 3.

Description: Test squattish, monoserial, circular in cross-section; length approximately two times of maximum width; strongly inflated chambers; sutures distinct; wall calcareous, hyaline, smooth, finely perforated; aperture terminal, radiate.

*Glandulina* sp.  
Plate 6, Figure 29

Description: Test elongate, monoserial, circular in cross-section; length approximately two or three times of maximum width; chambers increasing gradually in size; sutures distinct, strongly depressed; wall calcareous, hyaline, smooth, finely perforated; aperture terminal.

Suborder Rotaliina DELAGE and HÉROUARD, 1879  
Superfamily Bolivinaea GLAESSNER, 1937  
Family Bolivinidae GLAESSNER, 1937

Genus *Bolivina* D'ORBIGNY, 1839

Type species: *Bolivina plicata* D'ORBIGNY, 1839

*Bolivina cookei* CUSHMAN, 1922  
Plate 7, Figure 1

- 1922 *Bolivina cookei* n. sp. – CUSHMAN, p. 126, pl. 29, fig. 1.  
1961 *Bolivina cookei* CUSHMAN – KAASSCHIETER, p. 195, pl. 8, figs. 25–26.  
1962 *Bolivina cookei* CUSHMAN – KIESEL, p. 62, pl. 9, fig. 10.  
1985 *Bolivina cookei* CUSHMAN – KORECZNÉ-LAKY and NAGYNÉ-GELLAI, pl. 49, figs. 1–4.

Description: Test elongate, biserial, lanceolate in outline, oval, strongly compressed in cross-section; length approximately three to four times of maximum width; chambers increasing gradually in size; sutures invisible; wall calcareous, hyaline, smooth, finely perforated with numerous longitudinal ribs; aperture terminal fissure.

*Bolivina elongata* HANTKEN, 1875  
Plate 7, Figures 2–3

- 1875 *Bolivina elongata* n. sp. – HANTKEN, p. 55, pl. 7, fig. 14.  
1979 *Bolivina elongata* HANTKEN – SZTRÁKOS, pl. 18, fig. 22.  
1982 *Bolivina elongata* HANTKEN – SZTRÁKOS, pl. 28, fig. 3.

Description: Test elongate, biserial, lanceolate in outline, oval, strongly compressed in cross-section; length approximately two to three times of maximum width; chambers increasing gradually in size; wall calcareous, hyaline, smooth, finely perforated; sutures distinct, slightly depressed; aperture terminal fissure.

*Bolivina nobilis* HANTKEN, 1875  
Plate 7, Figure 4

- 1875 *Bolivina nobilis* n. sp. – HANTKEN, p. 56, pl. 15, figs. 4a–b.  
1937 *Bolivina nobilis* HANTKEN – CUSHMAN, p. 51, pl. 7, figs. 1–4.  
1947 *Bolivina nobilis* HANTKEN – SUBBOTINA, p. 96, pl. 9, fig. 3.  
1953 *Bolivina nobilis* HANTKEN – SUBBOTINA, p. 225, pl. 10, fig. 10.  
1967 *Bolivina nobilis* HANTKEN – HOFMANN, p. 173, pl. 1, fig. 12.  
1975 *Bolivina nobilis* HANTKEN – BRAGA et al., p. 106, pl. 5, figs. 1–2.  
1975 *Bolivina nobilis* HANTKEN – SAMUEL, p. 134, pl. 74, figs. 4–6.  
1979 *Bolivina nobilis* HANTKEN – SZTRÁKOS, pl. 19, fig. 2.  
1982 *Bolivina nobilis* HANTKEN – SZTRÁKOS, pl. 27, fig. 9.  
1983 *Bolivina nobilis* HANTKEN – KRHOVSKY, p. 77, pl. 1, fig. 8.  
1985 *Bolivina nobilis* HANTKEN – GRÜNIG, p. 265, pl. 5, figs. 12–14.  
1987 *Bolivina nobilis* HANTKEN – REISER, p. 90, pl. 9, figs. 10., 16.  
1992 *Bolivina nobilis* HANTKEN – DARAKCHIEVA and JURANOV, p. 21, pl. 4, figs. 1–2.  
1993 *Bolivina nobilis* HANTKEN – MATHELIN and SZTRÁKOS, p. 78, pl. 32, fig. 9.  
2006 *Bolivina nobilis* HANTKEN – CIMERMAN et al, p. 26, pl. 6, figs. 8–9.  
2006 *Bolivina nobilis* HANTKEN – ORTIZ and THOMAS, p. 113, pl. 4, figs. 7–8.

Description: Test elongate, biserial, lanceolate in outline, oval, mildly compressed in cross-section; length approximately three to four times of maximum width; chambers increasing gradually in size; wall calcareous, hyaline, smooth, finely perforated with numerous longitudinal ribs; aperture terminal fissure.

*Bolivina semistriata* HANTKEN, 1868  
Plate 7, Figure 5

- 1868 *Bolivina semistriata* n. sp. – HANTKEN, p. 95, pl. 2, fig. 34.  
1875 *Bolivina semistriata* HANTKEN – HANTKEN, p. 55, pl. 7, fig. 13.  
1979 *Bolivina semistriata* HANTKEN – SZTRÁKOS, pl. 19, fig. 3.  
1985 *Bolivina semistriata* HANTKEN – KORECZNÉ-LAKY and NAGYNÉ-GELLAI, pl. 15, fig. 2; 47, figs. 1–4.  
1987 *Bolivina semistriata* HANTKEN – SZTRÁKOS, pl. 7, fig. 3.  
1993 *Bolivina semistriata* HANTKEN – MATHÉLIN and SZTRÁKOS, p. 78, pl. 32, fig. 10.  
2006 *Bolivina semistriata* HANTKEN – CIMERMAN et al, p. 28, pl. 6, fig. 10.

Description: Test elongate, slightly curved, biserial, lanceolate in outline, oval, mildly compressed in cross-section; length approximately three to four times of maximum width; chambers increasing gradually in size; wall calcareous, hyaline, smooth, finely perforated with numerous longitudinal ribs; aperture terminal fissure.

*Bolivina* sp.  
Plate 7, Figure 6

Description: Test elongate, biserial, triangular in outline, oval in cross-section; length approximately two times of maximum width; chambers increasing gradually in size, strongly inflated final chamber; sutures distinct, strongly depressed; wall calcareous, hyaline, smooth, finely perforated; aperture terminal fissure.

Superfamily Cassidulinacea D'ORBIGNY, 1839  
Family Cassidulinidae D'ORBIGNY, 1839  
Subfamily Cassidulininae D'ORBIGNY, 1839

Genus *Globocassidulina* VOLOSHINOVA, 1960

Type species: *Globocassidulina globosa* (HANTKEN, 1875)

*Globocassidulina globosa* (HANTKEN, 1875)  
Plate 7, Figure 7–8

- 1875 *Cassidulina globosa* n. sp. – HANTKEN, p. 54, pl. 16, figs. 2a–b.  
1935 *Cassidulina globosa* HANTKEN – CUSHMAN, p. 49, pl. 26, fig. 12.  
1956 *Cassidulina globosa* HANTKEN – HAGN, p. 167, pl. 14, figs. 9–10.  
1975 *Globocassidulina globosa* (HANTKEN) – BRAGA et al., p. 108, pl. 6, fig. 4.  
1979 *Globocassidulina globosa* (HANTKEN) – SZTRÁKOS, pl. 27, fig. 13.  
1985 *Globocassidulina globosa* (HANTKEN) – GRÜNIG, p. 273, pl. 10, fig. 3.  
1987 *Globocassidulina globosa* (HANTKEN) – SZTRÁKOS, pl. 11, fig. 22.  
2006 *Globocassidulina globosa* (HANTKEN) – CIMERMAN et al, p. 28, pl. 6, figs. 12–13.

Description: Test squattish, planispiral, evolute, chambers increasing gradually in size; sutures invisible; wall calcareous, hyaline, smooth, finely perforated; aperture terminal.

Superfamily Buliminacea JONES, 1857  
Family Buliminidae JONES, 1875  
Subfamily Bulimininae KAASSCHIETER, 1961



Genus *Bulimina* D'ORBIGNY, 1826

Type species: *Bulimina marginata* D'ORBIGNY, 1826

*Bulimina affinis* D'ORBIGNY, 1839  
Plate 7, Figure 9

- 1839 *Bulimina affinis* n. sp. – D'ORBIGNY, p. 2, figs. 25–26.  
1926 *Bulimina affinis* D'ORBIGNY – CHAPMAN, p. 37, pl. 5, figs. 15–16.

Description: Test squattish, triserial, rapidly flaring towards apertural end, triangular in outline, oval in cross-section; chambers increasing gradually in size; length approximately two times of maximum width; sutures distinct, mildly depressed; wall calcareous, hyaline, smooth, finely perforated; aperture terminal.

*Bulimina truncana* GÜMBEL, 1868  
Plate 7, Figures 10–11

- 1868 *Bulimina truncana* n. sp. – GÜMBEL, p. 644, pl. 2, fig. 77.  
1875 *Bulimina truncana* GÜMBEL – HANTKEN, p. 61, pl. 7, fig. 5.  
1975 *Bulimina truncana* GÜMBEL – BRAGA et al., p. 106, pl. 4, figs. 15–16.  
1985 *Bulimina truncana* GÜMBEL – GRÜNIG, p. 267, pl. 7, figs. 26–27.  
1985 *Bulimina truncana* GÜMBEL – KORECZNÉ-LAKY – NAGYNÉ-GELLAI, pl. 14, fig. 23; pl. 56, fig. 4.  
1987 *Bulimina truncana* GÜMBEL – SZTRÁKOS, pl. 19, fig. 17.  
2006 *Bulimina truncana* GÜMBEL – CIMERMAN et al., p. 30, pl. 7, fig. 8.

Description: Test elongate, triserial, dropshape in outline, circular in cross-section; chambers increasing gradually in size; length approximately two-three times maximum width; sutures invisible; wall calcareous, hyaline, smooth, finely perforated with numerous strong longitudinal ribs, extending to apertural end; aperture terminal.

*Bulimina* sp.  
Plate 7, Figure 12

Description: Test squattish, triserial, rapidly flaring towards apertural end, dropshape in outline, circular in cross-section; chambers increasing gradually in size; length approximately two-three times maximum width; sutures invisible; wall calcareous, hyaline, smooth, finely perforated; aperture terminal.

Family Uvigerinidae HAECKEL, 1894  
Subfamily Uvigerininae HAECKEL, 1894

Genus *Uvigerina* D'ORBIGNY, 1826

Type species: *Uvigerina pigmea* D'ORBIGNY, 1826

*Uvigerina chirana* CUSHMAN and STONE, 1947  
Plate 7, Figure 13

- 1947 *Uvigerina chirana* n. sp. – CUSHMAN and STONE, p. 17, pl. 2, fig. 25.  
1951 *Uvigerina chirana* CUSHMAN and STONE – CUSHMAN and STAINFORTH, p. 155, pl. 26, fig. 60.  
1956 *Uvigerina chirana* CUSHMAN and STONE – HAGN, p. 150, pl. 13, figs. 14–15.  
1956 *Neouvigerina chirana* (CUSHMAN and STONE) – HOFKER, p. 929, Text-fig. 56.  
1975 *Uvigerina chirana* CUSHMAN and STONE – BRAGA and GRÜNIG, p. 106, pl. 5, fig. 10.  
1977 *Uvigerina chirana* CUSHMAN and STONE – PROTO DECIMA and DE BIASE, p. 95, pl. 2, fig. 10.  
1985 *Uvigerina chirana* CUSHMAN and STONE – GRÜNIG, p. 267, pl. 7, figs. 3–5.  
1987 *Uvigerina chirana* CUSHMAN and STONE – SZTRÁKOS, pl. 18, figs. 6–7.

Description: Test small, triserial, rapidly flaring towards apertural end, lobulate in outline, circular, lobulate in cross-section; chambers increasing gradually in size; length approximately two times maximum width; sutures strongly depressed, curved; wall calcareous, hyaline, smooth, finely perforated with numerous densely arranged nodes; aperture terminal on short neck with lip.

*Uvigerina cocoaensis* CUSHMAN, 1925

Plate 7, Figure 14

- 1925a *Uvigerina cocoaensis* n. sp. – CUSHMAN, p. 68, pl. 10, fig. 12.  
1926 *Uvigerina cocoaensis* CUSHMAN – CUSHMAN and APPLIN, p. 174, pl. 8, fig. 15.  
1933 *Uvigerina cocoaensis* CUSHMAN – ELLISOR, pl. 3, fig. 13.  
1935 *Uvigerina cocoaensis* CUSHMAN – CUSHMAN, p. 39, pl. 15, figs. 11–13.  
1982 *Uvigerina cocoaensis* CUSHMAN – SZTRÁKOS, pl. 25, figs. 3–4, pl. 29, fig. 4.

Description: Test large, triserial, lobulate in outline, circular, lobulate in cross-section; chambers increasing rapidly in size; length approximately two times maximum width; sutures mildly depressed, curved; wall calcareous, hyaline, smooth, finely perforated with numerous fine longitudinal ribs extending to central portion of test; aperture terminal on short neck with lip.

*Uvigerina cocoaensis jacksonensis* CUSHMAN, 1925

Plate 7, Figure 15

- 1925 *Uvigerina jacksonensis* n. sp. – CUSHMAN, p. 67, pl. 10, fig. 13.  
1983 *Uvigerina cocoaensis jacksonensis* CUSHMAN – SZTRÁKOS, p. 134, pl. 2, figs. 20–21; pl. 3, figs. 2, 8–12, 14–16, 20–24.

Description: Test squattish, triserial, rapidly flaring towards apertural end, lobulate in outline, circular, lobulate in cross-section; chambers increasing gradually in size; length approximately two times maximum width; sutures mildly depressed, curved; wall calcareous, hyaline, smooth, finely perforated with numerous fine longitudinal ribs extending to apertural end; aperture terminal on short neck with lip.

*Uvigerina eocaena* GÜMBEL, 1868

Plate 7, Figure 16

- 1868 *Uvigerina eocaena* n. sp. – GÜMBEL, p. 645, pl. 2, fig. 78.  
1975 *Uvigerina eocaena* GÜMBEL – BRAGA et al., p. 107, pl. 5, fig. 17.  
1983 *Uvigerina eocaena* GÜMBEL – SZTRÁKOS, p. 134, pl. 2, figs. 7–15 (cum syn).  
1985 *Uvigerina eocaena* GÜMBEL – GRÜNIG, p. 267, pl. 7, figs. 8–10.  
1993 *Uvigerina eocaena* GÜMBEL – MATHELIN and SZTRÁKOS, p. 79, pl. 34, figs. 9–10.  
2006 *Uvigerina eocaena* GÜMBEL – CIMERMAN et al, p. 30, pl. 7, fig. 12.

Description: Test elongate, triserial, rapidly flaring towards apertural end, lobulate in outline, circular, lobulate in cross-section; chambers increasing gradually in size, inflated, overlapping earlier chambers; length approximately three times maximum width; sutures distinct, mildly depressed, curved; wall calcareous, hyaline, smooth, finely perforated with several longitudinal ribs extending to apertural end; aperture terminal.

*Uvigerina gallowayi* CUSHMAN, 1929

Plate 7, Figure 17–18

- 1929 *Uvigerina gallowayi* n. sp. – CUSHMAN, p. 67, pl. 10., fig. 13.  
1979 *Uvigerina gallowayi* CUSHMAN – SZTRÁKOS, pl. 34, fig. 2.  
1983 *Uvigerina gallowayi* CUSHMAN – SZTRÁKOS, p. 134, pl.3, figs. 5–7, 13, 25 (cum syn).  
1985 *Uvigerina gallowayi* CUSHMAN – KORECZNÉ-LAKY and NAGYNÉ-GELLAI, pl. 16, figs. 5–8.  
2005 *Uvigerina gallowayi* CUSHMAN – NARAYAN et al., p. 138, pl. 5, fig. 24.

Description: Test elongate, triserial, lobulate in outline, circular, lobulate in cross-section; chambers increasing gradually in size, length approximately three times maximum width; sutures distinct, mildly

depressed, curved; wall calcareous, hyaline, smooth, finely perforated with several longitudinal ribs extending to apertural end; aperture terminal on short neck.

*Uvigerina hantkeni* CUSHMAN and EDWARDS, 1937  
Plate 7, Figure 19

- 1875 *Uvigerina pygmea* n. sp. – HANTKEN, p. 52, pl. 7, fig. 4.  
1937 *Uvigerina hantkeni* nomen nudum – CUSHMAN and EDWARDS, p. 60, pl. 8, figs. 15–16.  
1975 *Uvigerina hantkeni* CUSHMAN and EDWARDS – PAPP, p. 282, pl. 1, figs. 9–10.  
1979 *Uvigerina hantkeni* CUSHMAN and EDWARDS – SZTRÁKOS, pl. 24, fig. 3.  
1985 *Uvigerina hantkeni* CUSHMAN and EDWARDS – KORECZNÉ-LAKY and NAGYNÉ-GELLAI, pl. 16, figs. 1–4.  
1986 *Uvigerina hantkeni* CUSHMAN and EDWARDS – CÍCHA et al., p. 136, pl. 4, fig. 1–2, 6.  
1987 *Uvigerina hantkeni* CUSHMAN and EDWARDS – REISER, p. 81, pl. 7, figs. 18, 22.

Description: Test squattish, triserial, rapidly flaring towards apertural end, oval in outline, circular in cross-section; chambers increasing gradually in size; length approximately two times maximum width; sutures slightly depressed, curved; wall calcareous, hyaline, smooth, finely perforated with several longitudinal beads composing ribs; aperture terminal on short neck.

*Uvigerina hourcqi* GRAHAM, DE KLASZ and RÉRAT, 1965  
Plate 7, Figures 21–22

- 1965 *Uvigerina hourcqi* n. sp. – GRAHAM, DE KLASZ and RÉRAT, p. 75, pl. 1, figs. 9–10.  
1983 *Uvigerina hourcqi* GRAHAM, DE KLASZ and RÉRAT – SZTRÁKOS, p. 134, pl. 2, figs. 1–6

Description: Test elongate, triserial, rapidly flaring towards apertural end, lobulate in outline, circular, lobulate in cross-section; chambers increasing gradually in size, overlapping earlier chambers; length approximately two to three times of maximum width; sutures mildly depressed, curved; wall calcareous, hyaline, smooth, finely perforated with several longitudinal, sharp ribs extending to apertural end; aperture terminal on short neck with thickened lip.

*Uvigerina multistriata* HANTKEN, 1871  
Plate 7, Figure 20

- 1871 *Uvigerina multistriata* n. sp. – HANTKEN, p. 129, pl. 2, fig. 14.  
1983 *Uvigerina multistriata* HANTKEN – SZTRÁKOS, p. 136, pl. 1, figs. 7–13.

Description: Test elongate, triserial, lobulate in outline, circular, lobulate in cross-section; chambers increasing gradually in size, inflated, overlapping earlier chambers; length approximately three times maximum width; sutures mildly depressed, curved; wall calcareous, hyaline, smooth, finely perforated with numerous longitudinal ribs extending to apertural end; aperture terminal.

*Uvigerina pygmea* D'ORBIGNY, 1826  
Plate 7, Figure 23

- 1826 *Uvigerina pygmea* n. sp. – D'ORBIGNY, p. 269, pl. 12, figs. 8–9.  
1846 *Uvigerina pygmaea* D'ORBIGNY – D'ORBIGNY, p. 190, pl. 11, figs. 25–26.  
1875 *Uvigerina pygmea* D'ORBIGNY – HANTKEN, p. 62, pl. 7, fig. 4.  
1926 *Uvigerina pygmea* D'ORBIGNY – CHAPMAN, p. 70, pl. 14, fig. 7.  
1929 *Uvigerina pygmea* D'ORBIGNY – GALLOWAY and MORREY, p. 39, pl. 6, fig. 5.  
1932 *Uvigerina pygmaea* D'ORBIGNY – NUTTALL, p. 21, pl. 5, fig. 6.  
1953 *Uvigerina pygmea* D'ORBIGNY – SUBBOTINA, p. 239, pl. 12, figs. 1–6.  
1953 *Uvigerina pygmea* D'ORBIGNY – PAPP and TURNOVSKY, p. 131, pl. 5, fig. 4.  
1958 *Uvigerina pygmea* D'ORBIGNY – NYÍRŐ, p. 35, pl. 1, figs. 9a–d.  
1975 *Uvigerina pygmea* D'ORBIGNY – SAMUEL, p. 137, pl. 76, figs. 1–2.  
1982 *Uvigerina pygmea* D'ORBIGNY – AGIP, pl. 34, fig. 3.  
1984 *Uvigerina pygmaea* D'ORBIGNY – BOERSMA, p. 128, pl. 1, figs. 1–6; pl. 2, figs. 1–5.  
1985 *Uvigerina pygmaea* D'ORBIGNY – KORECZNÉ-LAKY and NAGYNÉ-GELLAI, pl. 147, figs. 1–4.  
1985 *Uvigerina pygmoides* D'ORBIGNY – PAPP and SCHMID, p. 74, pl. 65, figs. 1–5.  
1987 *Uvigerina pygmea* D'ORBIGNY – LOEBLICH and TAPPAN, p. 151, pl. 573, figs. 21–23.  
1992 *Uvigerina pygmea* D'ORBIGNY – DARAKCHIEVA and JURANOV, p. 25, pl. 4, fig. 7.

- 1993 *Uvigerina pigmaea* D'ORBIGNY – KATZ and MILLER, pl. 4, fig. 3.  
2006 *Uvigerina pigmaea* D'ORBIGNY – ORTIZ and THOMAS, p. 136, pl. 11, figs. 9–11.

Description: Test squattish, triserial, rapidly flaring towards apertural end, lobulate in outline, circular, lobulate in cross-section; chambers increasing gradually in size, overlapping earlier chambers; length approximately three times maximum width; sutures mildly depressed, curved; wall calcareous, hyaline, smooth, finely perforated; aperture terminal on short neck.

*Uvigerina rippensis* COLE, 1927  
Plate 7, Figure 24

- 1927 *Uvigerina rippensis* n. sp. – COLE, p. 11, pl. 2, fig. 27.  
1982 *Uvigerina rippensis* COLE – SZTRÁKOS, pl. 25, pl. 30, fig. 2.  
1983 *Uvigerina rippensis* COLE – SZTRÁKOS, p. 136, pl. 1, figs. 5. 14–23.  
1983 *Uvigerina rippensis* COLE – TIALSMA and LOHMANN, p. 38, pl. 14, figs. 6–7.  
1984 *Uvigerina rippensis* COLE – BOERSMA, p. 137, pl. 1, figs. 1–4; pl. 3, figs. 1–6.  
1987 *Uvigerina rippensis* COLE – SZTRÁKOS, pl. 18, fig. 14.  
1998 *Uvigerina rippensis* COLE – BIGNOT, p. 436, pl. 2, figs. 7–8.  
2006 *Uvigerina rippensis* COLE – ORTIZ and THOMAS, p. 136, pl. 11, figs. 12–14

Description: Test squattish, triserial, continually flaring towards apertural end, lobulate in outline, circular, lobulate in cross-section; chambers increasing gradually in size, overlapping earlier chambers; length approximately two times maximum width; sutures mildly depressed, curved; wall calcareous, hyaline, smooth, finely perforated with numerous longitudinal short ribs on earlier chambers and densely arranged oval to circular nodes on final chambers; aperture terminal.

*Uvigerina tenuistriata* REUSS, 1870  
Plate 7, Figure 25

- 1870 *Uvigerina tenuistriata* n. sp. – REUSS, p. 485, pl. 22, figs. 34–37.  
1926 *Uvigerina tenuistriata* REUSS – CHAPMAN, p. 70, pl. 14, fig. 9.

Description: Test squattish, triserial, rapidly flaring towards apertural end, lobulate in outline, circular, lobulate in cross-section; chambers increasing gradually in size, inflated, overlapping earlier chambers; length approximately two times maximum width; sutures distinct, strongly depressed, curved; wall calcareous, hyaline, smooth, finely perforated with several longitudinal, sharp ribs, extending to apertural end; aperture terminal.

Family Reussellidae CUSHMAN, 1933

Genus *Reussella* GALLOWAY, 1933

Type species: *Reussella spinulosa* (REUSS, 1850)

*Reussella elongata* (TERQUEM, 1882)  
Plate 7, Figures 27–28

- 1882 *Verneuilina elongata* n. sp. – TERQUEM, p. 106, pl. 11, fig. 326.  
1950 *Reussella elongata* (TERQUEM) – LE CALVEZ, p. 46, pl. 3, figs. 45–46.  
1961 *Reussella elongata* (TERQUEM) – KAASSCHIETER, p. 191, pl. 9, figs. 7–9.  
1970 *Reussella elongata* (TERQUEM) – LE CALVEZ, p. 119, pl. 22, fig. 4.

Description: Test large, elongate, triserial, rapidly flaring towards apertural end, pyramidal in outline, triangular in cross-section; periphery acute; chambers increasing gradually in size; sutures invisible; wall calcareous, hyaline, smooth, finely perforated; aperture terminal.

*Reussella terquemi* CUSHMAN, 1945  
Plate 7, Figure 26

- 1882 *Verneuilina spinulosa* REUSS – TERQUEM, p. 28, pl. 5, figs. 15–16.  
1945 *Reussella terquemi* n. sp. – CUSHMAN, p. 28, pl. 5, figs. 15–16.  
1961 *Reussella terquemi* (CUSHMAN) – KAASSCHIETER, p. 192, pl. 9, fig. 11.  
1970 *Reussella terquemi* (CUSHMAN) – LE CALVEZ, p. 121, pl. 24, fig. 7.

Description: Test large, wide, triserial, rapidly flaring towards apertural end, pyramidal in outline, triangular in cross-section; periphery acute with wide edges; chambers short, increasing gradually in size; sutures invisible; wall calcareous, hyaline, smooth, finely perforated; aperture terminal, circular.

*Reussella* sp.  
Plate 7, Figures 29–30

Description: Test squattish, triserial, rapidly flaring towards apertural end, pyramidal in outline, triangular in cross-section; periphery acute; chambers increasing gradually in size; sutures barely visible; wall calcareous, hyaline, smooth, finely perforated; aperture terminal.

Superfamily Fursenkoinacea LOEBLICH and TAPPAN, 1961  
Family Fursenkoinidae LOEBLICH and TAPPAN, 1961

Genus *Fursenkoina* LOEBLICH and TAPPAN, 1961

Type species: *Fursenkoina squamosa* (D'ORBIGNY, 1826)

*Fursenkoina hungarica* (HANTKEN, 1868)  
Plate 8, Figures 1–2

- 1868 *Virulina hungarica* n. sp. – HANTKEN, p. 92, pl. III, fig. 26.

Description: Test spindle-shaped, twisted biserial, slightly flaring towards apertural end; lanceolate, lobulate in outline, compressed in cross-section; length approximately four times maximum width; five pairs chambers, increasing rapidly in size; sutures distinct, mildly depressed; wall calcareous, hyaline, smooth, finely perforated; aperture terminal fissure between final two chambers.

*Fursenkoina schreibersiana* (CZIZEK, 1848)  
Plate 8, Figure 3

- 1848 *Virulina schreibersiana* n. sp. – CZIZEK, p. 147, pl. 13., figs. 18–21.  
1875 *Virulina schreibersiana* (CZIZEK) – HANTKEN, p. 53, pl. 7, fig. 15.  
1982 *Fursenkoina schreibersiana* (CZIZEK) – PETERS, p. 77, pl. 5, fig. 26.  
1999 *Fursenkoina schreibersiana* (CZIZEK) – HAYWARD et al., p. 131, pl. 257, figs. 1–12.  
2005 *Fursenkoina schreibersiana* (CZIZEK) – NARAYAN et al., p. 129, pl. 3, figs. 33–34.

Description: Test spindle-shaped, biserial, rapidly flaring towards apertural end; lobulate in outline, oval, mildly compressed in cross-section; length approximately two to three times of maximum width; chambers increasing rapidly in size; sutures distinct, mildly depressed; wall calcareous, hyaline, smooth, finely perforated; aperture terminal fissure between in final two chambers.

Superfamily Pleurostomellacea REUSS, 1860  
Family Pleurostomellidae REUSS, 1860  
Subfamily Pleurostomellinae REUSS, 1860

Genus *Ellipsoglandulina* SILVESTRI, 1900

Type species: *Ellipsoglandulina laevigata* SILVESTRI, 1900

*Ellipsoglandulina multicostata* (GALLOWAY and MORREY, 1929)  
Plate 8, Figure 4

- 1929 *Daucina multicostata* n. sp. – GALLOWAY and MORREY, p. 42, pl. 6, fig. 13.  
1932 *Ellipsoglandulina multicostata* (GALLOWAY and MORREY) – NUTTALL, p. 24, pl. 4, fig. 4.  
1945 *Ellipsoglandulina multicostata* (GALLOWAY and MORREY) – CUSHMAN and STAINFORTH, p. 58, pl. 10, figs. 6–7.  
1949 *Ellipsoglandulina multicostata* (GALLOWAY and MORREY) – BERMÚDEZ, p. 228, pl. 14, figs. 46–47.  
1952 *Ellipsoglandulina multicostata* (GALLOWAY and MORREY) – TODD and KNIKER, p. 23, pl. 4, fig. 11.  
1953 *Ellipsoglandulina multicostata* (GALLOWAY and MORREY) – BECKMANN, p. 380, pl. 23, fig. 13.  
1975 *Ellipsoglandulina multicostata* (GALLOWAY and MORREY) – PROTO DECIMA and DE BIASE, p. 96, pl. 2, fig. 16.  
1985 *Ellipsoglandulina multicostata* (GALLOWAY and MORREY) – GRÜNIG, p. 272, pl. 9, figs. 18–19.  
1998 *Ellipsoglandulina multicostata* (GALLOWAY and MORREY) – ROBERTSON, p. 168, pl. 63, figs. 2a–b.

Description: Test elongate, monoserial, circular in cross-section; length approximately two-three times maximum width; chambers increasing gradually in size, strongly inflated later chambers; sutures distinct, mildly depressed; wall calcareous, hyaline, smooth, finely perforated with numerous fine longitudinal ribs; aperture terminal elongate fissure.

Genus *Nodosarella* RZEHAK, 1895

Type species: *Nodosarella tuberosa* (GÜMBEL, 1868)

*Nodosarella lorifera* (HALKYARD, 1919)  
Plate 8, Figure 5

- 1919 *Nodosaria (Dentalina) lorifera* n. sp. – HALKYARD, p. 70, pl. 4, figs. 2–3.  
1987 *Nodosarella lorifera* (HALKYARD) – SZTRÁKOS, pl. 11, fig. 18.

Description: Test elongate, slightly curved monoserial, circular in cross-section; length approximately six times maximum width; seven chambers increasing gradually in size; sutures distinct, mildly depressed; wall calcareous, hyaline, smooth, finely perforated; aperture terminal, elongate fissure.

*Nodosarella tuberosa* (GÜMBEL, 1868)  
Plate 8, Figures 6–7

- 1868 *Lingulina tuberosa* n. sp. – GÜMBEL, p. 629, pl. 1, fig. 52.  
1956 *Nodosarella tuberosa* (GÜMBEL) – HAGN, p. 157, pl. 14, fig. 4.  
1975 *Nodosarella tuberosa* (GÜMBEL) – PROTO DECIMA and DE BIASE, p. 96, pl. 3, fig. 9.  
1987 *Nodosarella tuberosa* (GÜMBEL) – SZTRÁKOS, pl. 11, fig. 19.

Description: Test elongate, monoserial, circular in cross-section; length approximately four times maximum width; four chambers increasing gradually in size; wall calcareous, hyaline, smooth, finely perforated; aperture terminal elongate fissure.

*Nodosarella* sp.  
Plate 8, Figure 8

Description: Test squattish, monoserial, lobulate in outline, circular in cross-section; length approximately three times maximum width; chambers increasing gradually in size, strongly inflated later chambers; wall calcareous, hyaline, smooth; aperture terminal elongate fissure.

Genus *Pleurostomella* REUSS, 1860

Type species: *Pleurostomella subnodosa* (REUSS, 1846)

*Pleurostomella acuta* HANTKEN, 1875  
Plate 8, Figures 9–11

- 1875 *Pleurostomella acuta* n. sp. – HANTKEN, p. 37, pl. 13, fig. 18.  
 1953 *Pleurostomella* cf. *acuta* HANTKEN – BECKMANN, p. 372, pl. 22, fig. 1.  
 1956 *Pleurostomella acuta* HANTKEN – HAGN, p. 156, pl. 14, fig. 6.  
 1970 *Pleurostomella acuta* HANTKEN – PROTO DECIMA and DE BIASE, p. 96, pl. 3, fig. 7.  
 1979 *Pleurostomella acuta* HANTKEN – SZTRÁKOS, pl. 27, fig. 5.  
 1982 *Pleurostomella acuta* HANTKEN – SZTRÁKOS, pl. 21, fig. 1.  
 1987 *Pleurostomella acuta* HANTKEN – SZTRÁKOS, pl. 11, fig. 11.

Description: Test squattish, loosely biserial, oval in outline, circular in cross-section; length approximately two times maximum width; chambers increasing gradually in size, final pair of chambers extremely inflated; sutures barely visible, slightly depressed, strongly curved; wall calcareous, hyaline, smooth, finely perforated; aperture large oval opening of final chamber.

*Pleurostomella alternans* SCHWAGER, 1866

Plate 8, Figure 12

- 1866 *Pleurostomella alternans* n. sp. – SCHWAGER, p. 238, pl. 6, figs. 79–80.  
 1926 *Pleurostomella alternans* SCHWAGER – CHAPMAN, p. 41, pl. 9, fig. 9.  
 1953 *Pleurostomella alternans* SCHWAGER – BECKMANN, p. 373, pl. 22, fig. 3.  
 1987 *Pleurostomella alternans* HANTKEN – SZTRÁKOS, pl. 11, figs. 15–16.

Description: Test elongate, loosely biserial, lobulate in outline, circular in cross-section; length approximately three to four times of maximum width; four chambers increasing gradually in size; sutures barely visible, slightly depressed; wall calcareous, hyaline, smooth, finely perforated; aperture large oval opening of final chamber.

*Pleurostomella eocaena* GÜMBEL, 1868

Plate 8, Figure 13

- 1868 *Pleurostomella eocaena* n. sp. – GÜMBEL, p. 52, pl. 1, figs. 53a–b.  
 1875 *Pleurostomella eocaena* GÜMBEL – HANTKEN, p. 37, pl. 13, fig. 17.  
 1975 *Pleurostomella eocaena* GÜMBEL – PROTO DECIMA and DE BIASE, p. 96, pl. 3, figs. 6a–b.

Description: Test mildly elongate, loosely biserial, lobulate in outline, circular in cross-section; length approximately three to four times of maximum width; four chambers increasing gradually in size; sutures barely visible, slightly depressed; wall calcareous, hyaline, smooth, finely perforated; aperture oval opening of final chamber.

Remarks: The difference between *Pleurostomella alternans* SCHWAGER and *Pleurostomella eocaena* GÜMBEL is the aperture. *Pleurostomella alternans* SCHWAGER has large, oval aperture contrary to *Pleurostomella eocaena* GÜMBEL which has much smaller aperture.

*Pleurostomella incrassata* HANTKEN, 1884

Plate 8, Figures 15–16

- 1884 *Pleurostomella incrassata* n. sp. – HANTKEN, p. 146, pl. 1, figs. 4, 7.  
 1956 *Pleurostomella incrassata* HANTKEN – HAGN, p. 156, pl. 14, figs. 2, 5.  
 1975 *Pleurostomella incrassata* HANTKEN – PROTO DECIMA and DE BIASE, p. 96, pl. 3, fig. 5.  
 1987 *Pleurostomella incrassata* HANTKEN – SZTRÁKOS, pl. 11, fig. 14.

Description: Test squattish, loosely biserial, lobulate in outline, circular in cross-section; length approximately two to three times of maximum width; four chambers increasing gradually in size, third chamber extremely inflated and large, globular, inflated final chamber; sutures distinct, strongly depressed; wall calcareous, hyaline, smooth, finely perforated; aperture small oval opening of final chamber.

*Pleurostomella* sp.

Plate 8, Figure 14

Description: Test squattish, inflated, loosely biserial, lobulate in outline, circular in cross-section; length approximately two times maximum width; three chambers increasing gradually in size, second chamber extremely inflated; sutures distinct, mildly depressed; wall calcareous, hyaline, smooth, finely perforated; aperture small oval opening of final chamber

Superfamily Stilostomellacea FINLAY, 1947  
Family Stilostomellidae FINLAY, 1947  
Subfamily Stilostomellacea FINLAY, 1947

Genus *Orthomorphina* STAINFORTH, 1952

Type species: *Orthomorphina havanensis* (CUSHMAN and BERMÚDEZ, 1937)

*Orthomorphina rohri* (CUSHMAN and STAINFORTH, 1945)  
Plate 8, Figure 17

- 1945 *Nodogenerina rohri* n. sp. – CUSHMAN and STAINFORTH, p. 39, pl. 5, fig. 26.  
1953 *Orthomorphina rohri* (CUSHMAN and STAINFORTH) – BECKMANN, p. 365, pl. 21, fig. 8.  
1982 *Orthomorphina rohri* (CUSHMAN and STAINFORTH) – SZTRÁKOS, pl. 12, figs. 9–10.  
1985 *Orthomorphina rohri* (CUSHMAN and STAINFORTH) – KORECZNÉ-LAKY and NAGYNÉ-GELLAI, pl. 9, figs. 7–8.

Description: Test elongate, monoserial, circular in cross-section; length approximately four to five times of maximum width; chambers increasing gradually in size; sutures distinct, strongly depressed; wall calcareous, hyaline, smooth, finely perforated with numerous fine, longitudinal ribs extending to suture of final inflated chamber; aperture terminal, circular with lip.

Genus *Stilostomella* GUPPY, 1894

Type species: *Stilostomella rugosa* GUPPY, 1894

*Stilostomella abyssorum* (BRADY, 1881)  
Plate 8, Figures 18–19

- 1881 *Nodosaria abyssorum* n. sp. – BRADY, p. 63, pl. 63, figs. 8–9.  
1934 *Ellipsonodosaria nuttalli* n. sp. – CUSHMAN and JARVIS, p. 72, pl. 10, fig. 6.  
1953 *Stilostomella abyssorum* (BRADY) – BECKMANN, p. 369, pl. 21, fig. 33.  
1987 *Stilostomella* cf. *abyssorum* (BRADY) – SZTRÁKOS, pl. 7, figs. 6–7.

Description: Test elongate, monoserial, beads-like in outline, circular in cross-section; length approximately five to six times of maximum width; chambers increasing moderately in size; sutures broad, hoop-like, strongly depressed; wall calcareous, hyaline, smooth, finely perforated; aperture terminal, circular on short neck with thickened lip.

*Stilostomella adolphina* (D'ORBIGNY, 1846)  
Plate 8, Figures 20–22

- 1846 *Dentalina adolphina* n. sp. – D'ORBIGNY, p. 51, pl. 2, figs. 18–20.  
1868 *Dentalina adolphina* D'ORBIGNY – GÜMBEL, p. 45, pl. 1, fig. 32.  
1953 *Siphonodosaria adolphina* (D'ORBIGNY) – SUBBOTINA, p. 180, pl. 6, figs. 1–2.  
1969 *Nodosaria adolphina* (D'ORBIGNY) – KRAYEVA and ZERNECKII, p. 41, pl. 14, fig. 6.  
1969 *Stilostomella adolphina* (D'ORBIGNY) – RÖGL, p. 80, pl. 3, fig. 12.  
1975 *Nodosaria adolphina* (D'ORBIGNY) – SAMUEL, p. 119, pl. 70, figs. 2a–b.  
1979 *Stilostomella adolphina* (D'ORBIGNY) – SZTRÁKOS, pl. 19, fig. 8.  
1983 *Stilostomella adolphina* (D'ORBIGNY) – KRHOVSKY, p. 78, pl. 2, fig. 2.  
1985 *Stilostomella adolphina* (D'ORBIGNY) – KORECZNÉ-LAKY and NAGYNÉ-GELLAI, pl. 10, fig. 8., pl. 24. 4–5; pl. 118, figs. 1–2.  
1985 *Stilostomella adolphina* (D'ORBIGNY) – PAPP and SCHMID, p. 31, pl. 14, figs. 8–11.  
1987 *Stilostomella adolphina* (D'ORBIGNY) – WENGER, p. 287, pl. 10, figs. 21–22.



Description: Test elongate, monoserial, beads-like in outline, circular in cross-section; length approximately five - six times maximum width; chambers increasing gradually in size; sutures distinct strongly depressed; wall calcareous, hyaline, smooth, finely perforated with numerous random arranged circular to oval nodes; aperture terminal, circular on short neck with thickened lip.

*Stilostomella consobrina* (D'ORBIGNY, 1846)

Plate 8, Figure 23

- 1846 *Dentalina consobrina* n. sp. – D'ORBIGNY, p. 46, pl. 2, figs. 1–3.  
1953 *Stilostomella consobrina* (D'ORBIGNY) – BECKMANN, p. 370, pl. 21, figs. 24–25.  
1979 *Stilostomella consobrina* (D'ORBIGNY) – SZTRÁKOS, pl. 19, fig. 9.  
1982 *Stilostomella consobrina* (D'ORBIGNY) – SZTRÁKOS, pl. 16, fig. 5.  
1985 *Stilostomella consobrina* (D'ORBIGNY) – PAPP and SCHMID, p. 29, pl. 11, figs. 1–5.  
1987 *Stilostomella consobrina* (D'ORBIGNY) – SZTRÁKOS, pl. 7, fig. 11.

Description: Test elongate, monoserial, tubular in outline, circular in cross-section; length approximately five to six times of maximum width; chambers increasing moderately in size, first chamber strongly inflated with short tubular spine; sutures barely visible, slightly depressed; wall calcareous, hyaline, smooth, finely perforated; aperture terminal, circular.

*Stilostomella curvatura* (CUSHMAN, 1939)

Plate 8, Figure 24

- 1939 *Dentalina curvatura* n. sp. – CUSHMAN, p. 105, pl. 28, fig. 5.  
1953 *Stilostomella curvatura* (CUSHMAN) – BECKMANN, p. 370, pl. 21, figs. 26–27.  
1987 *Stilostomella curvatura* (CUSHMAN) – SZTRÁKOS, pl. 17, fig. 12.

Description: Test elongate, slightly curved, monoserial, beads-like in outline, circular in cross-section; length approximately five to six times of maximum width; chambers increasing gradually in size; sutures distinct, strongly depressed; wall calcareous, hyaline; smooth, finely perforated with numerous random arranged circular to oval nodes; aperture terminal, circular on short neck with thickened lip.

*Stilostomella elegans* (D'ORBIGNY, 1846)

Plate 8, Figures 25–26

- 1846 *Dentalina elegans* n. sp. – D'ORBIGNY, p. 45, pl. 1, figs. 52–56.  
1979 *Stilostomella elegans* (D'ORBIGNY) – SZTRÁKOS, pl. 19, fig. 12.  
1985 *Stilostomella elegans* (D'ORBIGNY) – KORECZNÉ-LAKY and NAGYNÉ-GELLAI, pl. 11, fig. 6; pl. 104, figs. 2, 13.  
1987 *Stilostomella elegans* (D'ORBIGNY) – SZTRÁKOS, pl. 7, fig. 12.

Description: Test elongate, slightly curved, monoserial, beads-like in outline, circular in cross-section; length approximately five to six times of maximum width; chambers increasing gradually in size, final four to five chambers spindle-shaped; sutures distinct, broad, strongly depressed; wall calcareous, hyaline; smooth, finely perforated; aperture terminal, circular on short neck with strongly thickened lip.

*Stilostomella emaciata* (REUSS, 1851)

Plate 8, Figure 27

- 1851 *Dentalina emaciata* n. sp. – REUSS, p. 63, pl. 3, fig. 9.  
1979 *Stilostomella emaciata* (REUSS) – SZTRÁKOS, pl. 19, fig. 14.

Description: Test elongate, curved, monoserial, tubular in outline, circular in cross-section; length approximately five to six times of maximum width; chambers increasing moderately in size, first chamber globular, inflated with short tubular spine; sutures distinct, slightly depressed; wall calcareous, hyaline; smooth, finely perforated; aperture terminal, circular.

*Stilostomella hoernesii* (HANTKEN, 1868)

Plate 8, Figure 28

1868 *Dentalina hoernesii* n. sp. – HANTKEN, p. 89, pl. 1. fig. 2.

1979 *Stilostomella hoernesii* (HANTKEN) – SZTRÁKOS, pl. 19, fig. 15.

1982 *Stilostomella hoernesii* (HANTKEN) – SZTRÁKOS, pl. 16, fig. 4.

1985 *Stilostomella hoernesii* (HANTKEN) – KORECZNÉ-LAKY and NAGYNÉ-GELLAI, pl. 10, figs. 5–6.

Description: Test elongate, monoserial, beads-like in outline, circular in cross-section; length approximately five to six times of maximum width; chambers increasing gradually in size; sutures distinct strongly depressed; wall calcareous, hyaline; smooth, finely perforated with numerous longitudinal ribs; aperture terminal, circular.

*Stilostomella pauperata* (D'ORBIGNY, 1846)

Plate 8, Figure 29

1846 *Dentalina pauperata* n. sp. – D'ORBIGNY, p. 46, pl. 1, figs. 57–58.

1979 *Stilostomella pauperata* (D'ORBIGNY) – SZTRÁKOS, pl. 19, fig. 13.

1985 *Stilostomella pauperata* (D'ORBIGNY) – KORECZNÉ-LAKY and NAGYNÉ-GELLAI, pl. 10, fig. 12; pl. 99, fig. 3; pl. 102, fig. 13; pl. 104, fig. 11.

Description: Test elongate, slightly curved, monoserial, beads-like in outline, circular in cross-section; length approximately five to six times of maximum width; chambers increasing gradually in size; sutures distinct strongly depressed; wall calcareous, hyaline; smooth, finely perforated with numerous irregularly arranged nodes; aperture terminal, circular.

*Stilostomella* sp.

Plate 8, Figure 30

Description: Test elongate, monoserial, beads-like in outline, circular in cross-section; length approximately four – five times maximum width; chambers increasing gradually in size; sutures distinct strongly depressed; wall calcareous, hyaline; smooth, finely perforated; aperture terminal, circular.

Superfamily Discorbacea EHRENBERG, 1838

Family Bagginidae CUSHMAN, 1927

Subfamily Baggininae CUSHMAN, 1927

Genus *Cancris* DE MONTFORT, 1808

Type species: *Cancris auriculatus* DE MONTFORT, 1808

*Cancris* sp.

Plate 9, Figures 1–2

Description: Test trochospiral, circular in outline, spiral side evolute, umbilical side involute, biconvex in cross-section; periphery acute; chambers increasing gradually in size, five to six chambers in final whorl; sutures slightly depressed, radial; wall calcareous, hyaline; smooth, perforated; aperture umbilical.

Genus *Valvulineria* CUSHMAN, 1926

Type species: *Valvulineria californica* CUSHMAN, 1926

*Valvulineria* sp.

Plate 9, Figure 3

Description: Test trochospiral, spiral side evolute, umbilical side involute, plano-convex in cross-section; periphery mildly undulate; chambers increasing gradually in size, five chambers in final whorl; sutures strongly depressed, radial, mildly curved; wall calcareous, hyaline; smooth, perforated; aperture umbilical.

Family Eponididae HOFKER, 1951  
Subfamily Eponidinae HOFKER, 1951

Genus *Eponides* DE MONTFORT, 1808

Type species: *Eponides repandus* FICHTEL and MOLL, 1798

*Eponides budensis* (HANTKEN, 1875)  
Plate 9, Figures 4–5

1875 *Truncatulina budensis* n. sp. – HANTKEN, p. 65, pl. 8, fig. 6.

Description: Test trochospiral, spiral side evolute, umbilical side involute, biconvex in cross-section; periphery rounded; chambers increasing gradually in size, six to seven chambers in final whorl; sutures slightly depressed, radial, mildly curved; wall calcareous, hyaline; smooth, coarsely perforated; aperture umbilical.

*Eponides haidingeri* (D'ORBIGNY, 1846)  
Plate 9, Figures 6–7

1846 *Rotalina haidingeri* n. sp. – D'ORBIGNY, p. 154, pl. 8, figs. 7–9.  
1875 *Pulvinulina haidingeri* (D'ORBIGNY) – HANTKEN, p. 67, pl. 15, figs. 10a–b.

Description: Test trochospiral, spiral side evolute, umbilical side involute, plano-convex in cross-section; periphery rounded; chambers increasing gradually in size, five to six chambers in final whorl; sutures strongly depressed, radial, mildly curved; wall calcareous, hyaline; smooth, coarsely perforated; aperture umbilical.

*Eponides umbonatus* (REUSS, 1851)  
Plate 9, Figures 8–9

1851 *Rotalina umbonata* n. sp. – REUSS, p. 75, pl. 5, figs. 35a–c.  
1875 *Pulvinulina umbonata* (REUSS) – HANTKEN, p. 67, pl. 9, figs. 8a–c.  
1926 *Pulvinulina umbonata* (REUSS) – CHAPMAN, p. 84, pl. 17, fig. 3.  
1928 *Eponides umbonata* (REUSS) – COLE, p. 15, pl. 2, fig. 6.  
1929 *Rotalia umbonata* (REUSS) – GALLOWAY and MORREY, p. 26, pl. 4, figs. 1a–c.  
1929 *Eponides umbonata* (REUSS) – CUSHMAN, p. 98, pl. 14, figs. 8a–c.  
1949 *Eponides umbonatus* (REUSS) – BERMÚDEZ, p. 249, pl. 17, figs. 22–24.  
1961 *Eponides umbonatus* (REUSS) – KAASSCHIETER, p. 211, pl. 13, fig. 1.  
1962 *Eponides umbonatus* (REUSS) – KIESEL, p. 71, pl. 10, fig. 10.  
1975 *Oridorsalis umbonatus* (REUSS) – PROTO DECIMA and DE BIASE, p. 97, pl. 2, figs. 14a–c.  
1979 *Eponides umbonatus* (REUSS) – SZTRÁKOS, pl. 25, fig. 12.  
1982 *Oridorsalis umbonatus* (REUSS) – AGIP, pl. 40, fig. 10.  
1987 *Oridorsalis umbonatus* (REUSS) – WENGER, p. 311, pl. 17, figs. 6–8.  
1993 *Oridorsalis umbonatus* (REUSS) – DARAKCHIEVA and JURANOV, p. 68, pl. 3, figs. 7, 9.  
1997 *Oridorsalis umbonatus* (REUSS) – BORNMALM, pl. 24, figs. j–l.  
2001 *Oridorsalis umbonatus* (REUSS) – MÜHLSTRASSER, pl. 3, figs. 4–5.

Description: Test trochospiral, spiral side evolute, umbilical side involute, biconvex in cross-section; periphery with thin keel; chambers increasing gradually in size, five to six chambers in final whorl; sutures strongly depressed, radial, mildly curved; wall calcareous, hyaline; smooth, coarsely perforated; aperture umbilical.

*Eponides* sp.  
Plate 9, Figure 10

Description: Test trochospiral, spiral side evolute, umbilical side involute, biconvex in cross-section; periphery rounded; chambers increasing gradually in size, five to six chambers in final whorl; sutures slightly depressed, radial, mildly curved; wall calcareous, hyaline; smooth, coarsely perforated; aperture umbilical.

Family Discorbidae EHRENBERG, 1838

Genus *Discorbis* LAMARCK, 1804

Type species: *Discorbis vesicularis* LAMARCK, 1804

*Discorbis elegans* HANTKEN, 1875  
Plate 9, Figure 11

1875 *Discorbina elegans* n. sp. – HANTKEN, p. 66, pl. 9, fig. 3, pl. 15, fig. 7.

Description: Test trochospiral, spiral side evolute, umbilical side involute, plano-convex in cross-section; periphery rounded; chambers increasing gradually in size, five to six chambers in final whorl; sutures slightly elevated, radial; wall calcareous, hyaline; smooth, coarsely perforated; aperture umbilical.

*Discorbis perplexa* LE CALVEZ, 1949  
Plate 9, Figures 12–13

1949 *Discorbis perplexa* LE CALVEZ – LE CALVEZ, p. 21, pl. 2, figs. 18–20.

1970 *Discorbis perplexa* LE CALVEZ – LE CALVEZ, p. 137, figs. 46–48.

Description: Test trochospiral, spiral side evolute, umbilical side involute, biconvex in cross-section; periphery with broad keel; chambers increasing gradually in size, five – six chambers in final whorl; sutures strongly depressed, radial; wall calcareous, hyaline; smooth, coarsely perforated; aperture umbilical.

Superfamily Discorbinellacea SIGAL, 1952  
Family Parrelloididae HOFKER, 1956

Genus *Cibicidoides* THALMANN, 1939

Type species: *Cibicidoides mundula* (BRADY, PARKER and JONES, 1888)

*Cibicidoides eocaenus* (GÜMBEL, 1868)  
Plate 9, Figures 14–16

1868 *Rotalia eocaena* n. sp. – GÜMBEL, p. 650, pl. 2, figs. 87 a–b.

1986 *Cibicidoides eocaenus* (GÜMBEL) – VAN MORKHOVEN et al., p. 256, pl. 86A, figs. 1–4; pl. 86B, figs. 1–2; pl. 86C, 1–3 (cum syn).

2006 *Cibicidoides eocaenus* (GÜMBEL) – ORTIZ and THOMAS, p. 115, pl. 5, fig. 2.

Description: Test trochospiral, spiral side evolute, umbilical side involute, biconvex in cross-section; periphery rounded; chambers increasing gradually in size, seven to eighth chambers in final whorl; sutures barely visible, radial; wall calcareous, hyaline; smooth, coarsely perforated; aperture umbilical.

*Cibicidoides perlucidus* (NUTTALL, 1932)

Plate 9, Figures 17–18

1932 *Cibicides perlucida* n. sp. – NUTTALL, p. 33, pl. 8, figs. 10–12.

1986 *Cibicidoides perlucidus* (NUTTALL) – VAN MORKHOVEN et al., p. 260, pl. 86D, figs. 1–2.

Description: Test trochospiral, spiral side evolute, umbilical side involute, biconvex in cross-section; periphery acute; large central umbilicus; chambers increasing gradually in size, eighth - nine chambers in final whorl; sutures thick, slightly depressed, gently curved, radial; wall calcareous, hyaline; smooth, coarsely perforated; aperture umbilical.

Superfamily Planorbulinacea SCHWAGER, 1877

Family Planulinidae BERMÚDEZ, 1952

Genus *Planulina* D'ORBIGNY, 1826

Type species: *Planulina ariminensis* (D'ORBIGNY, 1826)

*Planulina austriaca* (D'ORBIGNY, 1826)

Plate 9, Figures 19–20

1826 *Anomalina austriaca* n. sp. – D'ORBIGNY, p. 172, pl. 10, figs. 4–9.

1985 *Planulina austriaca* (D'ORBIGNY) – PAPP and SCHMID, p. 65, pl. 58, figs. 1–8; pl. 59, figs. 1–6.

Description: Test trochospiral, evolute, biconvex, mildly compressed in apertural view; periphery with rounded keel; chambers increasing gradually in size, eighth to nine chambers in final whorl; sutures strongly depressed, curved; wall calcareous, hyaline; smooth, perforated; aperture interiomarginal.

*Planulina compressa* (HANTKEN, 1875)

Plate 9, Figure 21

1875 *Truncatulina compressa* n. sp. – HANTKEN, p. 62, pl. 8, figs. 8a–b.

1987 *Truncatulina costata* HANTKEN – SZTRÁKOS, pl. 20, fig. 2.

Description: Test trochospiral, evolute, biconvex, strongly compressed in apertural view; periphery with rounded keel; chambers increasing gradually in size, sixth to eighth chambers in final whorl; sutures barely visible, curved; wall calcareous, hyaline; smooth, perforated; aperture interiomarginal.

*Planulina costata* (HANTKEN, 1875)

Plate 9, Figures 22–23

1875 *Truncatulina costata* n. sp. – HANTKEN, p. 63, pl. 9, fig. 2.

1987 *Planulina costata* HANTKEN – SZTRÁKOS, pl. 10, fig. 5.

Description: Test trochospiral, evolute, biconvex, mildly compressed in apertural view; periphery with rounded keel; chambers increasing gradually in size, eighth to nine chambers in final whorl; sutures strongly depressed, curved; wall calcareous, hyaline; smooth, coarsely perforated; aperture interiomarginal

Family Cibicididae CUSHMAN, 1927

Subfamily Cibicidinae CUSHMAN, 1927

Genus *Cibicides* DE MONTFORT, 1808

Type species: *Cibicides reflugens* DE MONTFORT, 1808

*Cibicides ammophilus* (GÜMBEL, 1868)

Plate 9, Figure 24

1868 *Rotalia ammophila* n. sp. – GÜMBEL, p. 652, pl. 2, figs. 90a–b.

1982 *Cibicides ammophilus* (GÜMBEL) – SZTRÁKOS, pl. 20, fig. 6.

Description: Test trochospiral, squattish, involute, plano-convex in cross-section; chambers increasing gradually in size, ten to eleven chambers in final whorl; deep umbilicus; sutures depressed, strongly curved; wall calcareous, hyaline; smooth, perforated; aperture interiomarginal.

*Cibicides boueanus* (D'ORBIGNY, 1846)

Plate 9, Figures 25–27

1846 *Truncatulina boueana* n. sp. – D'ORBIGNY, p. 169, pl. 9, figs. 24–26.

1951 *Cibicides boueanus* (D'ORBIGNY) – MARKS, p. 72, pl. 8, figs. 9a–b.

1985 *Cibicides boueanus* (D'ORBIGNY) – PAPP and SCHMID, p. 64, pl. 56, figs. 6–9.

Description: Test trochospiral, involute, plano-convex in cross-section; periphery with rounded keel; chambers increasing gradually in size, eighth to nine chambers in final whorl; deep umbilicus; sutures slightly depressed on umbilical side, strongly elevated on spiral side, strongly curved; wall calcareous, hyaline; smooth, perforated; aperture interiomarginal.

*Cibicides carinatus* (TERQUEM, 1882)

Plate 9, Figures 28–29

1882 *Truncatulina carinatus* n. sp. – TERQUEM, p. 94, pl. 10, figs. 1–2.

1949 *Cibicides carinatus* (TERQUEM) – LE CALVEZ, p. 45, pl. 4, figs. 72–74.

1968 *Cibicides carinatus* (TERQUEM) – POZARYSKA and SZCZECHURA, p. 75, pl. 11, fig. 1.

1970 *Cibicides carinatus* (TERQUEM) – LE CALVEZ, p. 180, figs. 66–68.

1970 *Cibicides carinatus* (TERQUEM) – MYATLYUK, p. 152, pl. 63, fig. 1.

1974 *Cibicides carinatus* (TERQUEM) – SZCZECHURA and POZARYSKA, p. 86, pl. 28, figs. 6–7.

1975 *Cibicides carinatus* (TERQUEM) – SAMUEL, p. 148, pl. 86, figs. 6a–b.

1987 *Cibicides carinatus* (TERQUEM) – SZTRÁKOS, pl. 20, figs. 8–9.

Description: Test trochospiral, spiral side evolute, umbilical side involute, plano-convex in cross-section; periphery with rounded, relatively broad keel; chambers increasing gradually in size, eighth to nine chambers in final whorl; sutures strongly depressed, curved; wall calcareous, hyaline; smooth, perforated; aperture interiomarginal.

*Cibicides mauricensis* HOWE and ROBERTS, 1939

Plate 10, Figure 1

1939 *Cibicides mauricensis* n. sp. – HOWE and ROBERTS in HOWE, p. 87, pl. 13, figs. 4–5.

Description: Test trochospiral, spiral side evolute, umbilical side involute, plano-convex in cross-section; periphery rounded; chambers increasing gradually in size, six to seven chambers in final whorl; deep umbilicus; sutures slightly depressed; wall calcareous, hyaline, smooth, perforated; aperture interiomarginal.

*Cibicides oligocenicus* SAMOILOVA, 1947

Plate 10, Figure 2

1947 *Cibicides dutemplei* var. *oligocenica* (SAMOILOVA) n. ssp. – SAMOILOVA, p. 96, figs. 34–36.

1977 *Cibicides oligocenicus* (SAMOILOVA) – POZARYSKA, p. 39, pl. 6, figs. 1–6; pl. 7, figs. 3a–c; pl. 13, figs. 8–9.

Description: Test trochospiral, spiral side evolute, umbilical side involute, plano-convex in cross-section; periphery rounded; chambers increasing gradually in size, eighth to nine chambers in final whorl; sutures strongly depressed, curved; wall calcareous, hyaline; smooth, perforated; aperture interiomarginal.

*Cibicides proprius* BROTZEN, 1948  
Plate 10, Figure 3

- 1948 *Cibicoides proprius* n. sp. – BROTZEN, p. 78, pl. 12, figs. 3–4.  
1961 *Cibicides proprius* BROTZEN – KAASSCHIETER, p. 222, pl. 13, figs. 9–10; pl. 14, fig. 7.

Description: Test trochospiral, squattish, umbilical side involute, plano-convex in cross-section; periphery rounded; chambers increasing gradually in size, eighth to nine chambers in final whorl; sutures slightly depressed, curved; wall calcareous, hyaline; smooth, coarsely perforated; aperture interiomarginal.

*Cibicides pseudolobatulus* PERELIS and REISS, 1975  
Plate 10, Figure 4

- 1975 *Cibicides pseudolobatulus* PERELIS and REISS – PERELIS and REISS, p. 77, pl. 4, figs. 1–7.  
1993 *Cibicides pseudolobatulus* PERELIS and REISS – HOTTINGER, HALICZ and REISS, p. 116, pl. 152, figs. 7–11.

Description: Test trochospiral, umbilical side involute, spiral side evolute, plano-convex in cross-section; periphery rounded; chambers increasing gradually in size, six chambers in final whorl; deep umbilicus; sutures depressed, radial; wall calcareous, hyaline; smooth, coarsely perforated; aperture interiomarginal.

*Cibicides pseudoungerianus* (CUSHMAN, 1922)  
Plate 10, Figure 5

- 1884 *Truncatulina ungeriana* (not D'ORBIGNY) – BRADY, p. 664, pl. 94, fig. 9.  
1899 *Truncatulina ungeriana* (not D'ORBIGNY) – FLINT, p. 333, pl. 77, fig. 2.  
1918 *Truncatulina ungeriana* (not D'ORBIGNY) – CUSHMAN, p. 69, pl. 24., fig. 1.  
1922 *Truncatulina pseudoungeriana* n. sp. – CUSHMAN, p. 97, pl. 20, fig. 9.  
1931 *Cibicides pseudoungeriana* (CUSHMAN) – CUSHMAN, p. 123, pl. 22, figs. 3–7.  
1951 *Cibicides ungerianus* (D'ORBIGNY) – MARKS, p. 73, pl. 8, figs. 2a–b.  
1953 *Cibicides pseudoungerianus* (CUSHMAN) – BECKMANN, p. 403, pl. 28, figs. 3–4.  
1959 *Cibicides pseudoungerianus* (CUSHMAN) – DIECI, p. 100, pl. 8, fig. 1.  
1960 *Cibicides pseudoungerianus* (CUSHMAN) – BARKER, p. 194, pl. 94, figs. 9a–c.  
1960 *Cibicides pseudoungerianus* (CUSHMAN) – CHRISTODOULOU, p. 94, pl. 14, fig. 7a–b.  
1970 *Cibicides pseudoungerianus* (CUSHMAN) – VERDENIUS, pl. 6, fig. 3.  
1971 *Cibicides pseudoungerianus* (CUSHMAN) – VERHOEVE, p. 63, pl. 3, fig. 5a–c.  
1976 *Cibicides* cf. *pseudoungerianus* (CUSHMAN) – PFLUM and FRERICHS, pl. 2, fig. 9, pl. 3, figs. 1–2.  
1978 *Cibicides pseudoungerianus* (CUSHMAN) – BROLSMA, pl. 3, figs. 7a–c.  
1979 *Cibicides ungerianus* (D'ORBIGNY) – HAGEMAN, p. 92, pl. 4, figs. 2a–c, non fig. 3a–b.  
1980 *Cibicoides floridanus* (CUSHMAN) – BREMER et al., p. 24, pl. 3, figs. 12–14.  
1982 *Cibicoides pseudoungerianus* (CUSHMAN) – AGIP, pl. 52, fig. 1.  
1991 *Cibicoides pseudoungerianus* (CUSHMAN) – CIMERMAN and LANGER, p. 69, pl. 74, figs. 2–3.  
1991 *Cibicides pseudoungerianus* (CUSHMAN) – VERHALLEN, p. 129, pl. 16, figs. 1–4.  
2000 *Cibicides pseudoungerianus* (CUSHMAN) – DEN DULK, pl. 6, figs. 3a–c.  
2000 *Cibicides pseudoungerianus* (CUSHMAN) – KOUWENHOVEN, pl. 1, figs. 3a–c.  
2001 *Cibicides pseudoungerianus* (CUSHMAN) – MÜHLSTRASSER, p. 73, pl. 4, figs. 9–11.  
2006 *Cibicides pseudoungerianus* (CUSHMAN) – SCHWEIZER, p. 125, pl. 7, figs. a–p.

Description: Test trochospiral, squattish, spiral side evolute, umbilical side involute, plano-convex in cross-section; periphery undulate; chambers increasing gradually in size, eighth to nine chambers in final whorl; sutures slightly depressed, curved; wall calcareous, hyaline, smooth, coarsely perforated; aperture interiomarginal, extending onto spiral side.

*Cibicides sublobatulus* (GÜMBEL, 1868)  
Plate 10, Figures 6–7

- 1868 *Truncatulina sublobatula* n. sp. – GÜMBEL, p. 659, pl. 2, figs. 103a–c.  
1988 *Cibicides sublobatulus* (GÜMBEL) – HORVÁTH-KOLLÁNYI, p. 81, pl. 22, figs. 1–3.

Description: Test trochospiral, umbilical side involute, plano-convex to biconvex in cross-section; periphery rounded; chambers increasing gradually in size, eighth to nine chambers in final whorl;

sutures slightly depressed, curved; wall calcareous, hyaline; smooth, coarsely perforated; aperture interiomarginal.

*Cibicides sulzensis* (HERRMANN, 1917)

Plate 10, Figure 8

1917 *Discorbina sulzensis* n. sp. – HERRMANN, p. 290, pl. 3, fig. 26.

1958 *Cibicides sulzensis* (HERRMANN) – BATJES, p. 143, pl. 9, fig. 5.

1961 *Cibicides sulzensis* (HERRMANN) – KAASSCHIETER, p. 223, pl. 13, fig. 11.

Description: Test trochospiral, umbilical side involute, plano-convex in cross-section; periphery rounded; chambers increasing gradually in size, eighth to nine chambers in final whorl; deep umbilicus; sutures barely visible; wall calcareous, hyaline; smooth, perforated; aperture interiomarginal.

*Cibicides ungerianus* (D'ORBIGNY, 1846)

Plate 10, Figure 9

1846 *Rotalina ungeriana* n. sp. – D'ORBIGNY, p. 157, pl. 8, figs. 16–18.

1951 *Cibicides ungerianus* D'ORBIGNY – MARKS, p. 73, pl. 8, figs. 2a–b.

1960 *Cibicides ungerianus* D'ORBIGNY – CÍCHA and ZAPLETOVA, p. 13, pl. 6, figs. 4–6.

1985 *Cibicides ungerianus* D'ORBIGNY – PAPP and SCHMID, p. 60, pl. 51, figs. 7–11.

Description: Test trochospiral, squattish, umbilical side involute, spiral side evolute, plano-convex in cross-section; periphery rounded; chambers increasing gradually in size, eighth to nine chambers in final whorl; sutures slightly depressed, curved; thick spiral line on spiral side; wall calcareous, hyaline; smooth, coarsely perforated; aperture interiomarginal.

*Cibicides westi* HOWE, 1939

Plate 10, Figures 10–12

1939 *Cibicides westi* n. sp. – HOWE, p. 88, pl. 13, figs. 20–22.

1949 *Cibicides westi* HOWE – BANDY, p. 112, pl. 20, fig. 7.

1961 *Cibicides westi* HOWE – KAASSCHIETER, p. 218, pl. 13, fig. 7.

Description: Test trochospiral, involute, plano-convex in cross-section; periphery rounded; chambers increasing gradually in size, eighth to nine chambers in final whorl; large-sized, triangular final chamber in lateral view; sutures slightly elevated, radial; relatively deep umbilicus; wall calcareous, hyaline; smooth, coarsely perforated; aperture interiomarginal, extending to spiral side.

*Cibicides* sp.

Plate 10, Figure 13

Description: Test trochospiral, involute, plano-convex in cross-section; periphery rounded; chambers increasing gradually in size, eighth to nine chambers in final whorl; sutures slightly depressed, radial; wall calcareous, hyaline; smooth, coarsely perforated; aperture interiomarginal, extending to spiral side.

Subfamily Stichocibicidinae SAIDOVA, 1981

Genus *Dyocibicides* CUSHMAN and VALENTINE, 1930

Type species: *Dyocibicides biserialis* CUSHMAN and VALENTINE, 1930

*Dyocibicides uniserialis* THALMANN, 1933

Plate 10, Figures 14–15

1884 *Truncatulina variabilis* D'ORBIGNY – BRADY, pl. 93, fig. 7.



1933 *Dyocibicides uniserialis* n. sp. – THALMANN, p. 254, pl. 93, fig. 7.

Description: Test elongate, trochospiral, involute in earlier chambers, becoming irregularly monoserial later (three or four monoserial chambers, compressed in cross-section; chambers increasing gradually in size, younger chambers much bigger than earlier; periphery rounded; sutures distinct, strongly depressed; wall calcareous, smooth, finely perforated; aperture terminal.

Superfamily Asterigerinacea D'ORBIGNY, 1839

Family Asterigerinidae D'ORBIGNY, 1839

Genus *Asterigerina* D'ORBIGNY, 1839

Type species: *Asterigerina carinata* D'ORBIGNY, 1839.

*Asterigerina* sp.

Plate 10, Figure 16

Description: Test trochospiral, umbilical side involute, spiral side evolute, plano-convex in cross-section; periphery rounded; chambers increasing gradually in size, eighth to nine chambers in final whorl; secondary chambers forming a star or rosette pattern around umbilical region; sutures slightly depressed, curved; wall calcareous, hyaline; smooth, finely perforated; aperture interiomarginal, extending to umbilical region.

Family Epistomariidae HOFKER, 1954

Subfamily Epistomariinae HOFKER, 1954

Genus *Nuttallides* FINLAY, 1939

Type species: *Nuttallides truempyi* (NUTTALL, 1930)

*Nuttallides* sp.

Plate 10, Figures 17–18

Description: Test trochospiral, triangular in lateral view, spiral side involute, umbilical side involute, plano-convex in cross-section; periphery rounded with thin keel; chambers increasing gradually in size, four to five chambers in final whorl wall calcareous, hyaline; smooth, finely perforated; aperture interiomarginal, extending to umbilical region.

Superfamily Nonionacea SCHULTZE, 1854

Family Nonionidae SCHULTZE, 1854

Subfamily Nonioninae SCHULTZE, 1854

Genus *Nonion* DE MONTFORT, 1808

Type species: *Nonion faba* (FICHTEL and MOLL, 1798)

*Nonion granosum* (D'ORBIGNY, 1846)

Plate 10, Figure 19

1846 *Nonionia perforata* n. sp. – D'ORBIGNY, p. 110, pl. 5, figs. 17–18.

1965 *Nonion granosus* (D'ORBIGNY) – SOUAYA, p. 326, pl. 3, fig. 5.

1985 *Elphidium granosum* (D'ORBIGNY) – PAPP and SCHMID, p. 46, pl. 36, figs. 6–8.

Description: Test planispiral, nautiloid-like, involute, compressed in cross-section; chambers increasing gradually in size, eighth to nine chambers in final whorl; strongly depressed sutures; wall

calcareous, hyaline; smooth with irregularly arranged tiny nodes, finely perforated; aperture interiomarginal.

*Nonion soldani* (D'ORBIGNY, 1846)  
Plate 10, Figure 20

1846 *Nonionia soldani* n. sp. – D'ORBIGNY, p. 109, pl. 5, figs. 15–16.  
1939 *Nonion soldani* (D'ORBIGNY) – CUSHMAN, p. 13, pl. 3, figs. 10–11.

Description: Test planispiral, nautiloid-like, involute, slightly inflated in cross-section; chambers increasing gradually in size, eighth to nine chambers in final whorl; sutures barely visible; wall calcareous, hyaline; smooth, finely perforated; aperture interiomarginal.

Genus *Nonionella* CUSHMAN, 1926

Type species: *Nonionella miocenica* (CUSHMAN, 1926)

*Nonionella* sp.  
Plate 10, Figure 21

Description: Test planispiral, involute, compressed in cross-section; chambers increasing gradually in size, eighth to nine chambers in final whorl; strongly depressed sutures; wall calcareous, hyaline; smooth, finely perforated; aperture interiomarginal.

Subfamily Pulleniinae SCHWAGER, 1877

Genus *Melonis* DE MONTFORT, 1808

Type species: *Melonis etruscus* DE MONTFORT, 1808

*Melonis* sp. 1  
Plate 10, Figures 22–23

Description: Test globular, inflated, planispiral, circular in lateral view, oval in cross-section; chambers increasing gradually in size, strongly inflated; wall calcareous, hyaline; smooth, finely perforated; aperture interiomarginal.

Genus *Pullenia* PARKER and JONES, 1862

Type species: *Pullenia sphaeroides* (D'ORBIGNY, 1826)

*Pullenia jarvisi* CUSHMAN, 1936  
Plate 10, Figures 24–25

1936 *Pullenia jarvisi* n. sp. – CUSHMAN, p. 77, pl. 13, fig. 6.  
1962 *Pullenia jarvisi* CUSHMAN – HILLEBRANDT, p. 94, pl. 7, figs. 1–3.  
1975 *Pullenia jarvisi* CUSHMAN – PROTO DECIMA and DE BIASE, p. 97, pl. 2, figs. 8a–b.  
1994 *Pullenia jarvisi* CUSHMAN – BOLLI et al., p. 128, pl. 10, fig. 9.  
1997 *Pullenia jarvisi* CUSHMAN – WIDMARK, p. 56, pl. 25, figs. E–F.  
2001 *Pullenia jarvisi* CUSHMAN – ALEGRET and THOMAS, p. 298, pl. 10, fig. 6.  
2006 *Pullenia jarvisi* CUSHMAN – ORTIZ and THOMAS, p. 128, pl. 10, fig. 9.

Description: Test globular, inflated, planispiral, circular in lateral view, oval in cross-section; chambers increasing gradually in size, strongly inflated; sutures distinct, slightly depressed; wall calcareous, hyaline; smooth, finely perforated; aperture interiomarginal.

*Pullenia quinqueloba* (REUSS), 1851  
Plate 10, Figure 26

- 1851 *Nonionia quinqueloba* n. sp. – REUSS, p. 71, pl. 5, fig. 31.  
1926 *Pullenia quinqueloba* (REUSS) – PLUMMER, p. 136, pl. 8, fig. 12.  
1927 *Pullenia quinqueloba* (REUSS) – COLE, p. 32, pl. 5, fig. 15.  
1929 *Pullenia quinqueloba* (REUSS) – GALLOWAY and MORREY, p. 44, pl. 6, fig. 17.  
1941 *Pullenia quinqueloba* (REUSS) – TOULMIN, p. 607, pl. 81, fig. 24.  
1943 *Pullenia quinqueloba* (REUSS) – CUSHMAN and TODD, p. 10, pl. 2, fig. 5; pl. 3, fig. 8.  
1947 *Nonionia quinqueloba* n. sp. – SUBBOTINA, p. 105, pl. 4, fig. 4.  
1949 *Pullenia quinqueloba* (REUSS) – BERMÚDEZ, p. 246, pl. 21, figs. 32–33.  
1951 *Pullenia quinqueloba* (REUSS) – MARKS, p. 69, pl. 7, figs. 19 a–b.  
1951 *Pullenia quinqueloba* (REUSS) – CUSHMAN, p. 59, pl. 17, fig. 6.  
1953 *Pullenia quinqueloba* (REUSS) – BECKMANN, p. 389, pl. 24, figs. 12–13.  
1961 *Pullenia quinqueloba* (REUSS) – HORNIBROOK, p. 90, pl. 11, figs. 207–208.  
1961 *Pullenia quinqueloba* (REUSS) – KAASSCHIETER, p. 202, pl. 202, pl. 11, figs. 1–2.  
1962 *Pullenia quinqueloba* (REUSS) – KIESEL, p. 67, pl. 10, fig. 1.  
1967 *Pullenia quinqueloba* (REUSS) – ROMEO, p. 58, pl. 3, figs. 6a–b.  
1974 *Pullenia quinqueloba* (REUSS) – MURRAY and WRIGHT, p. 120, pl. 18, figs. 13–14.  
1974 *Pullenia quinqueloba* (REUSS) – SZCZECURA and POŻARYSKA, p. 96, pl. 9, fig. 5.  
1975 *Pullenia quinqueloba* (REUSS) – SAMUEL, p. 145, pl. 78, figs. 6–7.  
1983 *Pullenia quinqueloba* (REUSS) – BASOV and KRASHENINNIKOV, p. 766, pl. 14, figs. 10–11.  
1983 *Pullenia quinqueloba* (REUSS) – TJALSMA and LOHMANN, p. 36, pl. 16, fig. 2.  
1987 *Pullenia quinqueloba* (REUSS) – WENGER, p. 299, pl. 13, figs. 16, 20.  
1993 *Pullenia quinqueloba* (REUSS) – DARAKCHIEVA and JURANOV, p. 65, pl. 2, figs. 1, 4.  
1993 *Pullenia quinqueloba* (REUSS) – MATHÉLIN and SZTRÁKOS, p. 82, pl. 19, fig. 17.  
2002 *Pullenia quinqueloba* (REUSS) – KUHN et al., p. 152, pl. 12, figs. 18–20.  
2005 *Pullenia quinqueloba* (REUSS) – NARAYAN et al., p. 133, pl. 4, figs. 31–32.  
2006 *Pullenia quinqueloba* (REUSS) – CIMERMAN et al., p. 38, pl. 10, fig. 10.  
2006 *Pullenia quinqueloba* (REUSS) – ORTIZ and THOMAS, p. 129, pl. 10, fig. 10.

Description: Test globular, inflated, planispiral, circular in lateral view, oval in cross-section; chambers increasing gradually in size, strongly inflated; four distinct sutures, slightly depressed; wall calcareous, hyaline; smooth, finely perforated; aperture interiomarginal.

Family Almaenidae MYATLYUK, 1959  
Subfamily Almaeninae MYATLYUK, 1959

Genus *Almaena* SAMOILOVA, 1940

Type species: *Almaena taurica* SAMOILOVA, 1941

*Almaena* sp.  
Plate 10, Figure 27

Description: Test circular in lateral view, compressed in cross-section, evolute; periphery rounded; chambers increasing gradually in size, eight – nine chambers in final whorl; wall calcareous, hyaline; smooth, coarsely perforated; aperture terminal, on outer edge of individual chambers.

Genus *Queraltina* MARIE, 1950

Type species: *Queraltina epistominoides* Marie, 1950

*Queraltina epistominoides* MARIE, 1950  
Plate 10, Figures 28–30

- 1950 *Queraltina epistominoides* n. sp. – MARIE, p. 74, figs. 1–3, 8–9.  
1983 *Queraltina epistominoides* MARIE – SETIAWAN, p. 133, pl. 15, figs. 2–3.  
1991 *Queraltina epistominoides* MARIE – BARBIN and KELLER-GRÜNIG, p. 241, pl. 2, fig. 16.  
1993 *Queraltina epistominoides* MARIE – SZTRÁKOS and MATHELIN, p. 56, pl. 39, fig. 20.  
2006 *Queraltina epistominoides* MARIE – CIMERMAN et al., p. 38, pl. 10, figs. 11–12.

Description: Test trochospiral, biconvex in cross-section, involute; periphery with relatively thick parallel keels; chambers increasing gradually in size, eighth to nine chambers in final whorl, strongly inflated; wall calcareous, hyaline; smooth, coarsely perforated; aperture terminal, on outer edge of individual chambers.

Superfamily Chilostomellacea BRADY, 1881  
Family Chilostomellidae BRADY, 1881  
Subfamily Chilostomellinae BRADY, 1881

Genus *Chilostomella* REUSS, 1849

Type species: *Chilostomella ovoidea* REUSS, 1850

*Chilostomella tenuis* BORNEMANN, 1855  
Plate 11, Figures 1–2

- 1855 *Chilostomella tenuis* n. sp. – BORNEMANN, p. 343, pl. 17, fig. 2.  
1987 *Chilostomella tenuis* BORNEMANN – SZTRÁKOS, pl. 11, fig. 26.

Description: Test oval, elongate, planispiral, ovate in outline, ovate in cross-section; two chambers in final whorl; invisible suture; wall calcareous, hyaline; smooth, finely perforated; aperture an interiomarginal large slit.

*Chilostomella* sp.  
Plate 11, Figure 3

Description: Test oval, elongate, planispiral, ovate in outline, ovate in cross-section; two chambers in final whorl; invisible suture; wall calcareous, hyaline; smooth, finely perforated; aperture an interiomarginal small slit.

Family Heterolepidae GONZÁLES – DONOSO, 1969

Genus *Anomalinoides* BROTZEN, 1942

Type species: *Anomalinoides plummerae* Brotzen, 1942

*Anomalinoides affinis* (HANTKEN, 1875)  
Plate 11, Figures 4–5

- 1875 *Pulvinulina affinis* n. sp. – HANTKEN, p. 68, pl. 10, fig. 6.  
1934 *Anomalina affinis* (HANTKEN) – CUSHMAN and APPLIN, p. 64, pl. 9, fig. 2.  
1935 *Anomalina affinis* (HANTKEN) – CUSHMAN, p. 51, pl. 21, figs. 11–12.  
1947 *Anomalina affinis* (HANTKEN) – SUBBOTINA, p. 134, pl. 6, figs. 20–25.  
1974 *Anomalinoides affinis* (HANTKEN) – SZCZUCHURA and POŻARYSKA, p. 109, pl. 26, figs. 4–5; pl. 27, figs. 6–7.  
1979 *Anomalina affinis* (HANTKEN) – SZTRÁKOS, pl. 29, fig. 7.  
1982 *Anomalina affinis* (HANTKEN) – SZTRÁKOS, pl. 22, fig. 14.  
1993 *Anomalinoides affinis* (HANTKEN) – DARAKCHIEVA and JURANOV, p. 71, pl. 3, figs. 4–5.  
1993 *Anomalinoides affinis* (HANTKEN) – MATHELIN and SZTRÁKOS, p. 82, pl. 20, fig. 8.

2006 *Anomalinoides affinis* (HANTKEN) – CIMERMAN et al., p. 38, pl. 10, fig. 16.

Description: Test trochospiral, circular in outline, oval, involute, biconvex in cross-section; deep umbilicus; chambers increasing gradually in size; wall calcareous, hyaline; smooth, finely perforated; aperture interiomarginal.

*Anomalinoides alazanensis* (NUTTALL, 1932)

Plate 11, Figure 6

- 1932 *Anomalina alazanensis* n. sp. – NUTTALL, p. 31, pl. 8, figs. 5–7.  
1966 *Anomalinoides alazanensis* (NUTTALL) – BERGGREN and AUBERT, p. 62, pl. 7, fig. 4.  
1983 *Anomalina alazanensis* (NUTTALL) – SETIAWAN, p. 132, pl. 14, fig. 2.  
1986 *Cibicides alazanensis* (NUTTALL) – VAN MORKHOVEN et al., p. 201, pl. 68, figs. 1–2.  
1994 *Anomalinoides alazanensis* (NUTTALL) – BOLLI et al., p. 373, pl. 59, figs 5–9; pl. 79, fig. 20.  
2006 *Anomalinoides alazanensis* (NUTTALL) – ORTIZ and THOMAS, p. 111, pl. 3, figs. 3a–c.

Description: Test low trochospiral, almost planispiral, circular in outline, involute, biconvex in cross-section; chambers increasing gradually in size; strongly depressed sutures; wall calcareous, hyaline; smooth, finely perforated; aperture interiomarginal.

*Anomalinoides grosserugosus* (GÜMBEL, 1868)

Plate 11, Figures 7–8

- 1868 *Truncatulina grosserugosa* n. sp. – GÜMBEL, p. 82, pl. 2, fig. 104.  
1961 *Anomalina grosserugosa* (GÜMBEL) – KAASSCHIETER, p. 217, pl. 12, fig. 14.  
1970 *Anomalina grosserugosa* (GÜMBEL) – LE CALVEZ, p. 198, pl. 40, fig. 2.  
1983 *Anomalina grosserugosa* (GÜMBEL) – SETIAWAN, p. 132, pl. 14, fig. 1.

Description: Test low trochospiral, almost planispiral, circular in outline, involute, biconvex in cross-section; chambers increasing gradually in size, strongly inflated; strongly depressed sutures; wall calcareous, hyaline; smooth, coarsely perforated; aperture interiomarginal.

*Anomalinoides* cf. *A. chiliana* (TODD and KNIKER, 1952)

Plate 11, Figures 9–10

- 1952 *Anomalina chiliana* n. sp. – TODD and KNIKER, p. 27, pl. 4, fig. 34.

Description: Test low trochospiral, almost planispiral, circular in outline, involute, biconvex in cross-section; chambers increasing gradually in size, ten to eleven chambers in final whorl; deep umbilicus; strongly depressed sutures, curved backward; wall calcareous, hyaline; smooth, finely perforated; aperture interiomarginal.

Genus *Heterolepa* FRANZENAU, 1884

Type species: *Heterolepa dutemplei* (D'ORBIGNY, 1846)

*Heterolepa dutemplei* (D'ORBIGNY, 1846)

Plate 11, Figures 11–13

- 1846 *Rotalina dutemplei* n. sp. – D'ORBIGNY, p. 157, pl. 8, figs. 19–21.  
1875 *Truncatulina dutemplei* (D'ORBIGNY) – HANTKEN, p. 71, pl. 8, fig. 5.  
1846 *Rotalina dutemplei* D'ORBIGNY – D'ORBIGNY, p. 157, pl. 8, figs. 19–21.  
1855 *Rotalia bruckneri* n. sp. – REUSS, p. 273, pl. 12, fig. 7.  
1857 *Rotalina dutemplei* D'ORBIGNY – EGGER, p. 274, pl. 7, fig. 8.  
1868 *Rotalia praecincta* n. sp. – KARRER, p. 189, pl. 5, fig. 7.  
1884 *Truncatulina praecincta* (KARRER) – BRADY, p. 667, pl. 95, figs. 1–3.  
1884 *Heterolepa costata* n. sp. – FRANZENAU, p. 216, pl. 5, figs. 2a–c.  
1958 *Cibicides dutemplei* (D'ORBIGNY) – BATJES, p. pl. 9, figs. 9–11.  
1960 *Cibicides dutemplei* (D'ORBIGNY) – CHRISTODOULOU, p. 92, pl. 13, figs. 8a–c.  
1961 *Cibicides dutemplei* (D'ORBIGNY) – KAASSCHIETER, p. 218, pl. 12, fig. 15.

- 1962 *Cibicides dutemplei* (D'ORBIGNY) – KIESEL, p. 73, pl. 11, fig. 1.  
 1964 *Heterolepa dutemplei* (D'ORBIGNY) – LOEBLICH and TAPPAN, p. C758, pl. 623, figs. 3a–c.  
 1966 *Cibicides dutemplei* (D'ORBIGNY) – BUTT, p. 68, pl. 4, figs. 9a–c.  
 1971 *Heterolepa dutemplei* (D'ORBIGNY) – POPESCU and IVA, p. 14, pl. 12, fig. 2.  
 1971 *Heterolepa dutemplei* (D'ORBIGNY) – VERHOEVE, p. 109, pl. 5, figs. 18a–c; pl. 10, fig. 7.  
 1975 *Heterolepa dutemplei* (D'ORBIGNY) – BRAGA et al., p. 109, pl. 6, figs. 1–3.  
 1979 *Cibicides dutemplei* (D'ORBIGNY) – HAGEMAN, p. 91, pl. 3, figs. 5a–b.  
 1982 *Cibicides dutemplei* (D'ORBIGNY) – VAN DER ZWAAN, p. 145, pl. 5, figs. 1a–c and 2a–c.  
 1983 *Cibicides dutemplei* (D'ORBIGNY) – SETIAWAN, p. 126, pl. 11, figs. 4a–c.  
 1984 *Cibicides dutemplei* (D'ORBIGNY) – JONKERS, pl. 4, figs. 2, 3a–b.  
 1985 *Heterolepa dutemplei* (D'ORBIGNY) – GRÜNIG, p. 275, pl. 11, figs. 4–6.  
 1986 *Cibicoides dutemplei* (D'ORBIGNY) – VAN MORKHOVEN et al., p. 112, pl. 35, figs. 1–2.  
 1993 *Heterolepa dutemplei* (D'ORBIGNY) – DARAKCHIEVA and JURANOV, p. 69, pl. 2, figs. 7–8.  
 2000 *Cibicides dutemplei* (D'ORBIGNY) – DEN DULK, pl. 7, figs. 2a–b.  
 2000 *Cibicides dutemplei* (D'ORBIGNY) – KOUWENHOVEN, pl. 2, figs. 2a–c.

Description: Test trochospiral, spiral side evolute, umbilical side involute, biconvex in cross-section; large central umbilicus; chambers increasing gradually in size, eighth to nine chambers in final whorl; sutures barely visible, slightly depressed, curved; wall calcareous, hyaline; smooth, coarsely perforated; aperture interiomarginal, extending to umbilical area.

*Heterolepa simplex* FRANZENAU, 1884  
 Plate 11, Figures 14–15

1884 *Heterolepa simplex* n. sp. – FRANZENAU, p. 215, pl. 5, figs. 1a–c.

Description: Test trochospiral, spiral side evolute, umbilical side involute, biconvex in cross-section; periphery rounded; chambers increasing gradually in size, eighth - nine chambers in final whorl; sutures barely visible, slightly depressed; wall calcareous, hyaline; smooth, coarsely perforated; aperture interiomarginal.

Family Gavelinellidae HOFKER, 1956  
 Subfamily Gyroidinoidinae SAIDOVA, 1981

Genus *Gyroidinoides* BROTZEN, 1942

Type species: *Gyroidinoides nitida* (REUSS, 1845)

*Gyroidinoides dissimilis* (CUSHMAN and RENZ, 1947)  
 Plate 11, Figures 16

- 1947 *Gyroidina dissimilis* n. sp. – CUSHMAN and RENZ, p. 32, pl. 3, fig. 4.  
 1982 *Gyroidinoides dissimilis* (CUSHMAN and RENZ) – SZTRÁKOS, pl. 22, fig. 15.  
 1987 *Gyroidinoides dissimilis* (CUSHMAN and RENZ) – SZTRÁKOS, pl. 12, fig. 17.

Description: Test trochospiral, spiral side evolute, umbilical side involute, triangular, plano-convex in cross-section; chambers increasing gradually in size; sutures barely visible; wall calcareous, hyaline; smooth, coarsely perforated; aperture interiomarginal, extending to umbilical area.

*Gyroidinoides soldanii* (D'ORBIGNY, 1826)  
 Plate 11, Figures 19–21

- 1826 *Rotalina soldanii* n. sp. – (D'ORBIGNY), p. 278, pl. 8, figs. 10–12.  
 1846 *Gyroidina soldanii* (D'ORBIGNY) – D'ORBIGNY, p. 155, pl. 8, figs. 10–12.  
 1953 *Gyroidina soldanii* (D'ORBIGNY) – MYATLYUK, p. 61, pl. 5, figs. 3–5.  
 1958 *Gyroidina soldanii* (D'ORBIGNY) – BATJES, p. 147, pl. 7, figs. 12–13.  
 1975 *Gyroidinoides soldanii* (D'ORBIGNY) – BRAGA et al., p. 109, pl. 6, figs. 10–11.  
 1975 *Gyroidina soldanii* (D'ORBIGNY) – SAMUEL, p. 151, pl. 81, figs. 3–4.

- 1983 *Gyroidinoides soldanii* (D'ORBIGNY) – SETIAWAN, p. 131, pl. 13, fig. 4.  
 1985 *Gyroidinoides soldanii* (D'ORBIGNY) – GRÜNIG, p. 275, pl. 10, figs. 12–14.  
 1987 *Gyroidinoides* ex. gr. *soldanii* (D'ORBIGNY) – SZTRÁKOS, pl. 13, fig. 4.  
 1991 *Gyroidinoides soldanii* (D'ORBIGNY) – CIMERMAN and LANGER, p. 75, pl. 85, figs. 5–6.

Description: Test trochospiral, spiral side evolute, umbilical side involute, plano-convex in cross-section; large central umbilicus; chambers increasing gradually in size, eighth - nine chambers in final whorl; sutures depressed around umbilicus and forming a rosette pattern around umbilical region; wall calcareous, hyaline; smooth, finely perforated; aperture interiomarginal, extending to umbilical area.

*Gyroidinoides* sp.  
 Plate 11, Figure 17

Description: Test trochospiral, spiral side evolute, umbilical side involute, plano-convex to biconvex in cross-section; chambers increasing gradually in size; sutures depressed around umbilicus and forming a rosette pattern around umbilical region; wall calcareous, hyaline; smooth, finely perforated; aperture interiomarginal, extending to umbilical area.

Family Gavelinellinae HOFKER, 1957

Genus *Gavelinella* BROTZEN, 1942

Type species: *Gavelinella pertusa* (MARSSON, 1878)

*Gavelinella micra* (BERMÚDEZ, 1949)  
 Plate 11, Figures 22–23

- 1949 *Cibicides micrus* n. sp. – BERMÚDEZ, p. 302, pl. 24, figs. 34–36.  
 1975 *Gavelinella micra* (BERMÚDEZ) – BRAGA et al., p. 109, pl. 6, fig. 15.  
 1982 *Cibicides micrus* (BERMÚDEZ) – AGIP, pl. 51, figs. 9d–v.  
 1985 *Gavelinella micra* (BERMÚDEZ) – GRÜNIG, p. 275, pl. 10, figs. 23–25.

Description: Test trochospiral in earlier three to four chambers, becoming planispiral in final chambers, test circular in outline, oval in cross-section; periphery rounded with thin keel; large umbilical knob; thick radial, elevated sutures; wall calcareous, hyaline; smooth, coarsely perforated; aperture interiomarginal.

Genus *Hanzawaia* ASANO, 1944

Type species: *Hanzawia nipponica* ASANO, 1944

*Hanzawaia ammophila* (GÜMBEL, 1868)  
 Plate 11, Figures 25–26

- 1868 *Rotalia ammophila* n. sp. – GÜMBEL, p. 652, pl. 2, figs. 90a–b.  
 1930 *Cibicides cushmani* n. sp. – NUTTALL, p. 291, pl. 25, figs. 3, 5–6.  
 1948 *Cibicides cushmani* NUTTALL – CUSHMAN and RENZ, p. 41, pl. 8, figs. 22–23.  
 1949 *Cibicides cushmani* NUTTALL – BERMÚDEZ, p. 297, pl. 26, figs. 4–6.  
 1949 *Cibicides cushmani* NUTTALL – CUSHMAN and STONE, p. 83, pl. 14, fig. 26.  
 1980 *Hanzawaia ammophila* (GÜMBEL) – SAPERSON and JANAL, p. 401, pl. 5, figs. 1–3.  
 1983 *Hanzawaia cushmani* (NUTTALL) – TJALSMA and LOHMANN, p. 32, pl. 17, figs. 1a–c.  
 1983 *Hanzawaia cushmani* (NUTTALL) – MILLER, p. 437, pl. 1, fig. 12.  
 1986 *Hanzawaia ammophila* (GÜMBEL) – VAN MORKHOVEN et al., p. 170, pl. 56, figs. 1–3.  
 1987 *Hanzawaia ammophila* (GÜMBEL) – MILLER and KATZ, p. 134, pl. 6, figs. 3a–b.  
 1992 *Hanzawaia cushmani* (NUTTALL) – MACKENSEN and BERGGREN, p. 620, pl. 3, figs. 1–4.  
 1994 *Cibicidina cushmani* (NUTTALL) – BOLLI et al., p. 370, pl. 57, figs. 15–16.  
 2006 *Hanzawaia ammophila* (GÜMBEL) – ORTIZ and THOMAS, p. 120, pl. 7, figs. 5–6.

Description: Test trochospiral, circular in outline, oval, plano-convex in cross-section, involute umbilical side, evolute spiral side; periphery rounded with thick keel; thick radial, elevated sutures; wall calcareous, hyaline; smooth, coarsely perforated; aperture interiomarginal.

*Hanzawaia producta* (TERQUEM, 1882)  
Plate 11, Figure 18

- 1882 *Truncatulina producta* n. sp. – TERQUEM, p. 92, pl. 9, figs. 20–21.  
1961 *Hanzawaia producta* (TERQUEM) – KAASSCHIETER, p. 266, pl. 13, fig. 13.  
1970 *Hanzawaia producta* (TERQUEM) – KIESEL, p. 3, pl. 21, fig. 2.  
1970 *Hanzawaia producta* (TERQUEM) – LE CALVEZ, p. 201, pl. 44, figs. 3, 9.  
1970 *Hanzawaia producta* (TERQUEM) – NYÍRŐ, p. 201, pl. 44, figs. 3, 9.  
1988 *Hanzawaia producta* (TERQUEM) – HORVÁTH-KOLLÁNYI, p. 93, pl. 29, figs. 3–5.

Description: Test trochospiral, circular in outline, plano-convex in cross-section, involute umbilical side, evolute spiral side; periphery rounded with keel; relatively large umbilicus, slightly compressed sutures; wall calcareous, hyaline; smooth, coarsely perforated; aperture interiomarginal.

Superfamily Orbitoidacea SCHWAGER, 1876  
Family Linderinidae LOEBLICH and TAPPAN, 1984

Genus *Eoannularia* COLE and BERMÚDEZ, 1944

Type species: *Eoannularia eocenica* COLE and BERMÚDEZ, 1944

*Eoannularia eocenica* COLE and BERMÚDEZ, 1944  
Plate 11, Figure 24

- 1944 *Eoannularia eocenica* n. sp. – COLE and BERMÚDEZ, p. 342, pl. 24, figs. 12–14.  
1969 *Eoannularia eocenica* COLE and BERMÚDEZ – ZILAHY, p. 158, pl. 7, figs. 10–13.  
1976 *Eoannularia eocenica* COLE and BERMÚDEZ – SIREL, p. 80, pl. 1, figs. 1–3, pl. 2, 1–2, 4–5, 8–9.  
1988 *Eoannularia eocenica* COLE and BERMÚDEZ – HORVÁTH-KOLLÁNYI, p. 83, pl. 23, figs. 1–2.  
2006 *Eoannularia eocenica* COLE and BERMÚDEZ – CIMERMAN et al., p. 40, pl. 12, fig. 3.

Description: Test small, circular in outline, strongly compressed in cross-section; large umbilicus; wall formed by calcareous meshwork.

Superfamily Rotaliacea EHRENBERG, 1839  
Family Rotaliidae EHRENBERG, 1839  
Subfamily Pararotaliinae REISS, 1963

Genus *Pararotalia* LE CALVEZ, 1949

Type species: *Pararotalia inermis* (TERQUEM, 1882)

*Pararotalia inermis* (TERQUEM, 1882)  
Plate 11, Figures 27–28

- 1882 *Rotalina inermis* n. sp. – TERQUEM, p. 38, pl. 6, fig. 1.  
1970 *Pararotalia inermis* (TERQUEM) – LE CALVEZ, p. 163, pl. 34, figs. 6–7

Description: Test trochospiral, evolute, circular in outline, biconvex in cross-section; periphery undulate with broad keel; deep, distinct umbilicus surrounded by numerous large-sized nodes; sutures strongly depressed; wall calcareous, hyaline; smooth, finely perforated; aperture interiomarginal, extending to umbilical area.



Genus *Rotalia* LAMARCK, 1804

Type species: *Rotalia trochidiformis* (LAMARCK, 1804)

*Rotalia* sp. cf. *R. calcar* (D'ORBIGNY, 1826)  
Plate 11, Figure 29

1826 *Calcarina calcar* n. sp. – D'ORBIGNY, p. 276, pl. 5, figs. 22–24.

Description: Test trochospiral, circular in outline, involute on umbilical side, evolute on spiral side, biconvex on cross-section; periphery acute, star-shaped; chambers increasing gradually in size; sutures distinct, depressed; wall calcareous, hyaline; smooth, finely perforated; aperture interiomarginal, extending to umbilical area.

*Rotalia* sp.  
Plate 11, Figure 30

Description: Test trochospiral, circular in outline, involute on umbilical side, evolute on spiral side, convex-plane in cross-section; umbilicus filled up by coarse grains; chambers increasing gradually in size, seven to eighth chambers in final whorl; sutures distinct, depressed; wall calcareous, hyaline; smooth, coarsely perforated; aperture interiomarginal, extending to umbilical area.

Subfamily Ammoniinae SAIDOVA, 1981

Genus *Ammonia* BRÜNNICH, 1772

Type species: *Ammonia beccari* (LINNÉ, 1758)

*Ammonia* sp.  
Plate 11, Figure 31

Description: Test trochospiral, circular in outline, biconvex in cross-section; periphery acute, star-shaped; chambers increasing gradually in size; sutures distinct, depressed; wall calcareous, hyaline; smooth, finely perforated; aperture interiomarginal, extending to umbilical area.

Family Elphidiidae Galloway, 1933  
Subfamily Elphidiinae GALLOWAY, 1933

Genus *Elphidium* DE MONTFORT, 1808

Type species: *Elphidium macellum* (FICHTEL and MOLL, 1798)

*Elphidium* sp. cf. *E. laeve* (D'ORBIGNY, 1826)  
Plate 11, Figures 32–33

1826 *Nonionina laevis* n. sp. – D'ORBIGNY, p. 294, fig. 46.  
1970 *Elphidium laeve* (D'ORBIGNY) – LE CALVEZ, p. 168, pl. 25, fig. 1.

Description: Test planispiral, involute, circular in outline, oval, slightly compressed in cross-section; periphery rounded with thin keel; deep circular umbilicus; chambers increasing gradually in size; sutures strongly depressed; wall calcareous, hyaline; smooth, coarsely perforated; aperture interiomarginal, extending to umbilical area.

Table 1. Geographical coordinates, total thickness of Eocene sequences, number of samples and defined age of the investigated boreholes and outcrops

Boreholes and outcrops	Code	Latitude	Longitude	Total thickness of Eocene sequences (m)	Number of samples	Age (Nannoplankton Zone)
Csetény 61	Cst-61	47°18'48.48"	18°00'55.72"	249	48	NP16-NP19
Bakonycsernye 18	Bkcs-18	47°18'12.33"	18°04'58.64"	90	18	NP16-NP17
Padrag 5	Pa-5	47°03'27.57"	17°32'52.76"	256	13	NP15?-NP18
Somlóvásárhely 1	Sv-1	47°05'00.70"	17°23'24.39"	294	207	NP14-NP19
Devecser 4	Dv-4	47°07'09.31"	17°28'40.59"	236	83	NP15?-NP17
Bakonyszentkirály 3	Bszk-3	47°21'53.54"	17°56'05.85"	178	40	NP16-NP18
Balinka 285	Ba-285	47°21'36.99"	18°10'50.57"	98	31	NP16-NP17
Csákberény 89	Csbr-89	47°20'03.88"	18°20'31.18"	197	76	NP16
Dudar 240	D-240	47°18'41.75"	17°58'34.08"	169	37	NP16-NP19
Halimba 1	Hgy-1	47°03'20.76"	17°32'31.74"	54	34	NP15?-NP16
Tarján 13	Tj-13	47°36'13.49"	18°32'52.52"	43	15	NP16-NP17
Tarján 14	Tj-14	47°35'56.43"	18°33'22.25"	86	19	NP16-NP17
Csordakút	Csk.	47°35'05.36"	18°32'31.97"	35	22	NP16-NP17
Mátyás-hegy	Mh.	47°32'01.33"	19°01'04.57"	21	12	NP18-NP19

Table 2. Presence/absence of foraminifer species in Eocene profiles of the Hungarian Paleogene Basin.

Boreholes	Species (Nannoplankton zones)	Mh		Tj-14		Tj-13		Csk		Csbr-89		Bszk-3			Cst-61			D-240			Bkcs-18			Ba-285			Sv-1			Pa-5			Hgy-1		Dv-4				
		NP 19	NP 18	?NP17	NP 16	?NP 17	NP 16	NP 17	NP 16	NP 16	NP 16	NP 18	NP 17	NP 16	NP 19	NP 18	NP 17	NP 16	NP 19	NP 18	NP 17	NP 16	NP 19	NP 18	NP 17	NP 16	NP 19	NP 18	NP 17	NP 16	NP 15	NP 14	NP 18	NP 16	NP 17				
<i>Acervulina</i> sp.																																							
<i>Adelosina</i> sp.																																							
<i>Almaena</i> sp.											+																												
<i>Ammomarginulina</i> sp.																																							
<i>Ammonia</i> sp.																																							
<i>Anomalinooides affinis</i>																																							
<i>Anomalinooides alazanensis</i>																																							
<i>Anomalinooides cf. chileana</i>																																							
<i>Anomalinooides grosserugosa</i>																																							
<i>Articulina curta</i>																																							
<i>Articulina laevigata</i>																																							
<i>Articulina nitida</i>																																							
<i>Asterigerina</i> sp.																																							
<i>Astrorhiza bakonycernyensis n. sp</i>																																							
<i>Bathysiphon eocenicus</i>																																							
<i>Bathysiphon saidi</i>																																							
<i>Bolivina cookei</i>																																							
<i>Bolivina elongata</i>																																							
<i>Bolivina nobilis</i>																																							
<i>Bolivina semistriata</i>																																							
<i>Bolivina</i> sp.																																							
<i>Bolivinopsis foliacea</i>																																							
<i>Bulimina affinis</i>																																							
<i>Bulimina parisensis</i>																																							
<i>Bulimina</i> sp.																																							
<i>Bulimina truncana</i>																																							
<i>Cancris</i> sp.																																							
<i>Chilostomella tenuis</i>																																							
<i>Chilostomella</i> sp.																																							
<i>Chrysalogonium tympanipectiformis</i>																																							
<i>Chrysalogonium</i> sp.																																							
<i>Cibicides ammophilus</i>																																							



Boreholes	Mh		Tj-14		Tj-13		Csk		Csdr-89		Bszk-3		Cst-61		D-240		Bkes-18		Ba-285		Sv-1		Pa-5		Hgy-1		Dv-4			
	NP 19	NP 18	?NP17	NP 16	?NP 17	NP 16	NP 17	NP 16	NP 16	NP 18	NP 17	NP 16	NP 19	NP 18	NP 17	NP 16	NP 17	NP 16	NP 17	NP 16	NP 19	NP 18	NP 17	NP 16	NP 18	NP 16	NP 17	NP 16	NP 17	
<i>Dentalina roemeri</i>																														
<i>Dentalina</i> sp.	+																													
<i>Dentalina subtilis</i>																														
<i>Discorbis elegans</i>																														
<i>Discorbis parisiensis</i>																														
<i>Discorbis perplexa</i>																														
<i>Discorinopsis</i> sp.																														
<i>Dorothia textilaroides</i>																														
<i>Dyocibicides uniserialis</i>																														
<i>Ellipsoglandulina multicosata</i>																														
<i>Elphidium</i> cf. <i>laeve</i>																														
<i>Eoannularia eocenica</i>																														
<i>Eponides budensis</i>																														
<i>Eponides haidingeri</i>																														
<i>Eponides polyganus</i>																														
<i>Eponides</i> sp.																														
<i>Eponides umbonatus</i>																														
<i>Fissurina</i> cf. <i>praeclara</i>																														
<i>Fissurina orbignyana</i>																														
<i>Fissurina</i> sp.																														
<i>Fissurina tricincta</i>																														
<i>Fronicularia budensis</i>																														
<i>Fronicularia semicosta</i>																														
<i>Fronicularia tenuissima</i>																														
<i>Fursenkoina hungarica</i>																														
<i>Fursenkoina schreibersiana</i>																														
<i>Gavelinella micra</i>																														
<i>Glandulina aequalis</i>																														
<i>Glandulina hantkeni</i>																														
<i>Glandulina inflata</i>																														
<i>Glandulina obtusissima</i>																														
<i>Glandulina</i> sp.																														
<i>Globocassidulina globosa</i>																														
<i>Globulina gibba</i>																														
<i>Globulina guttula</i>																														
<i>Globulina minuta</i>																														

Boreholes	Mh		Tj-14		Tj-13	Csk	Csbr-89		Bszk-3		Cst-61		D-240		Bkes-18		Ba-285	Sv-1		Pa-5	Hgy-1		Dv-4			
	NP 19	NP 18	?NP17	NP 16	?NP17		NP 16	NP 17	NP 16	NP 17	NP 18	NP 19	NP 16	NP 17	NP 18	NP 19		NP 16	NP 17		NP 18	NP 19		NP 14	NP 15	NP 16
<i>Gonatosphaera inflata</i>																										
<i>Guttulina irregularis</i>																										
<i>Gyroidinoides dissimilis</i>																										
<i>Gyroidinoides soldanii</i>																										
<i>Gyroidinoides koestleri</i>																										
<i>Gyroidionides</i> sp.																										
<i>Hanzawaia ammophila</i>																										
<i>Hanzawaia producta</i>																										
<i>Haplophragmoides</i> sp.																										
<i>Heterolepa dutemplei</i>																										
<i>Heterolepa simplex</i>																										
<i>Lagena clava</i>																										
<i>Lagena globosa</i>																										
<i>Lagena hexagona</i>																										
<i>Lagena laevis</i>																										
<i>Lagena sulcata</i>																										
<i>Lagena tenuis ornata</i>																										
<i>Lenticulina arcuatostrata</i>																										
<i>Lenticulina excisa</i>																										
<i>Lenticulina falcifer</i>																										
<i>Lenticulina granulata</i>																										
<i>Lenticulina gutticostata</i>																										
<i>Lenticulina platyptera</i>																										
<i>Lenticulina</i> sp.																										
<i>Marginulina behmi</i>																										
<i>Marginulina fragaria texasensis</i>																										
<i>Marginulina hantkeni</i>																										
<i>Marginulina pediformis</i>																										
<i>Marginulina propinqua</i>																										
<i>Marginulina</i> sp.																										
<i>Marginulina tumida</i>																										
<i>Marginulinopsis porvaensis</i>																										
<i>Marginulinopsis</i> sp.																										
<i>Martinottiella rhumbleri</i>																										
<i>Massilina</i> sp.																										
<i>Melonis</i> sp. 1																										

Boreholes	Mh		Tj-14		Tj-13	Csk	Csdt-89		Bszk-3		Cst-61		D-240		Bkes-18		Ba-285	Sv-1					Pa-5	Hgy-1		Dv-4	
	NP 19	NP 18	?NP17	NP 16	NP 16		NP 16	NP 16	NP 17	NP 16	NP 17	NP 18	NP 19	NP 16	NP 17	NP 16		NP 17	NP 18	NP 19	NP 16	NP 17		NP 18	NP 19		NP 16
<i>Miliola prisca</i>								+																			
<i>Miliola strigillata</i>								+																			
<i>Nodosarella lorifera</i>																											
<i>Nodosarella sp.</i>																											
<i>Nodosarella tuberosa</i>																											
<i>Nodosaria affinis</i>																											
<i>Nodosaria badenensis</i>																											
<i>Nodosaria crassa</i>																											
<i>Nodosaria elegans</i>																											
<i>Nodosaria exilis</i>																											
<i>Nodosaria intermedia</i>																											
<i>Nodosaria longiscata</i>																											
<i>Nodosaria pyrula</i>																											
<i>Nodosaria radricula</i>																											
<i>Nodosaria sp.</i>																											
<i>Nonion affinae</i>																											
<i>Nonion boueanum</i>																											
<i>Nonion granosum</i>																											
<i>Nonion scaphum</i>																											
<i>Nonion soldani</i>																											
<i>Nonionella sp.</i>																											
<i>Nonionella wemmelensis</i>																											
<i>Nuttalides sp.</i>																											
<i>Orthomorphina rohri</i>																											
<i>Pararotalia curry</i>																											
<i>Pararotalia inermis</i>																											
<i>Planularia sp. 1</i>																											
<i>Planularia sp. 2</i>																											
<i>Planulina austriaca</i>																											
<i>Planulina compressa</i>																											
<i>Planulina costata</i>																											
<i>Planulina sp.</i>																											
<i>Plectina eocenica</i>																											
<i>Pleurostomella acuta</i>																											
<i>Pleurostomella alternans</i>																											
<i>Pleurostomella eocaena</i>																											

Boreholes	Mh		Tj-14		Tj-13		Csk		Csdr-89		Bszk-3		Cst-61		D-240		Bkes-18		Ba-285		Sv-1		Pa-5		Hgy-1		Dv-4				
	NP 19	NP 18	?NP17	NP 16	?NP17	NP 16	NP 17	NP 16	NP 16	NP 18	NP 17	NP 16	NP 19	NP 18	NP 17	NP 16	NP 17	NP 16	NP 17	NP 16	NP 19	NP 18	NP 17	NP 16	NP 15	NP 14	NP18	NP 16	?NP 15	NP 17	
<i>Pleurostomella incrassata</i>																														+	
<i>Pleurostomella</i> sp.																															
<i>Pseudonodosaria discreta</i>																			+												
<i>Pullenia jarvisi</i>																															+
<i>Pullenia quinqueloba</i>																															+
<i>Pyrgo bulloides</i>																															
<i>Pyrgo simplex</i>																															
<i>Pyralina</i> sp.																															
<i>Queralina epistominoides</i>																															
<i>Quinqueloculina bicarinata</i>																															
<i>Quinqueloculina buchiana</i>																															
<i>Quinqueloculina carinata</i>																															
<i>Quinqueloculina affig. carinata</i>																															
<i>Quinqueloculina cf. contorta</i>																															
<i>Quinqueloculina juleana</i>																															
<i>Quinqueloculina seminula</i>																															
<i>Quinqueloculina ungeriana</i>																															
<i>Quinqueloculina</i> sp. 1																															
<i>Quinqueloculina</i> sp. 2																															
<i>Ramulina</i> sp.																															
<i>Reophax harrisi</i>																															
<i>Reussella elongata</i>																															
<i>Reussella</i> sp.																															
<i>Reussella terquemi</i>																															
<i>Rhabdammina abyssorum</i>																															
<i>Rotalia</i> cf. <i>R. calcar</i>																															
<i>Rotalia trochidiformis</i>																															
<i>Rotalia</i> sp.																															
<i>Saracenaria hantkeni</i>																															
<i>Sphaerogypsina globula</i>																															
<i>Spirolina mariei</i>																															
<i>Spirolina pedum</i>																															
<i>Spirolina</i> sp.																															
<i>Spiroloculina bicarinata</i>																															
<i>Spiroloculina jarvisi</i>																															
<i>Spiroloculina obscura</i>																															



Boreholes	Mh		Tj-14		Tj-13	Csk	Csbr-89		Bszk-3		Cst-61			D-240			Bkes-18		Ba-285	Sv-1					Pa-5	Hgy-1		Dv-4
	NP 19	NP 18	?NP17	NP 16	NP 16		NP 17	NP 16	NP 16	NP 17	NP 18	NP 19	NP 16	NP 17	NP 18	NP 19	NP 16	NP 17		NP 16	NP 17	NP 18	NP 19	NP 16		NP 17	NP 18	
<i>Spiroplectammina carinata</i>	+	+								+											+					+		
<i>Spiroplectammina subhaeringensis</i>	+	+																										
<i>Stilostomella abyssorum</i>	+									+																		
<i>Stilostomella adolphina</i>	+									+																		
<i>Stilostomella consobrina</i>								+																+				
<i>Stilostomella curvatura</i>	+																											
<i>Stilostomella elegans</i>	+																		+									
<i>Stilostomella emaciata</i>	+																											
<i>Stilostomella hoernesii</i>	+																											
<i>Stilostomella pauperata</i>																												
<i>Stilostomella</i> sp.	+																											
<i>Textularia cf. partschii</i>																												
<i>Textularia crookshanki</i>																												
<i>Textularia deperdita</i>																												
<i>Textularia globosa</i>	+																											
<i>Textularia halkyardi</i>																												
<i>Textularia lanceolata</i>																												
<i>Textularia pala</i>																												
<i>Textularia</i> sp. 1.	+																											
<i>Textularia</i> sp. 2.																												
<i>Textularia</i> sp. 3	+																											
<i>Triloculina gibba</i>																												
<i>Triloculina porvaensis</i>																												
<i>Triloculina</i> sp.																												
<i>Triloculina trigonula</i>																												
<i>Tritaxilina pupa</i>	+																											
<i>Tritaxilina</i> sp.	+																											
<i>Uvigerina chirana</i>	+																											
<i>Uvigerina cocoensis</i>																												
<i>Uvigerina cocoensis jacksonis</i>																												
<i>Uvigerina eocaena</i>	+																											
<i>Uvigerina gallowayi</i>	+																											
<i>Uvigerina gracilis</i>																												
<i>Uvigerina hantkeni</i>																												
<i>Uvigerina hourcqi</i>																												
<i>Uvigerina multistriata</i>	+																											

Boreholes	Species (Nannoplankton zones)		MH		Tj-14		Tj-13		Csk		Csdr-89		Bszk-3		Cst-61		D-240		Bkes-18		Ba-285		Sv-1		Pa-5		Hgy-1		Dv-4								
	NP 19	NP 18	?NP17	NP 16	?NP17	NP 16	NP 16	NP 17	NP 16	NP 16	NP 18	NP 17	NP 16	NP 19	NP 18	NP 17	NP 16	NP 17	NP 16	NP 17	NP 16	NP 19	NP 18	NP 17	NP 16	NP 15	NP 14	NP 18	NP 16	?NP 15	NP 17						
<i>Uvigerina pigma</i>				+																																	
<i>Uvigerina rippensis</i>				+																																	
<i>Uvigerina tenustriata</i>				+																																	
<i>Vaginulina legumen</i>																																					
<i>Vaginulina</i> sp. cf. <i>V. ex gr. mexicana</i>																																					
<i>Vaginulinopsis fragaria</i>																																					
<i>Vaginulinopsis hagni</i> n. sp.																																					
<i>Vaginulinopsis hantkeni</i>																																					
<i>Vaginulinopsis minimus</i>																																					
<i>Vaginulinopsis pseudodecorata</i>																																					
<i>Vaginulinopsis</i> sp.																																					
<i>Valvulineria</i> sp.																																					
<i>Verneuilina</i> sp.																																					
<i>Vulvulina advena</i>																																					
<i>Vulvulina jarvisi</i>																																					

## References

- AGIP S.p.A., 1982. Foraminiferi Padani (Terziario e Quaternario). AGIP SpA San Donato Milanese
- ALEGRET, L. and THOMAS, E. 2001. Upper Cretaceous and lower Paleogene benthic foraminifera from northeastern Mexico. *Micropaleontology* 47: 269–316.
- ANAN, H. S. 1994. Benthic foraminifera around Middle/Upper Eocene boundary in Egypt. Middle East Research Center, Ain Shams University, Earth Science Series, Cairo 8: 210–233.
- ANAN, H. S. 2005. Agglutinated Middle-Upper Eocene foraminifera in Jabal Hafit, Al Ain area, United Arab Emirates. *Revue de Paléobiologie* 24 (1): 17–27.
- ANDERSEN, H. V. 1961. Genesis and paleontology of the Mississippi river mudlumps, Part III: foraminifera of the mudlumps, lower Mississippi river delta. Louisiana Department of Conservation, Geological Bulletin 35: 1–208.
- BÁLDI, T. and BÁLDI-BEKE, M. 1985. The evolution of the Hungarian Paleogene Basins. *Acta Geologica Hungarica* 28: 5–28.
- BÁLDI, T. and BÁLDI-BEKE, M. 1991. Paleobathymetry and paleogeography of the Bakony Eocene basin in western Hungary. *Palaeogeography, Palaeoclimatology, Palaeoecology* 88: 25–52.
- BÁLDI-BEKE, M. 1972. The nannoplankton of the Upper Eocene bryozoan and Buda marl. – *Acta Geologica Hungarica* 16: 211–228.
- BÁLDI-BEKE, M. 1977. Stratigraphical and faciological subdivisions of the Oligocene as based on nannoplankton. *Földtani Közlöny [Bulletin of the Hungarian Geological Society]* 107: 59–89.
- BÁLDI-BEKE, M. 1984. The nannoplankton of the Transdanubian Palaeogene Formations. – *Geologica Hungarica, series Palaeontologica* 43: 1–307.
- BANDY, O. L. 1949. Eocene and Oligocene foraminifera from Little Stave, Clarke County, Alabama. *Bulletin of American Paleontology* 32 (131): 1–210.
- BARBIN, V. and KELLER, G. A. 1991. Benthic foraminiferal assemblages from the Brendola section (Priabonian stage stratotype area, northern Italy); distribution, palaeoenvironment and palaeoecology. *Marine Micropaleontology* 17(3–4): 237–254.
- BARKER, R. W. 1960. Taxonomic notes on the species figured by H. B. Brady in his report on the foraminifera dredged by the HMS Challenger during the years 1873–1876. *Society of Economic Paleontologists and Mineralogists, Special Publication* 9: 7–24.
- BASOV, I. A. and KRASHENINNIKOV, V. A. 1983. Benthic foraminifers in Mesozoic and Cenozoic sediments of the southwestern Atlantic as an indicator of paleoenvironment, Deep Sea Drilling Project Leg 71 In: LUDWIG, W. J., KRASHENINNIKOV, V. A. et al., Initial Reports DSDP 71: 739–788.
- BATJES, D. A. J. 1958. Foraminifera of the Oligocene of Belgium. *Institut Royal des Sciences Naturelles de Belgique, Mémoires* 143: 1–188.
- BECKMANN, J. P. 1953. Die Foraminiferen der Oceanic Formation (Eocæn-Oligocæn) von Barbados, Kl. Antillen. *Eclogae Geologicae Helvetiae* 46 (2): 301–412.
- BERGGREN, W. A. and AUBERT, J. 1976. Late Paleogene (Late Eocene and Oligocene) benthonic foraminiferal stratigraphy and paleobathymetry of Rockall Bank and Hatton-Rockall Basin. *Micropaleontology* 22: 307–326.
- BERMÚDEZ, P. 1949. Tertiary smaller foraminifera of the Dominican Republic. *Cushman Laboratory for Foraminiferal Research, Special Publication* 25: 1–322.
- BERNHARDT, B., BÁLDI-BEKE, M., LANTOS, M., HORVÁTH-KOLLÁNYI, K. and MÁRTON, P. 1988. Eocene magneto- and biostratigraphy at Somlóvásárhely, Hungary. *Acta Geologica Hungarica* 31: 33–52.
- BHATIA, S. B. 1955. The foraminiferal fauna of the late Palaeogene sediments of the Isle of Wight, England. *Journal of Palaeontology* 29(4): 665–693.
- BIGNOT, G. 1998. Middle Eocene benthic foraminifers from holes 960A and 960C, Central Atlantic Ocean In: MASCLE, J., LOHMANN, G. P. and MOULLADE, M. (Eds.): *Proceedings of the Ocean Drilling Program, Scientific Results* 159: 433–444.
- BOERSMA, A. 1984. *A Handbook of Common Tertiary Uvigerina*: Stony Point, NY (Microclimates Press): 1–207.
- BOLLI, H. M., BECKMANN, J. P. and SAUNDERS, J. B. 1994. *Benthic Foraminiferal Biostratigraphy of the South Caribbean region*. Cambridge University Press: 1–408.
- BORNEMANN, J. G. 1855. Die mikroskopische Fauna des Septarienthones von Hermsdorf bei Berlin. *Geologische Gesellschaft Zeitschrift* 7: 307–371.
- BORNMALM, L. 1997. Taxonomy and paleoecology of late Neogene benthic foraminifera from the Caribbean Sea and eastern equatorial Pacific Ocean. *Fossils and Strata* 41: 1–96.
- BRADY, H. B. 1881. Notes on some of the reticularian Rhizopoda of the “Challenger” Expedition; Part III. *Quarterly Journal of the Microscopical Society* 21: 31–71.

- BRADY, H. B. 1884. Report on the Foraminifera dredged by HMS Challenger, during the years 1873–1876. Reports, Scientific Results of the Challenger Expedition, Zoology 9: 1–814.
- BRAGA, G., DE BIASE, R., GRÜNIG, A. and PROTO DECIMA, F. 1975. Foraminiferi bentonici del Paleocene ed Eocene della sezione di Possagno. Schweizerische Palaeontologische Abhandlungen 97: 85–111.
- BRAGA, G. and GRÜNIG, A. 1975. Foraminiferi bentonici dell'Eocene superiore In: BRAGA et al. 1975. Foraminiferi bentonici del Paleocene ed Eocene della sezione di Possagno. Schweizerische Palaeontologische Abhandlungen 97: 98–111.
- BREMER, M. L., BRISKIN, M., BERGGREN, W. A. 1980. Quantitative paleobathymetry and paleoecology of the Late Pliocene-Early Pleistocene foraminifera of Le Castella (Calabria, Italy). Journal of Foraminiferal Research 10 (1): 1–30.
- BROLSMA, M. J. 1978. Quantitative foraminiferal analysis and environmental interpretation of the Pliocene and topmost Miocene on the south coast of Sicily. Utrecht Micropaleontological Bulletins 18: 1–159.
- BROTZEN, F. 1948. The Swedish Paleocene and its foraminiferal fauna. Sveriges Geologiska Undersökning 4: 19–52.
- BUTT, A. A. 1966. Late Oligocene foraminifera from Escornebéou, SW France. Schotanus & Jens, Utrecht: 1–123.
- CARPENTER, W. B. 1869. On the rhizopodal fauna of the deep sea. Proceedings of the Royal Society of London 18: 59–62.
- CARPENTER, W. B. 1881. The Microscope and its Revelations (6th edition) J. and A. Churchill, London: 1–882.
- CHAPMAN, F. 1926. Cretaceous and Tertiary Foraminifera of New Zealand. Palaeontological Bulletin of New Zealand Geological Survey Branch 11: 1–119.
- CHRISTODOULOU, G. 1960. Geologische und mikropaläontologische Untersuchung auf der Insel Karpathos (Dodekanes). Palaeontographica, Abteilung A 115: 1–143.
- CICHA, I., KRHOVSKÝ, J., BRZOBOHATÝ, R., ČTYROKA, J., VON DANIELS, C. H., HANNOLD, T., HORVÁTH, M., LUCZKOWSKA, E., REIZER, H., RUPP, CH., RIJAVEC, L. and WENGER, W. 1986. Oligocene and Miocene from the western and central Paratethys. Utrecht Micropaleontological Bulletin 35: 121–182.
- CIMERMAN, F., JELEN, B. and ŠKABERNE, D. 2006. Late Eocene benthic foraminiferal fauna from clastic sequence of the Socka. Geologija 49(1): 7–44.
- CIMERMAN, F. and LANGER, M. R. 1991. Mediterranean Foraminifera. Slovenska Akademija Znanosti, Ljubljana: 1–118.
- CITA, M. B. 1950. L'eocene della sponda occidentale del lago di Garda. Rivista Italiana di Paleontologia 56(3): 81–105.
- COLE, W. S. 1927. A foraminiferal fauna from Guayabal formation in Mexico. **Bulletins of American Paleontology** 14(51): 1–46.
- COLE, W. S. 1928. A foraminiferal fauna from the Chapopote formation in Mexico. **Bulletins of American Paleontology** 14(53): 202–232.
- COLE, W. S. and BERMÚDEZ, P. J. 1944. New foraminiferal genera from the Cuban Middle Eocene. **Bulletins of American Paleontology**, 28(113): 333–344.
- COSTA, A. 1856. Paleontologia del Regno di Napoli. Accad Pontaniana Napoli 7: 113–378.
- COLOM, G. 1954 - Estudio de las biozonas con foraminiferos del Terciario de Alicante. Inst. Geol. Min. España, Bol., vol. 66, pp. 1-279
- CUSHMAN, J. A. 1918. Some Miocene foraminifera of the coastal plain of the United States. U.S. Geological Survey, Bulletin: 1–676.
- CUSHMAN, J. A. 1921. Foraminifera of the Atlantic Ocean: Part 3 Textulariidae. Bulletin of the Natural History Museum, 104(3): 1–149.
- CUSHMAN, J. A. 1925. Some Textulariidae from the Miocene of California. Contributions from the Cushman Laboratory for Foraminiferal Research, 1(2): 29–35
- CUSHMAN, J. A. 1925a. Eocene foraminifera from the Cocoa Sand of Alabama. Contributions from the Cushman Laboratory for Foraminiferal Research 1(16): 65–69.
- CUSHMAN, J. A. 1926. The Foraminifera of the Velasco shale of the Tampico Embayment. American Association of Petroleum Geologists Bulletin, 10: 581–612.
- CUSHMAN, J. A. 1927. New and interesting foraminifera from Mexico and Texas. Contributions from the Cushman Laboratory for Foraminiferal Research 3(2): 111–117.
- CUSHMAN, J. A. 1928. Foraminifera, their classification and economic use. Cushman Laboratory for Foraminiferal Research Special Publication 1: 1–404.
- CUSHMAN, J. A. 1929. A late Tertiary fauna of Venezuela and other related regions Contributions from the Cushman Laboratory for Foraminiferal Research 5: 1–94.
- CUSHMAN, J. A. 1931. The foraminifera of the Atlantic Ocean, 8: Rotaliidae, Amphisteginidae, Calcarinidae, Cymbaloporetidae, Globorotaliidae, Anomalinidae, Planorbulinidae, Rupertiidae and Homotremidae. United States National Museum Bulletin 104: 1–144.

- CUSHMAN, J. A. 1932. The genus *Vulvulina* and its species. Contributions from the Cushman Laboratory for Foraminiferal Research 8(4): 1–84.
- CUSHMAN, J. A. 1933. New Foraminifera from the Upper Jackson Eocene of the southeastern Coastal Plain region of the United States. Contributions from the Cushman Laboratory for Foraminiferal Research, 9: 1–21.
- CUSHMAN, J. A. 1935. Upper Eocene foraminifera of the southeastern United States. United States Geological Survey, Professional Paper 181:1–88.
- CUSHMAN, J. A. 1936. Some American Cretaceous species of *Ellipsonodosaria* and *Chrysalogonium*. Contributions from the Cushman Laboratory for Foraminiferal Research, 12(3): 51–55.
- CUSHMAN, J. A. 1937. A monograph of the Foraminiferal family Valvulinidae. Cushman Laboratory for Foraminiferal Research, Special Publication 8: 1–209.
- CUSHMAN, J. A. 1939. A monograph of the foraminiferal family Buliminidae. Cushman Laboratory for Foraminiferal Research, Special Publication, 9: 1–228.
- CUSHMAN, J. A. 1945. The species of the subfamily Reussellinae of the foraminiferal family Buliminidae. Cushman Laboratory for Foraminiferal Research, Special Publication 21: 6–54.
- CUSHMAN, J. A. 1946. Upper Cretaceous Foraminifera of the Gulf Coastal Region of the United States and adjacent areas. United States Geological Survey, Professional Paper 206: 1–241.
- CUSHMAN, J. A. 1951. Paleocene foraminifera of the Gulf coastal Region of the United States and adjacent areas. United States Geological Survey, Professional Paper 232:1–75.
- CUSHMAN, J. A. and APPLIN, E R 1926. Texas Jackson Foraminifera. American Association of Petroleum Geologists Bulletin 10: 154–189.
- CUSHMAN, J. A. and BERMÚDEZ, P. J. 1937. Further new species of foraminifera from the Eocene of Cuba. Contributions from the Cushman Laboratory for Foraminiferal Research 13: 1–29.
- CUSHMAN, J. A. and HANNA, G D 1927. Foraminifera from the Eocene near Coalinga. California Academy of Sciences 16(8): 205–229.
- CUSHMAN, J. A. and EDWARDS, P G 1937. Notes on the early described Eocene species of *Uvigerina* and some new species. Contributions from the Cushman Laboratory for Foraminiferal Research 13(2): 54–61.
- CUSHMAN, J. A. and JARVIS, P W 1934. Some interesting new uniserial foraminifera from Trinidad. Contributions from the Cushman Laboratory for Foraminiferal Research 10(3): 71–75.
- CUSHMAN, J. A. and MCMASTERS, J H 1936. Middle Eocene Foraminifera from the Llajas Formation, Ventura County, California. Journal of Paleontology 10: 497–517
- CUSHMAN, J. A. and RENZ, H. H. 1946. The foraminiferal fauna of the Lizard Springs Formation of Trinidad, British West Indies. Cushman Laboratory for Foraminiferal Research, Special Publication 18: 1–48.
- CUSHMAN, J. A. and RENZ, H. H. 1947. The foraminiferal fauna of the Oligocene, Ste Croix Formation of Trinidad. Cushman Laboratory for Foraminiferal Research, Special Publication 22: 1–46.
- CUSHMAN, J. A. and STAINFORTH, R. M. 1945. The foraminifera of the Ciperio Marl Formation of Trinidad, British West Indies. Contributions from the Cushman Laboratory for Foraminiferal Research 14: 3–75.
- CUSHMAN, J. A. and SIMONSON, R. R. 1944. Foraminifera from the Tumey Formation, Fresno County, California. Journal of Paleontology 18: 186–203.
- CUSHMAN, J. A. and SIEGFUS, S. S. 1942. Foraminifera from the type area of the Kreyenhagen Shale of California. Transactions of the San Diego Society of Natural History 9(34): 385–426.
- CUSHMAN, J. A., STEWART, R. E. and STEWART, K. C. 1947. Eocene Foraminifera from Helmick Hill, Polk County, Oregon. Department of Geology and Mineral Industries, Bulletin 36, Part 5: 93–110.
- CUSHMAN, J. A. and STAINFORTH, R. M. 1951. Tertiary Foraminifera of coastal Ecuador; part I, Eocene. Journal of Paleontology 25(2): 129–164.
- CUSHMAN, J. A. and STONE, B. 1947. An Eocene foraminiferal fauna from the Chira shale of Perú. Cushman Laboratory for Foraminiferal Research, Special Publication 20: 1–27.
- CUSHMAN, J. A. and STONE, B 1949. Foraminifera from the Eocene Verdun Formation of Perú. Contributions from the Cushman Laboratory for Foraminiferal Research 25: 73–84
- CUSHMAN, J. A. and TODD, R. 1944. The Genus *Spiroloculina* and Its Species. Cushman Laboratory for Foraminiferal Research, Special Publication 11: 1–82.
- CUVILLIER, J. and SZAKÁLL, V. 1949. Foraminifères d'Aquitaine. Société Nationale des Pétroles d'Aquitaine: 1–112.
- CSÁSZÁR, G.. (ed) 1997. Basic lithostratigraphic units of Hungary. Charts and short descriptions. Published the Geological Institute of Hungary: 1–114.
- CSONTOS L., NAGYMAROSI A., HORVÁTH F., and KOVÁČ, M 1992. Tertiary evolution of the Intra-Carpathian area: a model. Tectonophysics 208: 221–241.
- CZIZEK, J. 1848. Beitrag zur Kenntniss der fossilen Foraminiferen des Wiener Beckens. Naturwissenschaftliche Abhandlungen 1:137–150.

- DARAKCHIEVA, S. and JURANOV, S. 1992. Small Foraminifers from the Middle and Upper Eocene in the Burgas District. *Calcareous benthic foraminifers. Part I. Geologica Balcanica* 22(4): 3–42.
- DEN DULK, M. 2000. Benthic foraminiferal response to late Quaternary variations in surface water productivity and oxygenation in the northern Arabian Sea. *Geologica Ultraiectiana* 188: 1–205.
- DIECI, G. 1959. I foraminiferi Tortoniani di Montegibbio e Castelvetro (Appenino Modenese). *Palaeontographia Italica* 54: 1–113.
- D'ORBIGNY, A. 1826. Tableau méthodique de la classe des Céphalopodes. *Annales des Sciences Naturelles* 1(7): 245–314
- D'ORBIGNY, A. 1839. Foraminifères. (Voyage dans l'Amérique méridionale) *Annales des Sciences Naturelles* 5: 1–86.
- D'ORBIGNY, A. 1846. Foraminiferes fossiles du Bassin Tertiaire de Vienne. Paris, Gide: 1–312.
- D'ORBIGNY, A. 1850. Prodrome de palaeontologie stratigraphique universelle des animaux mollusques & rayonnés. Faisant suite au cours elementaire de palaeontologie et de geologie stratigraphiques Paris: V. Masson 2: 1–409.
- D'ORBIGNY, A. 1852. Prodrome de paléontologie stratigraphique universelle des animaux mollusques et rayonnés fossils. Paris: V. Masson 3: 1–161.
- DOUGLAS, R.G. and WOODRUFF, F., 1981. Deep-sea benthic foraminifera. The oceanic Lithosphere. In: EMILIANI C. (Ed.), *The Sea*, Wiley-Interscience, New York: 1233–1327.
- EGGER, J.G. 1857. Die Foraminiferen der Miocän-Schichten bei Ortenburg in Nieder-Bayern. *Neues Jahrbuch für Mineralogie, Geognosie, Geologie und Petrefakten-Kunde* 1857: 266–311.
- EGGER, J. G. 1893. Foraminiferen aus Meeresgrundproben, gelohet von 1874 bis 1876 von S.M. Sch. *Gazelle. Königliche Bayerische Akademie der Wissenschaften, Mathematisch-Physikalische Klasse, Abhandlungen* 18(2): 139–458.
- EILAND, M. and GUDMUNDSSON, G. 2004. Taxonomy of some recent Nodosariinae (Foraminifera) from the North Atlantic, with notes on wall lamination. *Micropaleontology* 50: 195–210.
- ELLIS, B. F. and MESSINA, A. R. 1942–2007. *Catalogue of Foraminifera*. The American Museum of Natural History.
- ELLISOR, A.C. 1933. Jackson Group of formations in Texas with notes on Frio and Vicksburg. *American Association of Petroleum Geologists Bulletin* 17: 1293–1350.
- FLINT, J. M. 1899. Recent foraminifera: a descriptive catalogue of specimens dredged by the U.S. Fish Commission Steamer Albatross. Report of the United States National Museum 1897: 249–349.
- FODOR, L., MAGYARI, Á., FOGARASI, A., and PALOTÁS, K. 1994. Tercier szerkezetfejlődés és késő paleogén üledékképződés a Budai-hegységben A Budai-vonal új értelmezése [Tertiary tectonics and Late Paleogene sedimentation in the Buda Hills, Hungary. A new interpretation of the Buda Line]. *Földtani Közlöny [Bulletin of the Hungarian Geological Society]* 124(2): 129–305. [in Hungarian with English summary]
- FORNASINI, C. 1905. Illustrazione di specie orbignyane di Miliolidi istituite nel 1826. *Memorie della Reale Accademia della Scienze di Bologna* 6: 59–70.
- FRANZENAU, Á. 1894. Adatok Letkés faunájához. [Data to the fauna of Letkés] *Mathematikai and Természettudományi Közlemények* 26(1): 1–36. [in Hungarian]
- FRIESE, H. 1951. Zur Foraminiferen-Fauna der Meeresmolasse des unteren Inngebietes. *Abhandlungen des Geologischen Dienstes* 227: 1–52.
- GALLOWAY, J. J. 1933. *A Manual of Foraminifera*. Bloomington, The **Principia** Press: 1–483.
- GALLOWAY, J. J. and MORREY, M. 1929. A lower Tertiary foraminiferal fauna from Manta, Ecuador. *Bulletins of American Paleontology* 15: 7–56.
- GELLAI-NAGY, Á. 1988. Delineation of Hantkens's foraminiferal species from the original collection. *Magyar Állami Földtani Intézet Évi Jelentése [Annual Report of the Hungarian Geological Institute]* 1988(2): 133–173.
- GOODAY, A. J. 1988. A response by benthic Foraminifera to the deposition of phytodetritus in the deep sea. *Nature* 332: 70–73.
- GOODAY, A. J. 1993. Deep-sea benthic foraminiferal species which exploit phytodetritus: characteristic features and controls on distribution. *Marine Micropaleontology* 22: 187–205.
- GÖES, A. 1894. A synopsis of the Arctic and Scandinavian Recent marine Foraminifera hitherto discovered. *Konglischen Svenska Vetenskaps-Akademiens Handlingar* 25(9): 1–127.
- GRAHAM, J. J., DE KLASZ, I. and RÉRAT, D. 1965. Quelques importants Foraminiferes du Tertiaire du Gabon (Afrique équatoriale). *Revue Micropaléontologie* 8(2): 71–84.
- GRZYBOWSKI, J. 1896. Otwornice czerwonych ilów z Wadowic. *Rozprawy Akademia Umiejetnosci w Krakowie, Wydział Matematyczno-Przyrodniczego*, serya 2, 30: 261–308.
- GRZYBOWSKI, J. 1898. Otwornice pokładów naftonosnych okolicy Krosna. *Rozprawy Akademia Umiejetnosci w Krakowie, Wydział Matematyczno-Przyrodniczego*, serya 2, 33: 257–305.

- GULLENTOPS, F. 1968. Les Foraminifères des sables de Vieux Joncs (Tongrien supérieur). Mémoires de l'Institut géologique de l'Université de Louvain 20: 1–24.
- GÜMBEL, C. 1868. Beiträge zur Foraminiferenfuna der nordalpinen, älteren Eocängebilde oder Kressenberger Nummulitenschichten. Abhandlungen der Bayerische Akademie der Wissenschaften 10: 580–730.
- GRÜNIG, A., 1985. Systematical description of Eocene benthic foraminifera of Possagno (northern Italy), Sansoain (northern Spain) and Biarritz (Aquitaine, France). *Memorie di Scienze Geologiche*, Padova 37: 251–302.
- HAGEMAN, J. 1979. Benthic foraminiferal assemblages from Plio-Pleistocene open bay to lagoonal sediments of the Western Peloponnesus (Greece). *Utrecht Micropaleontological Bulletins* 20: 1–171.
- HAGN, H. 1956. Geologische und Palaeontologische Untersuchungen im Tertiär des Monte Brione und seiner Umgebung. *Paleontographica* 107: 1–67.
- HALKYARD, E. 1919. The fossil foraminifera of the Blue Marl of Côte des Basques Biarritz. *Memoirs and Proceeding of the Manchester Literary and Philosophical Society* 62(2): 1–145.
- HANTKEN M. 1868. A kis-czelli tállyag foraminiferái. [Foraminifers of the Kiscell Clay] *Magyarhoni Földtani Társulat Munkálatai* 4: 75–96.
- HANTKEN M. 1875. A *Clavulina Szabói* rétegek faunája I. Foraminiferák. [Fossils of the *Clavulina Szabói* layers. Part I. Foraminifers] *A Magyar Királyi Földtani Intézet Évkönyve* 4: 1–82.
- HAQUE, M. 1956. The smaller Foraminifera of the Ranikot and the Laki of the Nammal George, Salt Range. *Palaeontologia Pakistanica* 1: 1–300.
- HARDENBOL, J., THIERRY, J., FARLEY, M.B., JACQUIN, TH., DE GRACIANSKY, P.C. and VAIL, P.R. 1998. Mesozoic and Cenozoic Sequence Chronostratigraphic Framework of European basins. *SEPM Special Publication* 60: 1–364.
- HARRIS, R. W., and JOBE, B. I. 1951. Microfauna of basal Midway outcrops near Hope, Arkansas: Norman, Okla., Transcript Press: 1–113.
- HAYWARD, B. W., GRENFELL, H. R., REID, C. M. and HAYWARD, K. A. 1999. Recent New Zealand shallow-water benthic foraminifera: taxonomy, ecologic distribution, biogeography, and use in paleoenvironmental assessment. *Institute of Geological and Nuclear Sciences Monograph* 21:1–258.
- HERMANN, A., 1917. Vierter Beitrag zur Kenntnis der Vorkommens von Foraminiferen im Tertiär der Unterelsass. *Alsace-Lorraine, Geologischen Landesanstalt Mitteilungen* 10: 1–290.
- HILLEBRANDT, A. 1962. Das Paläozen und seine Foraminiferenfauna im Becken von Reichenhall und Salzburg. *Abhandlungen der Bayerische Akademie der Wissenschaften* 108: 1–182.
- HOFMANN G. W., 1967: Untersuchungen an der Gattung *Bolivina* (Foraminifera) im Oligozän und Miozän der ostbayerischen Molasse. *Geologica Bavarica* 57: 121–204.
- HOFKER, J. 1930. Foraminifera of the Siboga Expedition, Part 2: Families Astorhizidae, Rhizamminidae, Reophacidae, Anomalinidae, Peneroplidae. *Siboga-Expeditie, Monographie* 4: 79–170.
- HOFKER, J. 1956. Foraminifera Dentata, foraminifera of Santa Cruz and Thatch Island, Virgin Archipelago, West Indies. *Spolia Zoologica musei Hauniensis* 15: 1–237.
- HOFKER, J. 1966. Maestrichtian, Danian and Paleocene Foraminifera. *Paleontographica* 10: 1–376.
- HOFKER, J. 1968. Tertiary Foraminifera of Coastal Ecuador. Lower Oligocene and Lower Miocene. *Palaeontographica* 13: 1–59.
- HORNIBROOK, N. de B., 1961. Tertiary foraminifera from Oamaru District (N.Z.). *New Zealand Geological Survey, Paleontological Bulletin* 34: 1–194.
- HORVÁTH, M. 2002: Data to revision and distribution of small foraminifera species described by Hantken (1868, 1875). Part I. Textulariidae and Miliolidae. *Fragmenta Palaeontologica Hungarica* 20: 25–42.
- HORVÁTH, M. 2003. Data to revision and distribution of small Foraminifera species described by Hantken (1868, 1875). Part 2. Nodosaridae and Vaginulidae. *Fragmenta Palaeontologica Hungarica* 21: 5–32.
- HORVÁTH-KOLLÁNYI K. 1983. Az ÉK dunántúli terület eocén plankton Foraminifera zónái. [Eocene planktonic foraminifera zones from the NE part of Transdanubian Central Range] *Földtani Közlöny* [Bulletin of the Hungarian Geological Society] 113: 225–236. [in Hungarian]
- HORVÁTH-KOLLÁNYI K. 1988. Eocene benthonic smaller foraminifera fauna from Dudar. *A Magyar Állami Földtani Intézet Évkönyve* [Annals of the Hungarian Geological Institute] 63(4): 1–173.
- HOTTINGER, L., HALICZ, E. and REISS, Z. 1993. Recent foraminiferida from the Gulf of Aqaba, Red Sea. *Slovenska Akademija Znanosti in Umenosti (Ljubljana), classis IV*: 33: 1–179.
- HOWE, H. V. 1939. Louisiana Cook Mountain Eocene Foraminifera. *Bulletin of Louisiana Geological Survey* 14: 1–122.
- HOWE, H.V.W. and WALLACE, W. E. 1932. Foraminifera of the Jackson Eocene at Danville Landing on the Ouachita, Catahoula Parish, Louisiana. *Geological Bulletin, Louisiana Department of Conservation, Geological Survey* 2: 18–79.
- HUSSEIN, A. I. 1997. Some agglutinated foraminifera from the Middle Eocene rocks of Egypt. *Middle East Research Center, Ain Shams University, Earth Science Series, Cairo* 11: 106–114.

- JONKERS, H. A. 1984. Pliocene benthonic foraminifera from homogenous and laminated marls on Crete. Utrecht Micropaleontological Bulletin 31: 1–180.
- KAASSCHIETER, J. P. H. 1961. Foraminifera of the Eocene of Belgium. Institut Royal des Sciences Naturelles de Belgique, Mémoires 147: 1–271.
- KARRER, F. 1868. Die Miocene Foraminiferenfauna von Kostež im Banat. Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften in Wien 58: 121–193.
- KARRER, F. 1877. Geologie der Kaiser Franz Josefs Hochquellen-Wasserleitung. Eine Studie in den Tertiaer-Bildungen am Westrande des alpinen Theiles der Niederung von Wien. Abhandlungen der Kaiserlich-Königlichen Geologischen Reichsanstalt 9: 1–420.
- KÁZMÉR, M., DUNKL, I., FRISCH, W., KUHLEMANN, J. and OZSVÁRT, P., 2003. The Palaeogene forearc basin of the Eastern Alps and Western Carpathians: subduction erosion and basin evolution. Journal of the Geological Society 160: 413–428.
- KECSKEMÉTI, T. and VÖRÖS, A. 1975. Biostratigraphic and palaeoecologic studies on the Eocene transgressive series at Darvástó (Bakony Mts., Hungary). Fragmenta Mineralogica et Palaeontologica 6: 63–93.
- KHALILOV, D. M. 1951. New representatives of the foraminifera from the Paleogene deposits of Azerbaidzhan, Akademiya Nauk Azerbaidzhanskoi SSR, Izvestiya, Baku 2: 1–10.
- KIESEL, Y. 1962. Die Oligozänen Foraminiferen der Tiefbohrung Dobbertin (Mecklenburg). Freiburger Forschungshefte 122: 8–98.
- KITAZATO, H., SHIRAYAMA, Y., NAKATSUKA, T., FUJIWARA, S., SHIMANAGA, M., KATO, Y., OKADA, Y., KANDA, J., YAMAOKA, A., MASUZAWA, T. and SUZUKI, K. 2000. Seasonal phytodetritus deposition and responses of bathyal benthic foraminiferal populations in Sagami Bay, Japan: preliminary results from “Project Sagami 1996–1999”, Marine Micropalaeontology 40: 135–149.
- KOHL, B. 1985. Early Pliocene benthic foraminifers from the Salina Basin, southeastern Mexico. Bulletin of American Paleontology 88(322): 1–173.
- KOPEK, G. 1980. Eocene of north-eastern part of Bakony Mountains. Annals of the Hungarian Geological Institute 63: 1–176.
- KORECZNÉ LAKY I. and NAGYNÉ GELLAI Á. 1985: Foraminiferal fauna from the Oligocene and Miocene in the Börzsöny Mountains. A Magyar Állami Földtani Intézet Évkönyve [Annals of the Hungarian Geological Institute] 68: 1–527.
- KOUWENHOVEN, T. J. 2000. Survival under stress: benthic foraminiferal patterns and Cenozoic biotic crises. Geologica Ultraiectiana 186: 1–206.
- KOVÁCS, S., SZEDERKÉNYI, T., HAAS, J., BUDA, Gy., CSÁSZÁR, G., NAGYMAROSY, A. 2000. Tectonostratigraphic terranes in the pre-Neogene basement of the Hungarian part of the Pannonian area. Acta Geologica Hungarica 43(3): 225–328.
- KRAYEVA, E. 1965. Об объеме вида *Marginulina behmi* (Reuss) [On the scope of the species *Marginulina behmi* (Reuss)] Палеонтологический сборник [Paleontologicheskyy Sbornik] 2: 3–7. [in Russian with English summary]
- KRAYEVA, E. AND ZARNECKII, B. F. 1969. Палеонтологический йаравочник. Фораминиферы палеогена Украины. [Paleontological problems. Paleogene Foraminifers from Ukraine]. Akademia Nauka Ukraina 3: 1–199. [in Russian].
- KRHOVSKY, J., 1983. Foraminiferida of the Eocene/Oligocene boundary from the Pouzdrany Formation (the West Carpathians, CSSR). Knihovnička Zemního plynu a nafty 4: 71–98.
- KUHN, J.A., 1972. Stratigraphisch-mikropalaeontologische Untersuchungen in der Aeusseren Einsiedler Schuppenzone und im Waegitaler Flysch E und W des Sihlsees (Kt. Schwyz). Eclogae Geologicae Helveticae 65(3): 485–553.
- KÜMMERLE, E. 1963. Die Foraminiferenfauna des Kasseler Meeressandes (Oberoligozän) im Ahnetal bei Kassel. Abhandlungen des Hessischen Landesamtes für Bodenforschung 45: 1–72.
- LALICKER, C. G. 1935. New Tertiary Textulariidae. Contributions from the Cushman Laboratory for Foraminiferal Research 11(2): 39–51.
- LE CALVEZ, Y. 1947. Révision des Foraminifères lutétien du Bassin de Paris I. Miliolidae. Mémoire de la Carte Géologique France: 1–41.
- LE CALVEZ, Y. 1949. Révision des Foraminifères lutétien du Bassin de Paris II. Rotaliidae et familles affines. Mémoire de la Carte Géologique France: 1–54.
- LE CALVEZ, Y. 1950. Révision des Foraminifères lutétien du Bassin de Paris III. Polymorphinidae, Buliminidae, Nonionidae. Mémoire de la Carte Géologique France: 1–64.
- LE CALVEZ, Y. 1952. Révision des Foraminifères lutétien du Bassin de Paris IV. Valvulinidae, Peneroplidae, Ophthalmididae, Lagenidae. Mémoire de la Carte Géologique France: 1–64.
- LE CALVEZ, Y. 1970. Contribution à l'étude des Foraminifères Paléogènes du Bassin du Paris, Cahiers de Paléontologie. Éditions du CNRS: 1–326.



- LIEBUS, A. 1911. Die Foraminiferenfauna der mitteleocänen Mergel von Norddalmatien, Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften 120(1): 865–956.
- LINNÉ, C. 1758. Systema naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis. G. Engelmann (Lipsiae) 10: 1–824.
- LOEBLICH, A.R., and TAPPAN, H. 1988. Foraminiferal genera and their classification. Van Nostrand Reinhold Company, New York, 2 vols.: 1–970.
- LUCZKOWSKA, E. 1972. Miliolidae (Foraminiferida) from Miocene of Poland. Part I: Revision of the classification. Acta Palaeontologica Polonica 17: 341–377.
- MACKENSEN, A. and BERGGREN, W. A., 1992. Paleogene Benthic Foraminifers from the Southern Indian Ocean (Kerguelen Plateau): Biostratigraphy and Paleoecology. In: WISE, S. W., Jr., SCHLICH, R., et al., (Eds.) Proceedings of the Ocean Drilling Program, Scientific Results 120: 603–630.
- MAJZON, L. 1962. Hantken Miksa “*Clavulinoides Szabói* rétegek faunája” című művének nevezéktani módosítása. [Nomenclatural revision of “Fossils of the *Clavulinoides Szabói* layers” by Maximilian Hantken] Földtani Közlöny [Bulletin of the Hungarian Geological Society] 92: 268–273.
- MAJZON, L. 1966. Foraminifera vizsgálatok. [Foraminiferal examination]. Akadémiai Kiadó, Budapest: 1–939. [in Hungarian]
- MAJZON, L. 1972. Anmerkungen zur ungarischen *Clavulinoides* Arten. Földtani Közlöny 102: 109–121.
- MARKS, P. 1951. A revision of the smaller foraminifera from the Miocene of the Vienna Basin. Cushman Foundation for Foraminiferal Research 2: 33–73.
- MATHELIN, J. C. and SZTRÁKOS, K. 1993. L'Éocène de Biarritz (Pyrénées-Atlantiques, SW France). Stratigraphie et paléoenvironnement. Monographie des foraminifères. Cahiers de Micropaléontologie 8: 5–85.
- MCDUGALL, K. A., 1972. The Narizian-Refugian boundary in the Twin River Formation of the Northern Olympic Peninsula, Washington. MSc Thesis, University of Washington, Seattle, Washington: 1–223.
- MILLER, K. G., 1983. Eocene-Oligocene paleoceanography of the deep Bay of Biscay: benthic foraminiferal evidence. Marine Micropaleontology 7: 403–440.
- VAN MORKHOVEN, F., BERGGREN, W. A. and EDWARDS, A. S. 1986. Cenozoic cosmopolitan deep-water benthic foraminifera. Bulletin des Centres de Recherches Exploration-production Elf-Aquitaine, Memoires 11: 1–421.
- MURRAY, J. W. 1991. Ecology and Palaeoecology of Benthic Foraminifera. Longman, London: 1–397.
- MURRAY, J. W. and WRIGHT, C. A., 1974. Palaeogene foraminiferida and palaeoecology, Hampshire and Paris basins and the English Channel. Special Papers in Palaeontology 14: 1–171.
- MÜHLSTRASSER, T. 2001. Paleoceanographic history of the Northwestern Tethyan Realm for the Late Oligocene through Middle Miocene. Tübinger Mikropaläontologische Mitteilungen 24: 1–119.
- MYATLYUK, E. V. 1953. Iskopaemye foraminifery SSSR. Spirillinidy, rotaliidy, epistominidy i asterigerinidy. VNIGRI, Trudy 71: 1–274.
- MYATLYUK, E. V. 1970. Foraminiferi flishevikh otlozenii Vostochnikh Karpat. Vses. Nauchnoissled Geol. Institute, Trudy 282: 4–360.
- NAGYMAROSY, A. 1983. Calcareous nannofloras at the Eocene/Oligocene boundary and in the Early Oligocene sediments in Hungary. Discussiones Palaeontologicae 31: 25–32.
- NYÍRŐ, R. 1970. Study of the smaller Foraminifera of the Weim-pusztá Middle Eocene key section at Nagyesztergár (Bakony Mountains, Transdanubia, Hungary). Annales Historico-naturales Musei Nationalis Hungarici 62: 67–84.
- NAGY-GELLAI, Á. 1973. Oligocén Foraminiferák Dorog környékéről. [Oligocene foraminifera from Dorog] A Magyar Állami Földtani Intézet Évkönyve [Annals of the Hungarian Geological Institute] 55(3): 419–495. [in Hungarian]
- NARAYAN, Y. R., BARNES, C. R. AND JOHNS, M. J. 2005. Taxonomy and biostratigraphy of Cenozoic foraminifers from Shell Canada wells, Tofino Basin, offshore Vancouver Island, British Columbia. Micropaleontology 51(2): 101–168.
- NEUGEBORN, J. L. 1856. Die Foraminiferen aus der Ordnung der Stichostegier von Ober-Lapugy in Siebenbürgen. Denkschriften der Kaiserlichen Akademie der Wissenschaften, Wien, Mathematisch-Naturwissenschaftliche Klasse 12: 65–108.
- NEUMAYR, M. 1889. Die Stämme des Tierreiches, Band 1, Wirbellose Thiere: Wien, Prag, Verlag von F. Tempsky & Tempsky: 1–603.
- NUTTALL, W. L. F. 1932. Lower Oligocene foraminifera from Mexico. Journal of Paleontology 6: 3–35.
- NYÍRŐ, R. 1958. The studying of Foraminifera in thin section. Annales Universitatis Scientiarum Budapestinensis, sectio Geologica, 9: 31–38.
- KATZ, M. E. and MILLER, K. G. 1993. Miocene-Pliocene bathyal benthic foraminifera and the uplift of Buff Bay, Jamaica. In: WRIGHT, R. M. and ROBINSON, E., (Eds.) Biostratigraphy of Jamaica. Geological Society of America Memoir 182: 219–254.


- ORTIZ, S. and THOMAS, E. 2006. Lower-middle Eocene benthic foraminifera from the Fortuna Section (Betic Cordillera, southeastern Spain). *Micropaleontology* 52(2): 97–150.
- OZSVÁRT, P. 1999. Middle Eocene foraminifer, mollusc and ostracod fauna from the Csordakút Basin (Gerecse Mountains, Hungary): palaeoenvironments recorded in a transgressive sequence. *Annales Universitatis Scientiarum Budapestinensis, sectio Geologica* 32: 73–135.
- OZSVÁRT, P. 2003a. Five new species of benthic Foraminifera from Hungarian Paleogene Basin. *Fragmenta Paleontologica Hungarica* 21: 1–5.
- OZSVÁRT, P. 2003b. Palaeoceanographic history of the Hungarian Paleogene Basin during the Eocene. PhD dissertation, Eötvös University, Budapest. 182 pp.
- PAPP, A. 1975. Die Entwicklung der Uvigerinen in der Puchkirchener Schichtengruppe. In: *Chronostratigraphie und Neostatotypen V, Miozän OM Egerien*: 279–287.
- PAPP, A. and SCHMID, M. E. 1985. Die fossilen Foraminiferen des Tertiären Beckens von Wien—Abhandlungen der Geologischen Bundesanstalt 37: 1–107.
- PAPP, A. and TURNOVSKY, K. 1953. Die Entwicklung der Uvigerinen im Vindobon (Helvet und Torton) des Wiener Beckens. *Jahrbuch der Geologischen Bundesanstalt* 94(1): 117–142.
- PERELIS, L. and REISS, Z. 1975. Cibicididae in recent Sediments from the Gulf of Elat. *Israel Journal of Earth Sciences* 24: 73–96.
- PETERS, S. W. 1982. Central West African Cretaceous-Tertiary benthic foraminifera and stratigraphy. *Paleontographica, Abteilung A* 179: 1–104.
- PFLUM, C. E., and FRERICHS, W. E. 1976. Gulf of Mexico deep-water foraminifers. *Cushman Foundation for Foraminiferal Research, Special Publications* 14: 1–125.
- PLUMMER, H. J. 1926. Foraminifera of the Midway Formation in Texas. *University of Texas Bulletin* 2644: 9–206.
- POPESCU, G. 1975. Étude des Foraminifères du Miocène inférieur et moyen du nord-ouest de la Transylvanie. *Mémoire Institut de Géologie et Geophysique (Bucureşti)* 23: 1–121.
- POPESCU, G. and IVA, M. 1971. Contribution à la connaissance de la microfaune Oligocene des couches de Valea Lapusului. *Memoires etudes Micropaleontologie* 14: 35–52.
- POŻARYSKA, K. and SZCZETCHURA, J., 1968. Foraminifera from the Paleocene of Poland, their ecological and biostratigraphical meaning. *Palaeontologia Polonica* 20: 1–107.
- POŻARYSKA, K. 1977. Upper Eocene Foraminifera of East Poland and their palaeogeographical meaning. *Acta Palaeontologica Polonica* 22: 3–54.
- PROTESCU, O. 1923. La microfauna des marnes a glauconie de la region Tintea (Distr. Prahova) et l'importance stratigraphique de l'espèce *Clavulina szabói* Hantken. *Publ. Soc. Naturalist România* 11: 42–67.
- PROTO DECIMA, F. and DE BIASE R. 1975. Foraminiferi bentonici del Paleocene, dell'Eocene inferiore e medio. In BRAGA et al. 1975. *Foraminiferi bentonici del Paleocene ed Eocene della sezione di Possagno*. *Schweizerische Paleontologische Abhandlungen* 97: 87–98.
- REISER, H. 1987. Die Foraminiferen der bayerischen Oligozän-Molasse: Systematik, Stratigraphie und Paläobathymetrie. *Zitteliana* 16: 1–131.
- REUSS, A. E. 1850. Neue Foraminiferen aus den Schichten des österreichischen Tertiärbeckens. *Denkschriften der Kaiserlichen Akademie der Wissenschaften, Wien, Mathematisch–Naturwissenschaftliche Klasse* 1: 365–390.
- REUSS, A. E., 1851. Über die fossilen Foraminiferen und Entomostraceen der Septarienthone der Umgegend von Berlin, *Zeitschrift der Deutschen Geologischen Gesellschaft* 3: 49–92.
- REUSS, A. E. 1863. Beiträge zur Kenntniss der tertiären Foraminiferen-Fauna. *Denkschriften der Kaiserlichen Akademie der Wissenschaften, Wien, Mathematisch–Naturwissenschaftliche Klasse* 48: 36–71.
- REUSS, A. E. 1866. Die Foraminiferen, Anthozoen und Bryozoen des deutschen Septarienthones. *Denkschriften der Kaiserlichen Akademie der Wissenschaften, Wien, Mathematisch–Naturwissenschaftliche Klasse* 25: 117–214.
- REUSS A. E. 1870 Die Foraminiferen des Septarien-Thones von Pietzpuhl. *Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften in Wien* 62: 455–493.
- ROBERTSON, B. E. 1998. Systematics and paleoecology of the benthic Foraminiferida from the Buff Bay section, Miocene of Jamaica. *Micropaleontology* 44 (suppl. 2): 1–266.
- ROEMER, P. A. 1838. Die Cephalopoden des Nord-Deutschen tertiären Weersandes. *Neues Jahrbuch für Mineralogie, Geognosie, Geologie und Petrefakten-Kunde*: 1–386.
- ROMEO, M. 1967. Stratigrafia micropaleontologica del Messiniano di Rossano (Cosenza). *Palaeontographica Italica* 33: 1–74.
- RÖGL, F. 1969. Die miozäne Foraminiferenfauna von Laa an der Thaya in der Molassezone von Niederösterreich. *Mitteilungen der Geologischen Gesellschaft in Wien* 61: 63–123,
- RUIZ DE GAONA, M. and COLOM, G. 1950. Estudios sobre las sinecias de los foraminíferos eocénicos de la vertiente meridional del Pirineo (Cataluña-Vizcaya). *Estudios Geológicos* 12: 293–434.

- SARS, M. 1868. Forstatte Bemaerkninger over det dyriske livs Undbredningi Havest Dyber. Christiania Videnskaps-Selskabs Forhandlingar: 244–275
- SAMOILOVA, R. 1947. О некоторых новых и характерных видах фораминифер из верхнего палеогена Крыма. [On some new and characteristic species of foraminifera from Upper Paleogene in the Crimea]. Бюллетен Москва [Bulletin Moscou] 22(4): 78–101. [in Russian]
- SAMUEL, O. 1975. Foraminifera of Upper Priabonian from Lúbietová (Slovakia). Západné Karpaty sér. Paleontológia 1: 111–176.
- SAPERSON, E. and JANAL, M. 1980. Biostratigraphy of the Anomaliniidae and Cibicididae in the Soviet Tethyan Paleogene. Micropaleontology 36: 392–413.
- SCHOTT, W. 1935. Die Foraminiferen in dem Äquatorialen Teil des Atlantischem Ozeans. Deutsche Atlantische Expedition 6: 411–416.
- SCHUBERT, R. J. 1901. Neue und interessante Foraminiferen aus dem südtiroler Alteriär. Beiträge zur Paläontologie und Geologie Österreich-Ungarns und des Orients 14: 9–26.
- SCHWAGER, C. 1866. Fossile Foraminiferen von Kar Nikobar, Reise der Oesterreichischen Fregatte Novara um Erde in den Jahren 1857, 1858, 1859 unten den Befehlen des Commodore B. Von Wuellerstorff-Urbair. Geologischer Theil, Geologische Beobachtung no. 2, Palaeontologische Mittheilung 2(1): 187–268.
- SCHWEIZER, M. 2006. Evolution and molecular phylogeny of *Cibicides* and *Uvigerina* (Rotaliida, Foraminifera). Geologica Ultraiectina 261: 1–167.
- SEGUENZA, G. 1862 Dei terreni Terziarii del distretto di Messina; Parte II. Descrizione del foraminiferi monotalamici delle marne mioceniche del distretto di Mes.sina. Printed by Capra: 1–66.
- SEN GUPTA, B. K. 1999. Foraminifera in marginal marine environments. In: SEN GUPTA, B. K. (Ed.): Modern Foraminifera. Dordrecht (Kluwer): 141–159.
- SETIAWAN, J. R. 1983. Foraminifera and microfacies of the type Priabonian. Utrecht Micropaleontological Bulletin 29: 1–161.
- ŞIREL, E. 1976 Description of the new species *Eoannularia conica* n.sp. and new observations on the Upper Lutetian–Lower Priabonian limestone in the Darende–Gürün (west of Malatya) region. Bulletin of the Geological Society of Turkey 19: 79–82.
- SMITH, B.Y. 1957. Lower Tertiary Foraminifera from Contra Costa County California. California University, Publications of the Geological Society 32: 1–190.
- STACHE, G. 1864. Die Foraminiferen der tertiären Mergel des Whaingaroa-Hafens (Prov. Auckland). Novara-Expedition (1857–1859) Geologischer Theil 1(2): 159–304.
- SOUAYA, F. J. 1965. Miocene foraminifera of the Gulf of Suez region, U.A.R. Part 1 – Systematics (Astrorhizoidea – Buliminioidea). Micropaleontology 11: 301–344.
- SUBBOTINA, N. N. 1947 Foraminifers of the Danian and Paleogene deposits of the northern Caucasus.– In: Microfaun of the Caucasus Emba region and central Asia. Vsesoyuznyi nauchno-issledovatel'skii geologorazvedochni neftyanoi institut (VNI-GRI), new series 51: 39–160.
- SUBBOTINA, N. N. 1953. Upper Eocene Lagenidae and Buliminidae of the southern U.S.S.R. Vsesoyuznyi nauchno-issledovatel'skii geologorazvedochni neftyanoi institut (VNI-GRI), new series 69(6): 115–282.
- SZCZUCHURA, J. and POŻARYSKA, K. 1974. Foraminiferida from the Paleocene of Polish Carpathians (Babica clays). Paleontologia Polonica 31: 1–142.
- SZÓTS, E. 1956. Magyarország eocén (paleogén) képződményei. [Paleogene sequences of Hungary] Geologica Hungarica, Series Geologica 9: 1–320.
- SZTANÓ, O. and FODOR, L. 1997. Lejtőüledékek a paleogén medence peremén: A felső-eocén Piszkei Márga (Nyergesújfalu, Sánc-hegy) üledései és szerkezeti viszonyai. [Bathyal slope deposits in the Paleogene Basin: A case study of the Upper Eocene Piszke Marl (Nyergesújfalu, Sánc Hill, Hungary)] Földtani Közlöny [Bulletin of the Hungarian Geological Society] 127(3–4): 267–290.
- SZTRÁKOS, K. 1974. Paleogene planktonic foraminiferal zones in Northeastern Hungary. Fragmenta Mineralogica et Paleontologica 5: 29–81.
- SZTRÁKOS K. 1979. La stratigraphie, peléoécologie, paléogéographie et les foraminifères de l'oligocène du nord-est de la Hongrie. Cahiers de Micropaléontologie 1979(3): 1–95.
- SZTRÁKOS, K. 1982. Les Foraminifères de la Marne de Buda et la limite Eocene–Oligocene en Hongrie. Cahiers de Micropaléontologie 1982(4): 3–48.
- SZTRÁKOS, K. 1983. Le genre *Uvigerina* (foraminifères) dans le Paleogene de la Hongrie. Revue de Micropaleontologie 26: 132–142.
- SZTRÁKOS, K. 1987. Les Foraminifères bartoniens et priaboniens de couches à “Tritaxia szaboi” de Hongrie et essai de reconstruction paléogéographique de la Montagne Centrale de Hongrie au Bartonien et au Priabonien. Cahiers de Micropaléontologie 1987(2): 5–37.
- TARI G., BÁLDI T., and BÁLDI-BEKE M. 1993. Paleogene retroarc flexural basin beneath the Neogene Pannonian Basin: a geodynamic model. Tectonophysics 226: 433–455.

- TERQUEM, M. 1882. Les Foraminifères de l'Éocène des environs de Paris. Mémoires de la Société Géologique de France, Paléontologie 3(2): 1–193.
- THALMANN, H. E. 1933. Nachtrag zum Nomenclator zur Brady's Tafelband der Foraminiferen der Challenger-Expedition. *Eclogae Geologicae Helveticae* 26: 251–255.
- TJALSMA, R. C. and LOHMANN, G. P. 1983. Paleocene-Eocene bathyal and abyssal benthic foraminifera from the Atlantic Ocean. *Micropaleontology, Special Publication* 4: 1–90.
- TODD, R. and KNIKER, H. T. 1952. An Eocene foraminiferal fauna from the Agua Fresca shale of Magallanes Province, southernmost Chile. *Cushman Laboratory for Foraminiferal Research, Special Publication* 1: 1–28.
- TOULMIN, L. D. 1941. Eocene smaller foraminifera from the Salt Mountain Limestone of Alabama. *Journal of Paleontology* 15(6): 567–611.
- TOUMARKINE, M. 1971. Etude des Foraminifères planctoniques de deux sondages (H-849 et Pgyt-31) dans l'Éocène de la Montagne du Bakony (Transdanubie, Hongrie). *Annales Instituti Geologici Publici Hungarici* 54(4): 283–299.
- VAN DER ZWAAN, G. J. 1982. Paleocology of Late Miocene Mediterranean foraminifera. *Utrecht Micropaleontological Bulletins* 25: 1–202.
- VERDENIUS, J. G. 1970. Neogene stratigraphy of the Western Guadalquivir Basin (Southern Spain). *Utrecht Micropaleontological Bulletins* 3: 1–109.
- VERHALLEN, P. J. J. M. 1991. Late Pliocene to Early Pleistocene Mediterranean mud-dwelling foraminifera: influence of a changing environment on community structure and evolution. *Utrecht Micropaleontological Bulletins* 40: 1–219.
- VERHOEVE, D. 1971. Benthonic foraminifera of the Neogene Asteri, Dhramia and Francocastello formations from Western Crete, Greece. *Departement Aardwetenschappen, Leuven, Katholieke Universiteit Leuven*: 1–290.
- VILKS, G. 1969. Recent foraminifera in the Canadian Arctic. *Micropaleontology* 15(1): 35–60.
- VITÁLIS-ZILAHY, L. 1967. Felsőeocén Foraminiferák Felsőtárkány környékéről (DNy-Bükk). [Late Eocene foraminifera from felsőtárkány, Bükk Mountains] *A Magyar Állami Földtani Intézet Évi Jelentése [Annual Report of the Hungarian Geological Institute] 1965*: 393–441.
- VOGL, V. 1910. A piszkei bryozoás márga faunája. [Fossils of the Piszke Bryozoan Marl] *A Magyar Állami Földtani Intézet Évkönyve [Annals of the Hungarian Geological Institute] 1910*: 176–204.
- VÖRÖS A. 1989. Middle Eocene transgression and basin evolution in the Transdanubian Central Range, Hungary: sedimentological contributions. *Fragmenta Mineralogica et Paleontologica* 14: 63–72.
- WALKER, G. and BOYS, W. 1784. *Testacea minuta rariora, nupperine detecta in arena littoris Sandvicensis*. Printed by J. March. London
- WALKER, G. and JACOB, E. 1798. An arrangement and description of minute and rare shells. In: KANMACHER, F. *Adams' Essays on the microscope*. Printed by Dillon and Keating: 1–633.
- WENGER, W. F. 1987. Die Foraminiferen des Miozäns der bayerischen Molasse und ihre stratigraphische sowie paläogeographische Auswertung. *Zitteliana* 16: 173–340.
- WIDMARK, J. G. V., 1997. Deep-sea benthic foraminifera from Cretaceous-Paleogene boundary strata in the South Atlantic: taxonomy and paleoecology. *Fossils and Strata*, 43:1–94.
- WILLIAMSON, W. C. 1848. On the recent British species of the genus *Lagena*. *Annals and Magazine of Natural History, series 2*, 1: 1–20.
- WÓJCIK, K., 1903. Dolno oligoceńska fauna Kruhela Małego pod Przymysłem (warstwy z *Clavulina Szaboi*). Pt. I: Otornice i mięczaki. *Polska Wydział Akademiens Umiejetnosci Rozpr.* 43: 489–569.
- ZÁGORŠEK, K. & KÁZMÉR, M. (2001): Eocene Bryozoa from Hungary. – *Courier Forschungsinstitut Senckenberg* 231, 1–159, 7 figs, 2 tab., 36 pls, Frankfurt am Main.
- ZILAHY L. 1969. A Planorbulinidae, Acervulinidae, Cymvaloporidae család a magyarországi eocénben. [The family Planorbulinidae, Acervulinidae and Cymvaloporidae from the Eocene of Hungary] *A Magyar Állami Földtani Intézet Évi Jelentése [Annual Report of the Hungarian Geological Institute] 1966*: 153–178.

Plate 1

1. *Astrorhiza bakonycsernyensis* n.sp. (Inv. num.: M 2008.116.1) scale=1.1 mm
- 2–3. *Bathysiphon eocenicus* CUSHMAN and HANNA, 1927 scale=2 mm
4. *Bathysiphon saidi* (ANAN, 1994) scale=1.4 mm
5. *Rhabdammina abyssorum* SARS, 1869 scale=1.3 mm
6. *Reophax harrisi* nomen novum scale=2 mm
7. *Ammomarginulina* sp. scale=1.9 mm
8. *Haplophragmoides* sp. scale=1.23 mm
9. *Bolivinopsis foliacea* (GRZYBOWSKI, 1898) scale=1.9 mm
- 10–11. *Spiroplectammina carinata* (D'ORBIGNY, 1846) scale=0.43 mm
12. *Spiroplectammina subhaeringensis* (GRZYBOWSKI, 1896) scale=0.38 mm
13. *Vulvulina advena* CUSHMAN, 1926 scale=1.6 mm
14. *Vulvulina jarvisi* CUSHMAN, 1932 scale=1.5 mm
15. *Plectina dalmatina* (SCHUBERT, 1911) scale=0.75 mm
16. *Plectina eocenica* CUSHMAN, 1936 scale=1.63 mm
17. *Verneuilina* sp. scale=0.95 mm
18. *Dorothia textilaroides* (HANTKEN, 1875) scale=0.95 mm
19. *Martinottiella rhumberi* CUSHMAN, 1936 scale=0.55 mm
20. *Textularia crookshanki* (HAQUE, 1956) scale=1.8 mm



## Plate 2

- 1–2. *Textularia deperdita* D'ORBIGNY, 1826 scale=0.25 mm
3. *Textularia halkyardi* LALICKER, 1935 scale=0.95 mm
- 4–5. *Textularia globosa* (HANTKEN, 1875) scale=1 mm
- 6–7. *Textularia* cf. *partschii* CZIZEK, 1848 scale=0.85 mm
8. *Textularia lanceolata* (KARRER, 1861) scale=1.55 mm
9. *Textularia pala* CZIZEK, 1848 scale=0.45 mm
- 10–11. *Textularia* sp. 1 scale=1 mm
12. *Textularia* sp. 2 scale=1 mm
- 13–14. *Textularia* sp. 3 scale=0.25 mm
- 15–16. *Clavulinoides alpina* (CUSHMAN, 1936) scale=1.53 mm
17. *Clavulinoides lakiensis elongata* HAQUE, 1949 scale=1.6 mm
- 18–19. *Clavulinoides szabói* (HANTKEN, 1868) scale=1.55 mm
20. *Clavulinoides procerus* n. sp. (Inv. num.: M 2008.117.1) scale=0.6 mm
21. *Clavulina terterensis* KHALILOV, 1958 scale=1.3 mm
- 22–23. *Cylindroclavulina colomi* HAGN, 1956 scale=1 mm
24. *Cylindroclavulina rudislosta* (HANTKEN, 1875) scale=1.6 mm
- 25–27. *Tritaxilina pupa* (GÜMBEL, 1868) scale=1.85 mm






Plate 3

1. *Tritaxilina* sp. scale=1.85 mm
- 2–3. *Cornuspira involens* (REUSS, 1864) scale=0.65 mm
4. *Adelosina* sp. scale=0.68 mm
5. *Spiroloculina obscura* (CUSHMAN and TODD, 1882) scale=0.34 mm
- 6–7. *Spiroloculina bicarinata* TERQUEM, 1882 scale=0.8 mm
- 8–9. *Spiroloculina jarvisi* CUSHMAN and TODD, 1944 scale=0.75 mm
10. *Quinqueloculina buchiana* D'ORBIGNY, 1846 scale=0.9 mm
11. *Quinqueloculina juleana* D'ORBIGNY, 1846 scale=0.45 mm
12. *Quinqueloculina seminula* (LINNÉ, 1758) scale=0.8 mm
13. *Quinqueloculina* sp. 1. scale=0.95 mm
14. *Quinqueloculina* sp. 2. scale=1 mm
15. *Pyrgo bulloides* (D'ORBIGNY, 1826) scale=0.34 mm
16. *Articulina curta* LE CALVEZ, 1947 scale=0.45 mm
17. *Articulina laevigata* TERQUEM, 1882 scale=0.32 mm
18. *Articulina nitida* D'ORBIGNY, 1826 scale=0.18 mm
- 19–20. *Miliola prisca* (D'ORBIGNY, 1826) scale=0.85 mm
- 21–22. *Miliola strigillata* (D'ORBIGNY, 1850) scale=0.96 mm
23. *Massilina* sp. 1. scale=0.78 mm
- 24–25. *Spirolina mariei* LE CALVEZ, 1952 scale=0.42 mm
26. *Spirolina pedum* D'ORBIGNY, 1826 scale=1 mm
27. *Spirolina* sp. scale=0.45 mm




Plate 4

- 1–2. *Chrysalogonium tympanipectiformis* (SCHWAGER, 1866) scale=0.85 mm
3. *Chrysalogonium* sp. scale=0.45 mm
4. *Dentalina* cf. *acuta* D'ORBIGNY, 1846 scale=1 mm
5. *Dentalina approximata* REUSS, 1846 scale=1.3 mm
6. *Dentalina aboleta* SCHWAGER, 1865 scale=0.95 mm
- 7–8. *Dentalina cornicula* (D'ORBIGNY, 1826) scale=1 mm
9. *Dentalina debilis* HANTKEN, 1868 scale=1.1 mm
10. *Dentalina elegans* D'ORBIGNY, 1846 scale=1 mm
11. *Dentalina fissicostata* GÜMBEL, 1868 scale=0.95 mm
12. *Dentalina antennula* D'ORBIGNY, 1846 scale=1.2 mm
13. *Dentalina inornata* D'ORBIGNY, 1846 scale=1 mm
14. *Dentalina multilineata* BORNEMANN, 1855 scale=1 mm
15. *Dentalina karreri* (HANTKEN, 1868) scale=0.85 mm
16. *Dentalina havanensis* NEUGEBOREN, 1856 scale=1.2 mm
17. *Dentalina roemeri* NEUGEBOREN, 1856 scale=1 mm
18. *Dentalina subtilis* NEUGEBOREN, 1856 scale=1 mm
19. *Nodosaria affinis* REUSS, 1845 scale=1.1 mm
20. *Nodosaria badenensis* (D'ORBIGNY, 1846) scale=1.3 mm
21. *Nodosaria crassa* HANTKEN, 1868 scale=1 mm
22. *Dentalina budensis* HANTKEN, 1875 scale=1.1 mm
23. *Nodosaria elegans* (HANTKEN, 1875) scale=0.65 mm
24. *Nodosaria exilis* NEUGEBOREN, 1852 scale=1.1 mm
- 25–26. *Nodosaria acuminata* HANTKEN, 1875 scale=0.85 mm
27. *Nodosaria longiscata* D'ORBIGNY, 1846 scale=0.32 mm
28. *Nodosaria pyrula* (D'ORBIGNY, 1826) scale=0.75 mm
29. *Nodosaria radricula* LINNÉ, 1758 scale=1 mm
30. *Pseudonodosaria discreta* (REUSS, 1850) scale=1 mm
- 31–32. *Gonatosphaera inflata* BERMÚDEZ, 1949 scale=1.2 mm




Plate 5

1. *Frondicularia semicosta* KARRER, 1878 scale=1.2 mm
2. *Frondicularia budensis* (HANTKEN, 1875) scale=1.1 mm
- 3–4. *Frondicularia tenuissima* HANTKEN, 1875 scale=1.2 mm
- 5–6. *Lenticulina arcuatostrata* (HANTKEN, 1875) scale=1.3 mm
7. *Lenticulina excisa* (BORNEMANN, 1855) scale=1.85 mm
8. *Lenticulina falcifer* (STACHE, 1865) scale=0.76 mm
9. *Lenticulina granulata* (HANTKEN, 1875) scale=1.55 mm
10. *Lenticulina gutticastata* (GÜMBEL, 1870) scale=1.7 mm
11. *Lenticulina platyptera* (REUSS, 1870) scale=1.15 mm
12. *Lenticulina* sp. scale=1.1 mm
13. *Marginulinopsis porvaensis* (HANTKEN, 1875) scale=1.3 mm
14. *Marginulinopsis* sp. scale=1.3 mm
- 15–16. *Saracenaria hantkeni* CUSHMAN, 1933 scale=1.3 mm
17. *Marginulina behmi* (REUSS, 1866) scale=0.95 mm
18. *Marginulina fragaria texasensis* (CUSHMAN and APPLIN, 1926) scale=1.3 mm
- 19–20. *Marginulina hantkeni* BANDY, 1949 scale=1 mm
21. *Marginulina pediformis* BORNEMANN, 1855 scale=1.1 mm
22. *Marginulina propinqua* HANTKEN, 1883 scale=1.3 mm
23. *Marginulina tumida* REUSS, 1851 scale=1.2 mm
24. *Marginulina* sp. 1. scale=0.95 mm
25. *Vaginulinopsis hagni* n. sp., holotype (Inv. num.: M 2008.118.1) scale=0.55 mm
26. *Vaginulinopsis hagni* n. sp. (Inv. num.: M 2008.118.2) scale=0.85 mm
27. *Vaginulinopsis hagni* n. sp. (Inv. num.: M 2008.118.3) scale=0.8 mm
28. *Vaginulinopsis hantkeni* (HANTKEN, 1875) scale=1 mm
29. *Vaginulinopsis minimus* (HANTKEN, 1875) scale=1.1 mm




Plate 6

1. *Vaginulinopsis fragaria* (GÜMBEL, 1868) scale=1.2 mm
2. *Vaginulinopsis pseudodecorata* HAGN, 1956 scale=1.3 mm
3. *Vaginulinopsis* sp. scale=1.8 mm
- 4–5. *Planularia* sp. 1 scale=0.45 mm
6. *Planularia* sp. 2 scale=1.3 mm
- 7–8. *Vaginulina legumen* (LINNÉ, 1758) scale=1.2 mm
9. *Vaginulina* sp. cf. *V. ex gr. mexicana* NUTTALL, 1932 scale=1.3 mm
10. *Lagena globosa* (WALKER and BOYS, 1784) scale=0.85 mm
11. *Lagena hexagona* (WILLIAMSON, 1848) scale=0.15 mm
12. *Lagena laevis* (MONTAGU, 1803) scale=0.85 mm
13. *Lagena tenuis ornata* REUSS, 1863 scale=0.35 mm
14. *Lagena sulcata* (WALKER and JACOB, 1798) scale=0.25 mm
15. *Globulina guttula* REUSS, 1851 scale=0.86 mm
16. *Globulina minuta* (ROEMER, 1838) scale=0.54 mm
17. *Ramulina* sp. scale=0.32 mm
18. *Fissurina orbignyana* SEGUENZA, 1862 scale=0.95 mm
19. *Fissurina* sp. cf. *F. orbignyana praeclara* (CUSHMAN and RENZ, 1946) scale=1 mm
- 20–22. *Fissurina tricincta* (GÜMBEL, 1870) scale=0.85 mm
23. *Fissurina* sp. scale=0.85 mm
24. *Glandulina aequalis* REUSS, 1863 scale=0.35 mm
- 25–26. *Glandulina obtusissima* REUSS, 1863 scale=1.2 mm
27. *Glandulina hantkeni* (FRANZENAU, 1894) scale=0.85 mm
28. *Glandulina inflata* COSTA, 1853 scale=1 mm
29. *Glandulina* sp. scale=1.3 mm







Plate 7

1. *Bolivina cookei* CUSHMAN, 1922 scale=0.15 mm
- 2–3. *Bolivina elongata* HANTKEN, 1875 scale=0.1 mm
4. *Bolivina nobilis* HANTKEN, 1875 scale=0.23 mm
5. *Bolivina semistriata* HANTKEN, 1868 scale=0.32 mm
6. *Bolivina* sp. scale=0.56 mm
- 7–8. *Globocassidulina globosa* (HANTKEN, 1875) scale=0.75 mm
9. *Bulimina affinis* D'ORBIGNY, 1839 scale=0.43 mm
- 10–11. *Bulimina truncana* GÜMBEL, 1868 scale=0.5 mm
12. *Bulimina* sp. scale=0.6 mm
13. *Uvigerina chirana* CUSHMAN and STONE, 1947 scale=0.86 mm
14. *Uvigerina cocoaensis* CUSHMAN, 1925 scale=0.3 mm
15. *Uvigerina cocoaensis jacksonensis* CUSHMAN, 1925 scale=0.3 mm
16. *Uvigerina eocaena* GÜMBEL, 1868 scale=0.4 mm
- 17–18. *Uvigerina gallowayi* CUSHMAN, 1929 scale=0.5 mm
19. *Uvigerina hantkeni* CUSHMAN and EDWARDS, 1937 scale=0.15 mm
20. *Uvigerina multistriata* HANTKEN, 1871 scale=0.2 mm
- 21–22. *Uvigerina hourcqi* GRAHAM, DE KLASZ and RÉRAT, 1965 scale=0.4 mm
23. *Uvigerina pigmea* D'ORBIGNY, 1826 scale=0.15 mm
24. *Uvigerina rippensis* COLE, 1927 scale=0.35 mm
25. *Uvigerina tenuistriata* REUSS, 1870 scale=0.25 mm
26. *Reussella terquemi* CUSHMAN, 1945 scale=1 mm
27. *Reussella elongata* (TERQUEM, 1882) scale=0.65 mm
28. *Reussella elongata* (TERQUEM, 1882) scale=0.01 mm
- 29–30. *Reussella* sp. scale=0.6 mm



## Plate 8

- 1–2. *Fursenkoina hungarica* (HANTKEN, 1868) scale=0.1 mm
3. *Fursenkoina schreibersiana* (CZIZEK, 1848) scale=0.15 mm
4. *Ellipsoglandulina multicostata* (GALLOWAY and MORREY, 1929) scale=0.85 mm
5. *Nodosarella lorifera* (HALKYARD, 1919) scale=1.2 mm
6. *Nodosarella tuberosa* (GÜMBEL, 1868) scale=1.2 mm
7. *Nodosarella tuberosa* (GÜMBEL, 1868) scale=0.025 mm
8. *Nodosarella* sp. scale=1 mm
- 9–11. *Pleurostomella acuta* HANTKEN, 1875 scale=1.2 mm
12. *Pleurostomella alternans* SCHWAGER, 1866 scale=1.3 mm
13. *Pleurostomella eocaena* GÜMBEL, 1868 scale=1.3 mm
14. *Pleurostomella* sp. scale=1.4 mm
- 15–16. *Pleurostomella incrassata* HANTKEN, 1884 scale=1.4 mm
17. *Orthomorphina rohri* (CUSHMAN and STAINFORTH, 1945) scale=0.8 mm
18. *Stilostomella abyssorum* (BRADY, 1881) scale=1.2 mm
19. *Stilostomella abyssorum* (BRADY, 1881) scale=0.9 mm
20. *Stilostomella adolphina* (D'ORBIGNY, 1846) scale=1.2 mm
21. *Stilostomella adolphina* (D'ORBIGNY, 1846) scale=0.03 mm
22. *Stilostomella adolphina* (D'ORBIGNY, 1846) scale=1.1 mm
23. *Stilostomella consobrina* (D'ORBIGNY, 1846) scale=1.2 mm
24. *Stilostomella curvatura* (CUSHMAN, 1939) scale=1.6 mm
25. *Stilostomella elegans* (D'ORBIGNY, 1846) scale=1.7 mm
26. *Stilostomella elegans* (D'ORBIGNY, 1846) scale=0.05 mm
27. *Stilostomella emaciata* (REUSS, 1851) scale=1.2 mm
28. *Stilostomella hoernesii* (HANTKEN, 1868) scale=1 mm
29. *Stilostomella pauperata* (D'ORBIGNY, 1846) scale=1.3 mm
30. *Stilostomella* sp. scale=1.2 mm





Plate 9

- 1–2. *Cancris* sp. scale=0.95 mm
3. *Valvulineria* sp. scale=1 mm
- 4–5. *Eponides budensis* (HANTKEN, 1875) scale=1.1 mm
- 6–7. *Eponides haidingeri* (D'ORBIGNY, 1846) scale=1 mm
- 8–9. *Eponides umbonatus* (REUSS, 1851) scale=1.1 mm
10. *Eponides* sp. scale=0.6 mm
11. *Discorbis elegans* HANTKEN, 1875 scale=0.95 mm
- 12–13. *Discorbis perplexa* LE CALVEZ, 1949 scale=0.85 mm
- 14–16. *Cibicidoides eoceanus* (GÜMBEL, 1868) scale=1 mm
- 17–18. *Cibicidoides perlucidus* (NUTTALL, 1932) scale=1.1 mm
- 19–20. *Planulina austriaca* (D'ORBIGNY, 1826) scale=1.1 mm
21. *Planulina compressa* (HANTKEN, 1875) scale=1 mm
- 22–23. *Planulina costata* (HANTKEN, 1875) scale=0.35 mm
24. *Cibicides ammophilus* (GÜMBEL, 1870) scale=1 mm
- 25–27. *Cibicides boueanus* (D'ORBIGNY, 1846) scale=1.1 mm
- 28–29. *Cibicides carinatus* (TERQUEM, 1882) scale=0.95 mm



## Plate 10

1. *Cibicides mauricensis* HOWE and ROBERTS, 1939 scale=0.95 mm
2. *Cibicides oligocenicus* SAMOILOVA, 1947 scale=0.8 mm
3. *Cibicides proprius* BROTZEN, 1948 scale=1 mm
4. *Cibicides pseudolobatulus* PERELIS and REISS, 1975 scale=1 mm
5. *Cibicides pseudoungerianus* (CUSHMAN, 1922) scale=1.1 mm
- 6–7. *Cibicides sublobatulus* (GÜMBEL, 1868) scale=0.95 mm
8. *Cibicides sulzensis* (HERRMANN, 1917) scale=0.86 mm
9. *Cibicides ungerianus* D'ORBIGNY, 1846 scale=0.8 mm
- 10–12. *Cibicides westi* HOWE, 1939 scale=1.1 mm
13. *Cibicides* sp. scale=1 mm
- 14–15. *Dyocibicides uniserialis* Thalmann, 1933 scale=1.1 mm
16. *Asterigerina* sp. scale=0.95 mm
- 17–18. *Nuttallides* sp. scale=0.85 mm
19. *Nonion granosum* (D'ORBIGNY, 1846) scale=1 mm
20. *Nonion soldani* (D'ORBIGNY, 1846) scale=1.1 mm
21. *Nonionella* sp. scale=0.8 mm
- 22–23. *Melonis* sp. 1 scale=0.12 mm
- 24–25. *Pullenia jarvisi* CUSHMAN, 1936 scale=0.14 mm
26. *Pullenia quinqueloba* (REUSS, 1851) scale=0.23 mm
27. *Almaena* sp. scale=0.95 mm
- 28–30. *Queraltina epistominoides* MARIE, 1950 scale=1.1 mm






Plate 11

- 1–2. *Chilostomella tenuis* BORNEMANN, 1855 scale=0.45 mm
3. *Chilostomella* sp. scale=0.51 mm
- 4–5. *Anomalinoides affinis* (HANTKEN, 1875) scale=0.85 mm
6. *Anomalinoides alazanensis* (NUTTALL, 1932) scale=0.83 mm
- 7–8. *Anomalinoides grosserugosus* (GÜMBEL, 1868) scale=0.86 mm
- 9–10. *Anomalinoides* cf. *chilleana* (TODD and KNIKER, 1952) scale=0.78 mm
- 11–13. *Heterolepa dutemplei* (D'ORBIGNY, 1846) scale=1.1 mm
- 14–15. *Heterolepa simplex* FRANZENAU, 1884 scale=1.2 mm
16. *Gyroidinoides dissimilis* (CUSHMAN and RENZ, 1947) scale=1 mm
17. *Gyroidinoides* sp. scale=1.1 mm
18. *Hanzawaia producta* (TERQUEM, 1882) scale=1.2 mm
- 19–21. *Gyroidinoides soldanii* (D'ORBIGNY, 1826) scale=1.1 mm
- 22–23. *Gavelinella micra* (BERMÚDEZ, 1949) scale=1 mm
24. *Eoannularia eocenica* COLE and BERMÚDEZ, 1944 scale=0.85 mm
- 25–26. *Hanzawaia ammophila* (GÜMBEL, 1868) scale=1.1 mm
- 27–28. *Pararotalia inermis* (TERQUEM, 1882) scale=1.1 mm
29. *Rotalia* sp. cf. *R. calcar* (D'ORBIGNY, 1826) scale=1.1 mm
30. *Rotalia* sp. scale=0.8 mm
31. *Ammonia* sp. scale=0.46 mm
- 32–33. *Elphidium* sp. cf. *E. laeve* (D'ORBIGNY, 1826) scale=0.015 mm

